

THE ROYAL
SWEDISH
ACADEMY OF
SCIENCES



**INSTITUT
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89
info@mittag-leffler.se www.mittag-leffler.se

**The asymptotic behaviour in Schwarzschild
time of Vlasov matter in spherically
symmetric gravitational collapse**

H. Andréasson and G. Rein

REPORT No. 32, 2008/2009, fall

ISSN 1103-467X

ISRN IML-R- -32-08/09- -SE+fall

The asymptotic behaviour in Schwarzschild time of
Vlasov matter in spherically symmetric
gravitational collapse

Håkan Andréasson*
Mathematical Sciences
University of Gothenburg
Mathematical Sciences
Chalmers University of Technology
S-41296 Göteborg, Sweden
email: hand@chalmers.se

Gerhard Rein
Mathematisches Institut der Universität Bayreuth
D-95440 Bayreuth, Germany
email: gerhard.rein@uni-bayreuth.de

February 26, 2009

Abstract

Given a static Schwarzschild spacetime of ADM mass M , it is well-known that no ingoing causal geodesic starting in the outer domain $r > 2M$ will cross the event horizon $r = 2M$ in finite Schwarzschild time. In the present paper we show that in gravitational collapse of Vlasov matter this behaviour can be very different. We construct initial data for which a black hole forms and all matter crosses the event horizon as Schwarzschild time goes to infinity, and we show that this is a necessary condition for geodesic completeness of the event horizon. In addition to a careful analysis of the asymptotic behaviour of the matter characteristics our proof requires a new argument for global existence of solutions to the spherically symmetric Einstein-Vlasov system in an outer domain, since our initial data have non-compact support in the radial momentum variable and previous methods break down.

*Support by the Institut Mittag-Leffler (Djursholm, Sweden) is gratefully acknowledged.

1 Introduction

In a previous study [3] two classes of initial data for the spherically symmetric Einstein-Vlasov system were constructed which guarantee the formation of black holes. An additional argument to match the definition of a black hole in [8] is given [4]. In the present paper we denote any of these initial data classes by \mathcal{I} .

The analysis in [3] is carried out in Schwarzschild coordinates where the metric takes the form

$$ds^2 = -e^{2\mu(t,r)} dt^2 + e^{2\lambda(t,r)} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\varphi^2). \quad (1.1)$$

Here $t \in \mathbb{R}$ is the time coordinate, $r \in [0, \infty[$ is the area radius, i.e., $4\pi r^2$ is the area of the orbit of the symmetry group $\text{SO}(3)$ labeled by r , and the angles $\theta \in [0, \pi]$ and $\varphi \in [0, 2\pi]$ parameterize these orbits. The structure of the initial data \mathcal{I} is such that a possibly large fraction of its ADM mass M is necessarily located in the outer domain $r > 2M$. In [3] it was shown that solutions launched by such initial data have the following property: there exist constants $\alpha, \beta > 0$ such that spacetime is vacuum for

$$r \geq 2M + \alpha e^{-\beta t}, \quad t \geq 0. \quad (1.2)$$

Hence in this domain the metric equals the Schwarzschild metric

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\varphi^2),$$

representing a black hole of mass M . The generator of the event horizon approaches the surface $r = 2M$ asymptotically as Schwarzschild time goes to infinity, cf. [3, Thm. 2.4].

Although (1.2) gives information about the asymptotic location of the matter it does not answer the question whether or not matter crosses the surface $r = 2M$. As a matter of fact, the inequality (1.2) is not sufficient to conclude that any matter initially in the region $r > 2M$ ever crosses the surface $r = 2M$ since matter can pile up at $r = 2M$. On the other hand it is known that not all matter can cross the surface $r = 2M$ in finite Schwarzschild time. Indeed, if this were to happen the Einstein equations would imply that the metric function λ became infinite at $r = 2M$. But according to [14] this cannot happen for the solutions considered in [3]. It follows that on any finite time interval some matter must remain in the region $r > 2M$. The purpose of the present paper is to investigate the asymptotic behaviour of Vlasov matter in Schwarzschild time in the neighbourhood of

the event horizon. Note that if matter crosses the surface $r = 2M$ in finite time it also crosses the event horizon in finite time. Our main motivations are the following.

- In Proposition 2.3 it is shown that a necessary condition for completeness of the outgoing radial null geodesic which generates the event horizon is that all matter crosses the surface $r = 2M$ as Schwarzschild time goes to infinity.
- In a static Schwarzschild spacetime of ADM mass M no ingoing causal geodesic starting in the outer domain $r > 2M$ will cross the event horizon $r = 2M$ in finite Schwarzschild time. It is interesting to know if this remains true in evolutionary gravitational collapse. The result in the present paper shows that for the initial data we construct the behaviour is indeed very different.
- In [6, p. 13] some open problems about gravitational collapse are stated. For instance, for a scalar field it is known that

$$\sup_H r = 2 \sup_H m, \tag{1.3}$$

but for other matter models this issue is open. Here m is the quasi-local mass, H is the event horizon, and r the area radius.

- [9, Thm. 1.5] relates the asymptotic behaviour of the matter at the event horizon to the question of strong cosmic censorship, see also [9, Question 15.3].
- The asymptotic behaviour of matter in Schwarzschild time is directly related to what earth bound observers of gravitational collapse observe, which is not the case using other standard coordinates, e.g. Eddington-Finkelstein coordinates. If in Schwarzschild time all the matter crosses $r = 2M$, an earth bound observer will "see" all matter eventually swallowed by the emerging black hole.
- An important open problem is the question whether or not solutions can break down in finite Schwarzschild time. It is often conjectured that Schwarzschild coordinates are singularity avoiding and that in these coordinates solutions of the spherically symmetric Einstein-Vlasov system exist globally for general initial data. We expect that to understand the asymptotics of Vlasov matter in gravitational collapse is going to be useful for understanding the global existence issue in general.

In the present paper we construct a class \mathcal{J} of initial data for which the results in [3] apply, and such that all the matter asymptotically crosses the surface $r = 2M$. In particular (1.3) holds for Vlasov matter for this class of initial data. The class \mathcal{J} is different from the class \mathcal{I} in that the support of the momentum variables is not compact. This is a technical condition needed for our method of proof, but we believe that the conclusion holds for compactly supported data as well. Below, we always have in mind the radial momentum variable when we discuss compactly or non-compactly supported initial data. Our method of proof does imply that matter crosses the surface $r = 2M$ also in the compactly supported case, but we are not able to conclude that *all matter* eventually crosses $r = 2M$ in this case. We point out that the condition of non-compact support is required in some works in the cosmological case, cf. [8] and [16]. For compactly supported initial data the result in [14] guarantees that solutions exist as long as matter stays in a region $r \geq \epsilon > 0$. The proof in [14] breaks down for non-compactly supported data. But for applying the method in [3] it is crucial that solutions are global in an outer domain, and so we need to establish such a global existence result in the case of non-compactly supported data as well. Non-compactly supported data have been considered for other kinetic equations such as the Vlasov-Poisson system [11] and the Vlasov-Maxwell system [10, 15]. These methods do not directly apply in the case of the Einstein-Vlasov system, and we have not been able to find a result analogous to [14] for non-compactly supported data. However, the initial data set \mathcal{J} constructed below is such that the matter continues to move inward for all times. This crucial feature allows us to obtain the necessary global existence proof for the corresponding, non-compactly supported data in an outer domain.

The outline of the paper is as follows. In the next section we introduce the Einstein-Vlasov system, recall the set up and the construction of the class of initial data in [3], and formulate the main results of the present paper. Sections 4, 5, and 6 are devoted to their proofs.

2 Set up and main results

In this section we recall the Einstein-Vlasov system and the set up in [3] and formulate the main results. For more background on kinetic theory and the Einstein-Vlasov system we refer to [1]. We consider the asymptotically flat spherically symmetric Einstein-Vlasov system. We use Schwarzschild coordinates (t, r, θ, φ) and parameterize the metric as in (1.1). Asymptotic flatness means that the metric quantities λ and μ have to satisfy the bound-

ary conditions

$$\lim_{r \rightarrow \infty} \lambda(t, r) = \lim_{r \rightarrow \infty} \mu(t, r) = 0. \quad (2.1)$$

Vlasov matter is a collisionless ensemble of particles which is described by a density function f on phase space. In order to exploit the symmetry it is useful to introduce non-canonical variables on momentum space and write $f = f(t, r, w, L)$. The variables $w \in]-\infty, \infty[$ and $L \in [0, \infty[$ can be thought of as the radial component of the momentum and the square of the angular momentum respectively.

The Vlasov equation is given by

$$\partial_t f + e^{\mu-\lambda} \frac{w}{E} \partial_r f - \left(\lambda_t w + e^{\mu-\lambda} \mu_r E - e^{\mu-\lambda} \frac{L}{r^3 E} \right) \partial_w f = 0, \quad (2.2)$$

where

$$E = E(r, w, L) := \sqrt{1 + w^2 + L/r^2},$$

and where subscripts indicate partial derivatives. The Einstein equations read

$$e^{-2\lambda}(2r\lambda_r - 1) + 1 = 8\pi r^2 \rho, \quad (2.3)$$

$$e^{-2\lambda}(2r\mu_r + 1) - 1 = 8\pi r^2 p, \quad (2.4)$$

$$\lambda_t = -4\pi r j, \quad (2.5)$$

and the matter quantities are given by

$$\rho(t, r) = \frac{\pi}{r^2} \int_{-\infty}^{\infty} \int_0^{\infty} E f(t, r, w, L) dL dw, \quad (2.6)$$

$$p(t, r) = \frac{\pi}{r^2} \int_{-\infty}^{\infty} \int_0^{\infty} \frac{w^2}{E} f(t, r, w, L) dL dw, \quad (2.7)$$

$$j(t, r) = \frac{\pi}{r^2} \int_{-\infty}^{\infty} \int_0^{\infty} w f(t, r, w, L) dL dw. \quad (2.8)$$

The equations (2.2)–(2.8) constitute the spherically symmetric Einstein-Vlasov system in Schwarzschild coordinates. For a detailed derivation of this system we refer to [12].

As initial data we need to prescribe an initial distribution function $\mathring{f} = \mathring{f}(r, w, L) \geq 0$ such that

$$\int_0^r 4\pi \eta^2 \mathring{\rho}(\eta) d\eta = 4\pi^2 \int_0^r \int_{-\infty}^{\infty} \int_0^{\infty} E \mathring{f}(\eta, w, L) dL dw d\eta < \frac{r}{2}. \quad (2.9)$$

Here we denote by $\mathring{\rho}$ the energy density induced by the initial distribution function \mathring{f} . If in addition the initial data is C^1 we say that it is regular. In previous investigations the condition of compact support was included in the definition of regular data, but compact support in w is not required in the present paper and is replaced by a suitable fall-off condition, cf. (3.3) below. The Cauchy problem is well defined for regular initial data. We will restrict ourselves to a smaller class of regular initial data which guarantee the formation of black holes. Clearly, black holes do not form for any initial data, e.g., if the data is sufficiently small matter disperses and spacetime is geodesically complete, cf. [13].

Let us recall the set up and the properties of one of the initial data sets constructed in [3]. We fix $0 < r_0 < r_1$, and let γ^+ be the outgoing radial null geodesic originating from $r = r_0$, i.e.,

$$\frac{d\gamma^+}{ds}(s) = e^{(\mu-\lambda)(s,\gamma^+(s))}, \quad \gamma^+(0) = r_0. \quad (2.10)$$

We consider solutions of the spherically symmetric Einstein-Vlasov system (2.2)–(2.8) on the outer region

$$D := \{(t, r) \in [0, \infty[^2 \mid r \geq \gamma^+(t)\}. \quad (2.11)$$

Note that characteristics of the Vlasov equation can pass from the region D into the region $\{r < \gamma^+(t)\}$ but not the other way around so that initial data \mathring{f} posed for $r > r_0$ completely determine the solution on D .

Let $M := r_1/2$ be the total ADM mass and define the quasi-local mass by

$$m(t, r) = M - 4\pi \int_r^\infty \rho(t, \eta) \eta^2 d\eta. \quad (2.12)$$

Let $M_{\text{out}} < M$ be given and such that

$$\frac{2(M - M_{\text{out}})}{r_0} < \frac{8}{9}. \quad (2.13)$$

Take $R_1 > r_1$ such that

$$R_1 - r_1 < \frac{r_1 - r_0}{6},$$

and define

$$R_0 := \frac{1}{2}(r_1 + R_1).$$

We require that all the matter in the region $[r_0, \infty[$ is initially located in the strip $[R_0, R_1]$, with M_{out} being the corresponding fraction of the ADM mass

M , i.e.,

$$\int_{r_0}^{\infty} 4\pi r^2 \mathring{\rho}(r) dr = \int_{R_0}^{R_1} 4\pi r^2 \mathring{\rho}(r) dr = M_{\text{out}}.$$

Furthermore, the remaining fraction $M - M_{\text{out}}$ should be initially located within the ball of area radius r_0 , i.e.,

$$\int_0^{r_0} 4\pi r^2 \mathring{\rho}(r) dr = M - M_{\text{out}}.$$

If one considers the Einstein-Vlasov system on the whole spacetime $r \geq 0$, then the definition (2.12) for the quasi-local mass is equivalent to the more standard one, namely $m(t, r) = 4\pi \int_0^r \rho(t, \eta) \eta^2 d\eta$. This is because the ADM mass $M = m(t, \infty)$ is conserved. On the outer domain D the definition (2.12) is more suitable, since it does not refer to the matter inside $\{r < \gamma^+(t)\}$ except for the fact that this matter is there and contributes to the total mass.

The properties above concern the structure of the initial data in space. We also need to specify conditions on the momentum variables. Let $W_- < 0$ and $L_1 > 0$ be given. In [3] we introduced the **general support condition**: For all $(r, w, L) \in \text{supp } \mathring{f}$,

$$r \in]0, r_0] \cup [R_0, R_1],$$

and if $r \in [R_0, R_1]$ then

$$w \leq W_-, \quad 0 \leq L \leq L_1,$$

and

$$0 \leq L < \frac{3L}{\eta} \mathring{m}(\eta) + \eta \mathring{m}(\eta), \quad \eta \in [r_0, R_1].$$

Here we use the notation \mathring{m} when $\rho = \mathring{\rho}$ in (2.12). In addition to the conditions above the initial data \mathcal{I} in [3] were assumed to have compact support. If W_- is sufficiently negative, a black hole of ADM mass M forms and $\lim_{t \rightarrow \infty} \gamma^*(t) = 2M$ for a certain radially outgoing null geodesic which is the generator of the event horizon, cf. [3, Thm. 2.4] and [4, Sect. 4.3]. The initial data we construct below do have the properties specified above, but the support in the radial momentum variable w is not compact. However, the proof in [3] goes through unchanged also for such data *provided* the solutions are global on the domain D . With respect to global existence in the domain D the following holds.

Theorem 2.1 *Let regular initial data f be given with the properties specified above, and such that the fall-off condition (3.3) is satisfied. Then the corresponding solutions of the spherically symmetric Einstein-Vlasov system (2.2)–(2.8) in the domain D exist for all $t \geq 0$.*

We can now state the main result of the present paper.

Theorem 2.2 *There exists a class of regular initial data for the spherically symmetric Einstein-Vlasov system such that the corresponding solutions have the asymptotic property that*

$$\lim_{t \rightarrow \infty} m(t, 2M) = \lim_{t \rightarrow \infty} m(t, \gamma^*(t)) = M. \quad (2.14)$$

As mentioned in the introduction the condition (2.14) is a necessary condition for completeness of the generator γ^* of the event horizon. We state this in a proposition.

Proposition 2.3 *A necessary condition for future completeness of the generator γ^* of the event horizon is that (2.14) holds.*

3 Proof of Theorem 2.1

Compactly supported, regular initial data launch a local regular solution which can be extended as long as the momentum support of the solution can be controlled [12, 13]. We do not give a complete proof for the corresponding result for non-compactly supported data and restrict ourselves to establishing the main a-priori bounds. To this end it is convenient to introduce the Cartesian coordinates $x = r(\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) \in \mathbb{R}^3$ with corresponding momentum variable $v \in \mathbb{R}^3$ so that

$$w = \frac{x \cdot v}{r}, \quad L = |x \times v|^2, \quad |v|^2 = w^2 + \frac{L}{r^2}. \quad (3.1)$$

Here \cdot denotes the Euclidean scalar product and $|v|$ the induced norm. We denote by $(X, V)(s, t, x, v)$ the solution of the characteristic system of the Vlasov equation, written in the variables x and v ,

$$\begin{aligned} \dot{x} &= e^{(\mu-\lambda)(s,r)} \frac{v}{\sqrt{1+|v|^2}}, \\ \dot{v} &= - \left(\lambda_t(s, r) \frac{x \cdot v}{r} + e^{(\mu-\lambda)(s,x)} \mu_r(s, r) \sqrt{1+|v|^2} \right) \frac{x}{r}, \end{aligned}$$

with $(X, V)(t, t, x, v) = (x, v)$; here $\dot{\cdot}$ denotes the derivative with respect to s . We define

$$\begin{aligned} Q(t) &:= \sup \left\{ \frac{1 + |v|}{1 + |V(0, s, x, v)|} \mid 0 \leq s \leq t, (x, v) \in \text{supp } f(s) \right\} \\ &= \sup \left\{ \frac{1 + |V(s, 0, x, v)|}{1 + |v|} \mid 0 \leq s \leq t, (x, v) \in \text{supp } \mathring{f} \right\}. \end{aligned} \quad (3.2)$$

We require that the initial data satisfy the fall-off condition

$$\|\mathring{f}\| := \sup_{(x, v) \in \mathbb{R}^6} (1 + |v|)^5 |\mathring{f}(x, v)| < \infty. \quad (3.3)$$

Since

$$f(t, x, v) = \mathring{f}((X, V)(0, t, x, v)),$$

we get the estimate

$$f(t, x, v) \leq \|\mathring{f}\| (1 + |V(0, t, x, v)|)^{-5} \leq Q^5(t) \|\mathring{f}\| (1 + |v|)^{-5}.$$

Hence

$$\int_{\mathbb{R}^3} (1 + |v|) f(t, x, v) dv \leq \|\mathring{f}\| Q^5(t) \int_{\mathbb{R}^3} (1 + |v|)^{-4} dv \leq C \|\mathring{f}\| Q^5(t). \quad (3.4)$$

By the characteristic system,

$$\frac{d}{ds} (1 + |V(s, 0, x, v)|) \leq (\|e^{\mu-\lambda} \mu_r(s)\|_\infty + \|\lambda_t(s)\|_\infty) (1 + |V(s)|).$$

This implies that for $0 \leq s \leq t$,

$$\frac{1 + |V(s, 0, x, v)|}{1 + |v|} \leq e^{\int_0^t (\|e^{\mu-\lambda} \mu_r(\tau)\|_\infty + \|\lambda_t(\tau)\|_\infty) d\tau},$$

and we obtain the estimate

$$Q(t) \leq e^{\int_0^t (\|e^{\mu-\lambda} \mu_r(s)\|_\infty + \|\lambda_t(s)\|_\infty) ds}.$$

The field equations (2.3) and (2.4) together with the boundary condition (2.1) imply that

$$(\mu + \lambda)(t, r) = - \int_r^\infty (\mu_r + \lambda_r)(t, \eta) d\eta \leq 0,$$

and

$$e^{\mu-\lambda}\mu_r(t, r) = e^{\mu+\lambda} \left(\frac{m(t, r)}{r^2} + 4\pi r p(t, r) \right) \leq 4\pi r (\|\rho(t)\|_\infty + \|p(t)\|_\infty).$$

Together with (2.5) and (3.4) we have

$$\|e^{\mu-\lambda}\mu_r(s)\|_\infty + \|\lambda_t(s)\|_\infty \leq C(1+s)\|\mathring{f}\|Q^5(s),$$

so that

$$Q(t) \leq e^{\int_0^t C\|\mathring{f}\|(1+s)Q^5(s)ds}.$$

This implies that Q is bounded on some time interval $[0, T[$. A standard iterative procedure then shows that there is a local, regular solution which can be extended as long as the function Q does not blow up, cf. [12, 13].

Global existence in the outer domain D will now follow if we can establish a bound on $Q(t)$ in D . For this argument we use the variables (r, w, L) . By (3.3),

$$\mathring{f}(r, w, L) \leq C|w|^{-3}.$$

Let us define a quantity as in (3.2). By abuse of notation we let

$$\begin{aligned} Q(t) &:= \sup \left\{ \frac{|w|}{|W(0, s, r, w, L)|} \mid 0 \leq s \leq t, (r, w, L) \in \text{supp } f(s) \right\} \\ &= \sup \left\{ \frac{|W(s, 0, r, w, L)|}{|w|} \mid 0 \leq s \leq t, (r, w, L) \in \text{supp } \mathring{f} \right\}; \quad (3.5) \end{aligned}$$

notice that in the outer domain D the area radius $r \geq r_0 > 0$ so that by (3.1) a bound on Q as defined in (3.5) implies a bound on Q as defined in (3.2). The following lemma taken from [3] shows that when the general support condition holds, then the particles in the outer domain D keep moving inward in a controlled way.

Lemma 3.1 *Let \mathring{f} be regular and satisfy the general support condition for some suitable $W_- < 0$. Then for all characteristics $(R(t), W(t), L)$ with $(R(0), W(0), L) \in \text{supp } \mathring{f}$ and $R(0) \in [R_0, R_1]$,*

$$W(t) \leq e^{\lambda(t, R(t))} e^{-\lambda(0, R(0))} W(0) \leq e^{-\lambda(0, R(0))} W(0)$$

as long as $(t, R(t)) \in D$. In particular $w < 0$ for all $(r, w, L) \in \text{supp } f(t)$ and $(t, r) \in D$, and $j \leq 0$ on D .

Following [14] we find that along any characteristic in $\text{supp } f$,

$$\frac{d}{ds}w^2 \leq Cw^2 + C \int_{-\infty}^{\infty} \int_0^{L_1} |\tilde{w}| f(s, r, \tilde{w}, \tilde{L}) d\tilde{L} d\tilde{w}; \quad (3.6)$$

for this estimate it is essential that all particles are moving inward. We estimate the last term. Since

$$f(s, r, w, L) = \mathring{f}(R(0, s, r, w, L), W(0, s, r, w, L), L),$$

we have

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_0^{L_1} |\tilde{w}| f(s, r, \tilde{w}, \tilde{L}) d\tilde{L} d\tilde{w} \\ & \leq C \iint_{\text{supp } f(s, r, \cdot, \cdot)} |\tilde{w}| |W(0, s, r, \tilde{w}, \tilde{L})|^{-3} d\tilde{L} d\tilde{w}. \end{aligned}$$

By the definition of Q and the general support condition,

$$|W(0, s, r, \tilde{w}, \tilde{L})| \geq \max \left\{ |W_-|, \frac{|\tilde{w}|}{Q(s)} \right\}.$$

Hence we find that

$$\begin{aligned} \int_{-\infty}^{\infty} \int_0^{L_1} |\tilde{w}| f(s, r, \tilde{w}, \tilde{L}) d\tilde{L} d\tilde{w} & \leq C \int_0^{\infty} \int_0^{L_1} \tilde{w} \left(\max \left\{ |W_-|, \frac{\tilde{w}}{Q(s)} \right\} \right)^{-3} d\tilde{w} \\ & \leq C \int_0^{|W_-|Q(s)} \tilde{w} d\tilde{w} + \int_{|W_-|Q(s)}^{\infty} \tilde{w} \frac{Q^3(s)}{\tilde{w}^3} d\tilde{w} \\ & \leq C Q^2(s). \end{aligned}$$

We have thus derived the estimate

$$\frac{d}{ds}w^2 \leq Cw^2 + CQ^2(s),$$

and hence for $0 \leq s \leq t$,

$$\frac{w^2(s)}{w^2(0)} \leq e^{Ct} \left(1 + C \int_0^t Q^2(\tau) d\tau \right).$$

This implies that

$$Q^2(t) \leq e^{Ct} \left(1 + C \int_0^t Q^2(s) ds \right),$$

hence Q is bounded on bounded time intervals, and global existence in D follows. \square

4 Proof of Theorem 2.2

We aim to show that all characteristics starting in the domain $[R_0, R_1[$ enter the region $\{r \leq 2M\}$ in finite time, and we need an estimate for the required time. Let $(R(s), W(s), L)$ be a characteristic emanating from the support of \mathring{f} with $R(0) \in [R_0, R_1]$; all the following estimates are valid as long as $R(s) \geq 2M$. By Lemma 3.1 and the characteristic equation,

$$\dot{R}(s) = \frac{W(s)}{E(s)} e^{(\mu-\lambda)(s, R(s))} \leq -B(R(0), W(0)) e^{(\mu-\lambda)(s, R(s))},$$

where

$$B(r, w) := \frac{e^{-\lambda(0, r)} |w|}{\sqrt{1 + e^{-2\lambda(0, r)} w^2 + L_1 (2M)^{-2}}}.$$

We require that on $\text{supp } \mathring{f}$,

$$w \leq -e^{\lambda(0, r)} K(r) \tag{4.1}$$

where $K : [R_0, R_1[\rightarrow]0, \infty[$ is an increasing function which will be specified below. Hence

$$B(r, w) \geq \frac{K(r)}{\sqrt{1 + K^2(r) + L_1 (2M)^{-2}}} := B(r),$$

and

$$\dot{R}(s) \leq -B(R(0)) e^{(\mu-\lambda)(s, R(s))}. \tag{4.2}$$

By [3, Lemma 4.1 (a)],

$$\mu - \lambda \geq 2\hat{\mu}$$

where

$$\hat{\mu}(t, r) := - \int_r^\infty \frac{m(t, \eta)}{\eta^2} e^{2\lambda(t, \eta)} d\eta.$$

Inserting this into (4.2) implies that

$$\dot{R}(s) \leq -B(R(0)) e^{2\hat{\mu}(s, R(s))}. \tag{4.3}$$

In order to estimate the right hand side we compute $\hat{\mu}_t$, cf. [3, Lemma 4.1 (d)], and observe that

$$\begin{aligned} \hat{\mu}_t(s, r) &= \int_r^\infty 4\pi j(s, \eta) e^{(\mu+\lambda)(s, \eta)} e^{2\lambda(s, \eta)} d\eta \\ &\geq \frac{1}{2r} \int_r^\infty 4\pi \eta 2j(s, \eta) e^{(\mu+\lambda)(s, \eta)} e^{2\lambda(s, \eta)} d\eta; \end{aligned}$$

note that by Lemma 3.1, $j \leq 0$. Since

$$E + 2w + \frac{w^2}{E} = \left(\sqrt{E} + \frac{w}{\sqrt{E}} \right)^2 \geq 0$$

the expressions for the matter terms imply that $2j \geq -(\rho + p)$ so that by [3, Lemma 4.2],

$$\begin{aligned} \hat{\mu}_t(s, r) &\geq -\frac{1}{2r} \int_r^\infty 4\pi\eta(\rho + p)(s, \eta) e^{(\mu+\lambda)(s, \eta)} e^{2\lambda(s, \eta)} d\eta \\ &= -\frac{1}{2r} \left(1 - e^{(\mu+\lambda)(s, r)} \right). \end{aligned}$$

Moreover,

$$\hat{\mu}_r(s, r) = \frac{m(s, r)}{r^2} e^{2\lambda(s, r)}$$

and $|\dot{R}(s)| \leq e^{(\mu-\lambda)(s, R(s))}$. Hence

$$\begin{aligned} \hat{\mu}(t, R(t)) - \hat{\mu}(0, R(0)) &= \int_0^t \frac{d}{ds} \hat{\mu}(s, R(s)) ds \\ &= \int_0^t \left(\hat{\mu}_t(s, R(s)) + \hat{\mu}_r(s, R(s)) \dot{R}(s) \right) ds \\ &\geq \int_0^t \left(-\frac{1}{2R(s)} \left(1 - e^{(\mu+\lambda)(s, R(s))} \right) - \frac{m(s, R(s))}{R(s)^2} e^{(\mu+\lambda)(s, R(s))} \right) ds \\ &= \int_0^t \left(-\frac{1}{2R(s)} + \left(\frac{1}{2R(s)} - \frac{m(s, R(s))}{R(s)^2} \right) e^{(\mu+\lambda)(s, R(s))} \right) ds. \quad (4.4) \end{aligned}$$

The right hand side of this inequality will be estimated using the following lemma.

Lemma 4.1 *Let $\rho \in L^1([2M, R_1])$ be such that $\rho \geq 0$ and $0 \leq 2m(r)/r < 1$, where*

$$m(r) := M - \int_r^{R_1} 4\pi\eta^2 \rho(\eta) d\eta,$$

and let

$$e^{2\lambda(r)} := \left(1 - \frac{2m(r)}{r} \right)^{-1}, \quad r \in [2M, R_1].$$

Then for all $r \in [2M, R_1]$,

$$e^{-2 \int_r^{R_1} 4\pi\eta \rho(\eta) e^{2\lambda(\eta)} d\eta} \geq \frac{r - 2M}{r - 2m(r)}.$$

Proof. For $r \in [2M, R_1]$ we define

$$h(r) := (r - 2m(r)) e^{g(r)} - r + 2M, \quad g(r) := -2 \int_r^{R_1} 4\pi\eta\rho(\eta) e^{2\lambda(\eta)} d\eta.$$

Then

$$\begin{aligned} h'(r) &= (1 - 8\pi r^2 \rho(r)) e^{g(r)} + (r - 2m(r)) e^{g(r)} g'(r) - 1 \\ &= (1 - 8\pi r^2 \rho(r)) e^{g(r)} + r \left(1 - \frac{2m(r)}{r}\right) e^{g(r)} 8\pi r \rho(r) e^{2\lambda(r)} - 1 \\ &= (1 - 8\pi r^2 \rho(r)) e^{g(r)} + r e^{-2\lambda(r)} e^{g(r)} 8\pi r \rho(r) e^{2\lambda(r)} - 1 \\ &= e^{g(r)} - 1 \leq 0, \quad r \in [2M, R_1]. \end{aligned}$$

Hence for $r \in [2M, R_1]$,

$$h(r) \geq h(R_1) = 0,$$

which is the assertion. \square

Remark. It is interesting to note that the configuration for which equality holds in the inequality in the lemma can be shown to be an infinitely thin shell. This should be compared to the situation considered in [2] where an infinitely thin shell is the maximizer of a similar integral expression as above. Let us return to the proof of Theorem 2.2. By the field equations (2.3) and (2.4) and the form (2.6) and (2.7) of the matter terms,

$$\mu_r + \lambda_r = 4\pi r e^{2\lambda} (\rho + p) \leq 8\pi r e^{2\lambda} \rho.$$

Hence, Lemma 4.1 implies that for $r \in [2M, R_1]$,

$$e^{(\mu+\lambda)(s,r)} \geq e^{-2 \int_r^{R_1} 4\pi\eta\rho(s,\eta) e^{2\lambda(s,\eta)} d\eta} \geq \frac{r - 2M}{r - 2m(s,r)}.$$

We insert this into the estimate (4.4) and find that as long as $R(t) \geq 2M$,

$$\hat{\mu}(t, R(t)) \geq \hat{\mu}(0, R(0)) - \int_0^t \frac{M}{R^2(s)} ds.$$

By (4.3) this implies that

$$\dot{R}(s) \leq -B(R(0)) C(R(0)) \exp\left(-2 \int_0^s \frac{M}{R^2(\tau)} d\tau\right),$$

where $C(r) := e^{2\hat{\mu}(0,r)}$. This implies that

$$\begin{aligned} \frac{d}{ds} \frac{1}{R(s)} &= -\frac{\dot{R}(s)}{R^2(s)} \\ &\geq \frac{BC}{R^2(s)} \exp\left(-2 \int_0^s \frac{M}{R^2(\tau)} d\tau\right) = -\frac{BC}{2M} \frac{d}{ds} \exp\left(-2 \int_0^s \frac{M}{R^2(\tau)} d\tau\right) \end{aligned}$$

which upon integration yields the estimate

$$\begin{aligned} \frac{1}{R(t)} &\geq \frac{1}{R(0)} - \frac{BC}{2M} \left(\exp\left(-2 \int_0^t \frac{M}{R^2(s)} ds\right) - 1 \right) \\ &\geq \frac{1}{R(0)} + \frac{(BC)(R(0))}{2M} \left(1 - e^{-2Mt/R_1^2}\right). \end{aligned} \quad (4.5)$$

In order to proceed the functions C and B must be related properly. We require that

$$\dot{\rho}(r) > 0 \text{ for } r \in]R_0, R_1[$$

so that $\dot{m}(r) < M$ for $r \in [0, R_1[$, and

$$\begin{aligned} C(r) &= e^{2\hat{\mu}(0,r)} = \exp\left(-\int_r^\infty \frac{2\dot{m}(\eta)}{\eta^2(1-2\dot{m}(\eta)/\eta)} d\eta\right) \\ &> \exp\left(-\int_r^\infty \frac{2M}{\eta(\eta-2M)} d\eta\right) = 1 - \frac{2M}{r}. \end{aligned}$$

We can therefore choose the function K which specifies our support condition (4.1) in such a way that for $r \in [R_0, R_1[$,

$$B(r) C(r) = \frac{K(r)}{\sqrt{1+K^2(r)+L_1(2M)^{-2}}} C(r) > 1 - \frac{2M}{r}; \quad (4.6)$$

note that this necessarily implies that $K(r) \rightarrow \infty$ as $r \rightarrow R_1$, and that the function C is determined by the initial data. Given any $r^* \in]R_0, R_1[$ there now exists $\kappa > 0$ such that for $r \in [R_0, r^*]$,

$$B(r) C(r) > 1 - \frac{2M}{r} + \kappa.$$

The estimate (4.5) therefore implies that for any characteristic as above, but with $R(0) \in [R_0, r^*]$, and as long as $R(t) \geq 2M$,

$$\begin{aligned} \frac{1}{R(t)} &\geq \frac{1}{R(0)} + \frac{1}{2M} \left(1 - \frac{2M}{R(0)} + \kappa\right) \left(1 - e^{-2Mt/R_1^2}\right) \\ &= \frac{1+\kappa}{2M} \left(1 - e^{-2Mt/R_1^2}\right) + \frac{1}{R(0)} e^{-2Mt/R_1^2} \\ &> \frac{1+\kappa}{2M} \left(1 - e^{-2Mt/R_1^2}\right), \end{aligned}$$

which implies that

$$R(t) < \frac{2M}{1 + \kappa} \left(1 - e^{-2Mt/R_1^2}\right)^{-1}.$$

This shows that there is a time $t^* > 0$ such that $R(t^*) \leq 2M$ for all characteristics starting with $R(0) \in [R_0, r^*]$.

To complete the proof we fix $\epsilon > 0$ and let $r_\epsilon < R_1$ be sufficiently close to R_1 such that $\hat{m}(r_\epsilon) \geq M - \epsilon$. Then there is a finite time t_ϵ such that all characteristics $(R(t), W(t), L)$ with $R(0) \in [R_0, r_\epsilon]$ reach $r = 2M$ at some time $t \leq t_\epsilon$.

We construct a curve $(t, \alpha(t))$ with the property that $m(t, \alpha(t)) \geq M - 2\epsilon$ for $0 \leq t \leq t_\epsilon$ and $\alpha(t) = 2M$ for some $t \leq t_\epsilon$. To this end, let

$$\delta := \frac{\epsilon}{4\pi R_1^2 t_\epsilon}, \quad (4.7)$$

and let α be the solution of

$$\dot{\alpha} = \frac{j(s, \alpha) - \delta}{\rho(s, \alpha) + \delta} e^{(\mu-\lambda)(s, \alpha)}, \quad \alpha(0) = r_\epsilon. \quad (4.8)$$

The reason for introducing the δ parameter is to avoid any potential problems with uniqueness of solutions if $\rho = 0$. Taking the partial derivative of $e^{-2\lambda} = 1 - 2m/r$ with respect to t and using (2.5) we find that $m_t = -4\pi r^2 e^{\mu-\lambda} j$, and hence

$$\begin{aligned} \frac{d}{dt} m(t, \alpha(t)) &= -4\pi e^{(\mu-\lambda)(t, \alpha(t))} \alpha^2(t) j(t, \alpha(t)) + 4\pi \alpha^2(t) \rho(t, \alpha(t)) \dot{\alpha}(t) \\ &= 4\pi \alpha^2 e^{(\mu-\lambda)(t, \alpha(t))} \left(\frac{j - \delta}{\rho + \delta} \rho - j \right) \\ &\geq -4\pi \alpha^2(t) e^{(\mu-\lambda)(t, \alpha(t))} \delta \geq -\frac{\epsilon}{t_\epsilon}. \end{aligned}$$

Here we used that $j \leq 0$. Thus for all $0 \leq t \leq t_\epsilon$,

$$m(t, \alpha(t)) \geq M - 2\epsilon,$$

and it remains to show that $\alpha(t) \leq 2M$ for some $0 < t \leq t_\epsilon$. Define for each $r \in [R_0, R_1]$ the barrier curve $(t, R_B(t))$ by

$$\dot{R}_B = -B(r) e^{(\mu-\lambda)(s, R_B)}, \quad R_B(0) = r.$$

We use the term barrier curve since the area radius along this curve is larger than the area radius along any characteristic $(R(t), W(t), L)$ starting in the

support of \mathring{f} with $R(0) = r$. This is clear from the differential estimate (4.2) and the estimates which followed. In addition, all barrier curves starting at some $r \in [R_0, r_\epsilon]$ reach the region $r \leq 2M$ within the time interval $[0, t_\epsilon]$. For $r \in [R_0, R_1]$ the definition of $B(r)$ and the condition on the support of \mathring{f} imply that,

$$\begin{aligned} |j(0, r)| &\geq e^{\lambda(0, r)} B(r) \frac{\pi}{r^2} \int_{-\infty}^{\infty} \int_0^{\infty} \sqrt{1 + e^{-2\lambda(0, r)} w^2 + L_1(2M)^{-2}} \mathring{f} dL dw \\ &\geq B(r) \rho(0, r), \end{aligned}$$

and since $B < 1$ we thus have

$$\frac{|j(0, r)| + \delta}{\rho(0, r) + \delta} > B(r).$$

Consider the barrier curve $R_B(t)$ with $R_B(0) = r_\epsilon$. Then

$$\dot{\alpha}(0) = \frac{j(0, r_\epsilon) - \delta}{\rho(0, r_\epsilon) + \delta} e^{(\mu-\lambda)(0, r_\epsilon)} < -B(r_\epsilon) e^{(\mu-\lambda)(0, r_\epsilon)} = \dot{R}_B(0),$$

and hence $\alpha(s) < R_B(s)$ on a time interval $]0, s_1]$. Assume that $\alpha(t) > 2M$ for all $0 \leq t \leq t_\epsilon$. We define

$$t^* := \inf \{t \in [0, t_\epsilon] \mid R_B < \alpha \text{ on } [t, t_\epsilon] \text{ for all barriers starting in } [R_0, r_\epsilon]\}.$$

Then there exists some barrier curve $R_B(t)$ starting at some $r \in [R_0, r_\epsilon]$ such that $\alpha(t^*) = r^* := R_B(t^*)$. By definition of t^* , $\alpha(t) > R_B(t)$ for $t > t^*$, and hence

$$-\frac{|j(t^*, r^*)| + \delta}{\rho(t^*, r^*) + \delta} e^{(\mu-\lambda)(t^*, r^*)} = \dot{\alpha}(t^*) \geq \dot{R}_B(t^*) = -B(R_B(0)) e^{(\mu-\lambda)(t^*, r^*)},$$

which implies that

$$|j(t^*, r^*)| < B(R_B(0)) \rho(t^*, r^*).$$

The latter inequality is only possible if there is at least one characteristic $(R(t), W(t), L)$ with $R(t^*) = r^*$, and

$$\begin{aligned} \frac{K(R_B(0))}{\sqrt{1 + K^2(R_B(0)) + L_1(2M)^{-2}}} &= B(R_B(0)) > \frac{|W(t^*)|}{\sqrt{1 + |W(t^*)|^2 + LR(t^*)^{-2}}} \\ &\geq B(R(0), W(0)) \geq B(R(0)) = \frac{K(R(0))}{\sqrt{1 + K^2(R(0)) + L_1(2M)^{-2}}}. \end{aligned}$$

Since the function K is taken to be increasing this estimate implies that $R(0) < R_B(0) \leq r_\epsilon$. The barrier curve $(t, \tilde{R}_B(t))$ with $\tilde{R}_B(0) = R(0)$, must satisfy the estimate $\alpha(t^*) = r^* = R(t^*) < \tilde{R}_B(t^*)$, and this is a contradiction to the definition of t^* . Hence $\alpha(t) \leq 2M$ for some $0 < t \leq t_\epsilon$, which proves that $\lim_{t \rightarrow \infty} m(t, 2M) = M$.

We can chose the parameter δ in (4.7) such that $m(t, \alpha(t)) \geq M - 2\epsilon$ for all $0 \leq t \leq t_\epsilon + 1$. Since α is strictly decreasing, $\alpha(t_\epsilon + 1) < 2M$, and since $\lim_{t \rightarrow \infty} \gamma^*(t) = 2M$ there exists some time $t \geq t_\epsilon + 1$ such that $\gamma^*(t) \geq \alpha(t_\epsilon + 1)$ and hence $m(t, \gamma^*(t)) \geq M - 2\epsilon$. This completes the proof of Theorem 2.2. \square

To conclude our main result we show that initial data which satisfy the conditions required above do exist. To this end, let $\rho = \rho(r)$ be a C^1 function supported in $[R_0, R_1]$ with $\rho(r) > 0$ on $]R_0, R_1[$, and such that

$$M_{\text{out}} = 4\pi \int_{R_0}^{R_1} r^2 \rho(r) dr < M,$$

satisfies (2.13). Define

$$m(r) := M - 4\pi \int_r^\infty r^2 \rho(r) dr, \quad e^{-2\lambda(r)} := 1 - \frac{2m(r)}{r}.$$

If $r \in [R_0, R_1[$, then $M - m(r) > 0$, and hence there exists a function K as introduced in (4.1), which satisfies the condition (4.6). Now let $\tilde{h} = \tilde{h}(r, w, L)$ be a C^1 function supported in $[R_0, R_1] \times]-\infty, \infty[\times]0, L_1]$ and such that $\tilde{h}(r, w, L) = 0$ if $e^{-\lambda(r)} w > -K(r)$, and

$$\int_{-\infty}^\infty \int_0^{L_1} \tilde{h}(r, w, L) dL dw = \frac{r^2}{\pi}.$$

Let

$$h(r, w, L) = \frac{\tilde{h}(r, w, L)}{\sqrt{1 + w^2 + L/r^2}}$$

so that $\rho(r)h(r, w, L)$ induces the energy density ρ and the quasi-local mass m . Let f_i be a density function supported in $[0, r_0]$ such that the assumptions of Lemma 3.1 hold. Then $\mathring{f} = f_i + \rho h$ defines initial data which have all the required properties stated above.

5 Proof of Proposition 2.3

The function $m(t, r)$ is increasing in both variables. We assume that $m(t, r) \leq m(t, 2M) \leq C < M$ for all $t \geq 0$ and $r \leq 2M$ and have to

show that the generator γ^* of the event horizon is incomplete. According to [3, Thm. 2.4], γ^* approaches $r = 2M$ as $t \rightarrow \infty$. It follows that for all $t \geq 0$,

$$\lambda(t, \gamma^*(t)) \leq C. \quad (5.1)$$

Let $\tau \mapsto (t, r, \theta, \phi)(\tau)$ be an affine parameterization of γ^* with corresponding momenta $(p^0, p^1, p^2, p^3)(\tau)$. Since γ^* is radial, let $\theta = \pi/2$, $\phi = 0$ and $p^2 = p^3 = 0$. Since γ^* is null,

$$e^{2\mu}(p^0)^2 = e^{2\lambda}(p^1)^2,$$

and we get

$$p^0 = e^{\lambda-\mu}p^1. \quad (5.2)$$

By the geodesic equations,

$$\begin{aligned} \frac{dp^1}{d\tau} &= -e^{2(\mu-\lambda)}\mu_r(p^0)^2 - \lambda_r(p^1)^2 - 2\lambda_t p^0 p^1 \\ &= 4\pi\gamma^* e^{2\lambda}(p^1)^2[-p - \rho + 2j]. \end{aligned}$$

Here we used (5.2) to express p^0 in terms of p^1 . Since $dt/d\tau = p^0$,

$$\frac{dp^1}{dt} = 4\pi\gamma^* e^{\mu+\lambda}p^1[-(p + \rho) + 2j], \quad (5.3)$$

and γ^* is incomplete if

$$\int_0^\infty \frac{ds}{p^0} < \infty.$$

By (5.2) and (5.3),

$$\int_0^t \frac{ds}{p^0} = \int_0^t \frac{e^{\mu-\lambda}}{p^1} ds = \int_0^t e^{\mu-\lambda} \frac{1}{p^1(0)} e^{\int_0^s 4\pi\gamma^* e^{\mu+\lambda}(p+\rho-2j)d\eta} ds. \quad (5.4)$$

For the interior integral expression we have since $p \leq \rho$,

$$\int_0^s 4\pi\gamma^* e^{\mu+\lambda}(p + \rho - 2j)d\tau \leq \int_0^s 4\pi\gamma^* e^{\mu+\lambda}(2\rho - 2j)d\tau.$$

Along the null geodesic we get

$$\frac{d}{ds}\lambda(s, \gamma^*(s)) = \lambda_t + \lambda_r \frac{d\gamma^*}{ds} = 4\pi\gamma^*(\rho - j)e^{\mu+\lambda} - \frac{m}{(\gamma^*)^2}e^{\mu+\lambda}.$$

Hence by (5.1),

$$\begin{aligned} \int_0^s 4\pi\gamma^* e^{\mu+\lambda}(2\rho - 2j)d\tau &= 2\lambda(s, \gamma^*(s)) - 2\lambda(0, \gamma^*(0)) \\ &\quad + \int_0^s \frac{2m(\tau, \gamma^*(\tau))}{(\gamma^*(\tau))^2} e^{(\mu+\lambda)(\tau, \gamma^*(\tau))} d\tau \\ &\leq C + C \int_0^s e^{\mu(\tau, \gamma^*(\tau))} d\tau. \end{aligned}$$

By [3, Thm. 2.4] there exist positive constants α and β such that if (t, r) satisfies

$$r \geq 2M + \alpha e^{-\beta t} := \sigma(t),$$

then there is vacuum at (t, r) . By monotonicity of μ and the fact that there is vacuum for $r \geq \sigma$,

$$e^{\mu(\tau, \gamma^*(\tau))} \leq e^{\mu(\tau, \sigma(\tau))} \leq e^{-\int_{\sigma(\tau)}^{\infty} \frac{M}{\eta(\eta-2M)} d\eta} \leq C e^{-\frac{\beta\tau}{2}},$$

and thus

$$\int_0^s 4\pi\gamma^* e^{\mu+\lambda}(2\rho - 2j)d\tau \leq C.$$

From (5.4) we therefore obtain the estimate

$$\int_0^t \frac{ds}{p^0} \leq C \int_0^t e^{(\mu-\lambda)(s, \gamma^*(s))} ds = C \int_0^t \frac{d\gamma^*(s)}{ds} ds < 2CM,$$

which says that γ^* is incomplete, and the proof is complete. \square

Acknowledgement : The authors want to thank Helmut Friedrich whose question ‘‘Where is the matter?’’ was a starting point for this work, and Mihalis Dafermos for discussions.

References

- [1] H. ANDRÉASSON, The Einstein-Vlasov System/Kinetic Theory, *Living Rev. Relativity* **8** (2005).
- [2] H. ANDRÉASSON, Sharp bounds on $2m/r$ of general spherically symmetric static objects, *J. Differential Equations* **245**, 2243-2266 (2008).
- [3] H. ANDRÉASSON, M. KUNZE, G. REIN, The formation of black holes in spherically symmetric gravitational collapse, arXiv:0706:3787.

- [4] H. ANDRÉASSON, M. KUNZE, G. REIN, Gravitational collapse and the formation of black holes for the spherically symmetric Einstein-Vlasov system, *Quarterly of Appl. Math.*, to appear, arXiv:0812:1645.
- [5] D. CHRISTODOULOU, On the global initial value problem and the issue of singularities, *Class. Quantum Gravity* **16**, A23–A35 (1999).
- [6] M. DAFERMOS, Spherically symmetric spacetimes with a trapped surface, *Class. Quantum Gravity* **22**, 2221–2232 (2005).
- [7] M. DAFERMOS, A. D. RENDALL, An extension principle for the Einstein-Vlasov system in spherical symmetry, *Ann. Henri Poincaré* **6**, 1137–1155 (2005).
- [8] M. DAFERMOS, A. D. RENDALL, Strong cosmic censorship for T^2 -symmetric cosmological spacetimes with collisionless matter, arXiv:gr-qc/0610075.
- [9] M. DAFERMOS, A. D. RENDALL, Strong cosmic censorship for surface-symmetric cosmological spacetimes with collisionless matter, arXiv:gr-qc/0701034.
- [10] R. GLASSEY, W. STRAUSS, Large velocities in the relativistic Vlasov-Maxwell equations, *J. Fac. Sci. Univ. Tokyo Sect. IA Math* **36**, 615–627 (1989).
- [11] E. HORST, On the asymptotic growth of the solutions of the Vlasov-Poisson system, *Math. Meth. Appl. Sci.* **16**, 75–85 (1993).
- [12] G. REIN, *The Vlasov-Einstein System with Surface Symmetry*, Habilitationsschrift, München 1995.
- [13] G. REIN, A. D. RENDALL, Global existence of solutions of the spherically symmetric Vlasov-Einstein system with small initial data, *Comm. Math. Phys.* **150**, 561–583 (1992). Erratum: *Comm. Math. Phys.* **176**, 475–478 (1996).
- [14] G. REIN, A. D. RENDALL, J. SCHAEFFER, A regularity theorem for solutions of the spherically symmetric Vlasov-Einstein system, *Comm. Math. Phys.* **168**, 467–478 (1995).
- [15] J. SCHAEFFER, A small data theorem for collisionless plasma that includes high velocity particles, *Indiana University Math. J.* **53**, 1–34 (2004).

- [16] J. SMULEVICI, Strong cosmic censorship for T^2 -symmetric spacetimes with positive cosmological constant and matter. arXiv:0710:1351.