

THE ROYAL
SWEDISH
ACADEMY OF
SCIENCES



**INSTITUT
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89
info@mittag-leffler.se www.mittag-leffler.se

**On closed cosmological models that satisfy
the strong energy condition but do not
recollapse**

S. Calogero and M. Heinzle

REPORT No. 39, 2008/2009, fall

ISSN 1103-467X

ISRN IML-R- -39-08/09- -SE+fall

On closed cosmological models that satisfy the strong energy condition but do not recollapse

Simone Calogero^{1,*} and J. Mark Heinzle^{2,†}

¹*Departamento de Matemática Aplicada, Facultad de Ciencias, Universidad de Granada 18071 Granada, Spain*

²*Gravitational Physics, Faculty of Physics, University of Vienna, 1090 Vienna, Austria*

We show the existence of a rather general class of closed cosmological models of Bianchi type IX that do not exhibit recollapse but expand for all times. This is despite the fact that these models satisfy the strong energy condition by a wide margin.

PACS numbers:

In relativistic cosmology, the trichotomy of the Friedmann-Robertson-Walker (FRW) models is of prime importance. The spatial geometry determines the evolution of the cosmological model: In the hyperboloidal ($k = -1$) or the spatially flat ($k = 0$) case, the universe exhibits an initial singularity ('big bang'), and from that 'moment' on, the universe is forever expanding. In the case of a closed cosmological model ($k = +1$) we observe a fundamentally different behavior: "[...] the dynamical equations of general relativity show that the spatially closed 3-sphere universe will exist for only a finite span of time. [...] at a finite time after the big bang, the universe will achieve a maximum size [...], and then will begin to recontract. [...] a finite time after recontraction begins, a 'big crunch' will occur." The quotation is taken from [7]. The recollapse of closed FRW cosmologies holds because the *strong energy condition* is imposed on the matter (which is assumed to be a perfect fluid), i.e., $\rho + 3p \geq 0$, where ρ is the energy density and p the pressure.

On this basis it would be tempting to consider the recollapse of the spatially closed FRW cosmologies satisfying the strong energy condition as a paradigm for models with less symmetries. That this belief is erroneous has been demonstrated in [1]. At least in the case of cosmological models of Bianchi type IX, into which the closed FRW models are naturally embedded, there exists a rigorous result by Lin and Wald [5]: Assuming the dominant energy condition, i.e., $|p_i| \leq \rho$ for the anisotropic pressures p_1, p_2, p_3 , and non-negative average principal pressure p , i.e., $3p = (p_1 + p_2 + p_3) \geq 0$, then the 'closed-universe-recollapse conjecture' [1] holds. In the locally rotationally symmetric (LRS) case with isotropic matter, it is sufficient to require that $\rho + 3p \geq \epsilon\rho$ for an arbitrarily small $\epsilon > 0$, see [4]. (Note that if $p/\rho \rightarrow -1/3$, there exist models that expand forever approaching the Einstein static universe in the limit.)

In this letter, we approach the problem from a different direction by investigating cases where 'closed-universe-recollapse' does not hold. We prove that there exist cosmological models (of Bianchi type IX) satisfying the strong energy condition that do not recollapse but expand forever. Two points are important to emphasize:

(i) For these models the strong energy condition is satisfied 'by a wide margin'. The assumption we make is that $w = p/\rho$ is a constant, i.e., $w = \text{const} > -1/3$; however, $w < (1 - \sqrt{3})/3 \approx -0.244$, hence the average pressure is not positive. (ii) We prove the existence of a *generic* class of models that expand forever (where 'generic' is understood in the sense of an open set of initial data of the Einstein equations). An interesting observation is that these cosmological models exhibit partial (i.e., directional) accelerated expansion for late times.

As a matter of course, we do not propose the cosmological models we analyze as actual models of the universe. However, we want to emphasize that the matter model we consider is not 'exotic' but as physically realistic as possible. First, the energy conditions are satisfied and there do not form 'matter singularities' in the spacetime, and second, the properties of the matter source we consider resemble those of collisionless matter, elastic matter, and magnetic fields [3]. Restriction to matter sources of this type might modify the 'closed-universe-recollapse conjecture' in general, and, in view of the results of this paper, lead to specific bounds on the matter quantities. Finally let us comment on the approach we take. We use the dynamical systems approach to spatially homogeneous cosmologies [6]. However, as we will see, it is essential to avoid the standard Hubble-normalized variables—the results we present here are rather elusive in that approach.

Equations. For models of Bianchi type IX the metric can be written as

$$ds^2 = -dt^2 + g_{ij}(t) \hat{\omega}^i \hat{\omega}^j, \quad (1)$$

where $\{\hat{\omega}^1, \hat{\omega}^2, \hat{\omega}^3\}$ is a symmetry-adapted coframe satisfying $d\hat{\omega}^1 = -\hat{\omega}^2 \wedge \hat{\omega}^3$ (and cyclic permutations). The Einstein equations comprise evolution equations for the metric, $\partial_t g_{ij} = -2g_{il}k^l_j$, and for the extrinsic curvature k^i_j , see, e.g., [6]. The Gauss constraint reads ${}^3R + (\text{tr } k)^2 - k^i_j k^j_i = 2\rho$, where $\rho = -T^0_0$ is the energy density associated with the energy-momentum tensor T^μ_ν and 3R the scalar three-curvature. In the vacuum case or orthogonal perfect fluid case the metric is diagonal, i.e., $g_{ij}(t) = \text{diag}(g_{11}(t), g_{22}(t), g_{33}(t))$; in the locally rotationally symmetric (LRS) case we have $g_{22}(t) \equiv g_{33}(t)$.

We restrict ourselves to diagonal metrics even if the matter source is anisotropic.

In the diagonal case, an isotropic matter source is characterized by an energy-momentum tensor with $T^i_j = \text{diag}(p_1, p_2, p_3)$. The ‘isotropic pressure’ p is the average of the anisotropic pressures p_1, p_2, p_3 , i.e., $\text{tr} T = 3p$. Define w and w_i , $i = 1, 2, 3$, according to

$$p = w\rho, \quad p_i = w_i\rho; \quad (2)$$

obviously, $w_1 + w_2 + w_3 = 3w$. Matter that is consistent with LRS symmetry satisfies $w_2 = w_3$. For perfect fluids, $w_1 = w_2 = w_3 = w$, where w is typically assumed to be a constant. In this paper we consider anisotropic matter that resembles a perfect fluid in this way: We assume that the energy density and the isotropic pressure satisfy a linear equation of state, $w = \text{const}$, where the strong energy condition is supposed to hold, i.e., $w > -1/3$. The rescaled anisotropic pressures, w_1, w_2, w_3 , are assumed to be functions of (s_1, s_2, s_3) , i.e.,

$$w_k = w_k(s_1, s_2, s_3) \quad (k = 1, 2, 3), \quad (3)$$

where

$$s_k = g^{kk}(g^{11} + g^{22} + g^{33})^{-1}; \quad (4)$$

obviously, $s_1 + s_2 + s_3 = 1$. The functions w_k are such that there exists a distinguished (‘isotropic’) state (s_1^F, s_2^F, s_3^F) of the matter where $w_1 = w_2 = w_3 = w$, and remain bounded (and take limits) under extreme conditions (when one or more of the s_i are zero). In particular, there exists a constant v_- such that $w_1(0, s_2, s_3) = w_2(s_1, 0, s_3) = w_3(s_1, s_2, 0) = v_-$. There exist excellent examples for matter models of this type, e.g., collisionless matter, elastic matter, or magnetic fields; for a detailed discussion we refer to [2].

Let us define the Hubble scalar H as $H = -\text{tr} k/3$ and the traceless part of the extrinsic curvature, $(\sigma_1, \sigma_2, \sigma_3)$, by $k^i_i = -H - \sigma_i$ (no sum over i); $\sigma_1 + \sigma_2 + \sigma_3 = 0$. Furthermore, let $n_k = g_{kk}(\det g)^{-1/2}$; note that $n_k > 0$. Then the Einstein equations can be expressed as evolution equations for H , $(\sigma_1, \sigma_2, \sigma_3)$, and (n_1, n_2, n_3) plus one constraint, which can be used to express ρ in terms of the other variables, which leads to the fact that the matter enters the equations only via w and w_1, w_2, w_3 .

In the LRS case, which we will focus on henceforth, we observe $n_2 = n_3$ and $\sigma_2 = \sigma_3$ as well as $s_2 = s_3$; consistently, the matter satisfies $w_2 = w_3$. Let $\sigma = \sigma_2$ and let $s = s_2$; then

$$(\sigma_1, \sigma_2, \sigma_3) = (-2\sigma, \sigma, \sigma), \quad (s_1, s_2, s_3) = (1-2s, s, s). \quad (5)$$

Eq. (4) implies $s = (2 + n_2/n_1)^{-1}$, so that $s \in (0, 1/2)$. Finally we abbreviate the rescaled anisotropic pressure in the plane of rotational symmetry by u ; more specifically,

$$u(s) := w_2(1 - 2s, s, s). \quad (6)$$

The Einstein equations can be expressed in the variables H , σ , n_1 , and n_2 , and $u(s)$ appears in these equations. However, in the dynamical system approach to cosmology these equations are rather expressed in a normalized version of these variables. We follow an approach that uses a well adapted normalization, see [4]: We define the ‘dominant variable’ D by

$$D = \sqrt{H^2 + \frac{n_1 n_2}{3}}, \quad (7a)$$

and we introduce normalized variables according to

$$H_D = \frac{H}{D}, \quad \Sigma_D = \frac{\sigma}{D}, \quad N_{1D} = \frac{n_1}{D}, \quad N_{2D} = \frac{n_2}{D}. \quad (7b)$$

Note that these variables are not independent because

$$1 - H_D^2 = \frac{N_{1D} N_{2D}}{3}. \quad (8)$$

Further, we define a normalized density $\Omega_D = \rho/(3D^2)$, and we replace the cosmological time t by a rescaled time variable τ through

$$\frac{d}{d\tau} = \frac{1}{D} \frac{d}{dt}. \quad (9)$$

Henceforth, a prime denotes differentiation w.r.t. τ . Expressed in these variables the Einstein evolution equations split into a decoupled equation for D ,

$$D' = -D \left(H_D(1 + q_D) + \Sigma_D(1 - H_D^2) \right), \quad (10)$$

and a system of coupled equations for the normalized variables (7b),

$$H'_D = -(1 - H_D^2)(q_D - H_D \Sigma_D), \quad (11a)$$

$$\Sigma'_D = -(2 - q_D)H_D \Sigma_D + (1 - H_D^2)\Sigma_D^2 + \frac{1}{3}(N_{1D}^2 - N_{1D}N_{2D}) + 3\Omega_D(u(s) - w), \quad (11b)$$

$$N'_{1D} = N_{1D}(q_D H_D - 4\Sigma_D + (1 - H_D^2)\Sigma_D), \quad (11c)$$

$$N'_{2D} = N_{2D}(q_D H_D + 2\Sigma_D + (1 - H_D^2)\Sigma_D), \quad (11d)$$

where q_D is given by $q_D = 2\Sigma_D^2 + (1/2)(1 + 3w)\Omega_D$. The normalized matter density, Ω_D , can be expressed in terms of the variables (7b) by the Gauss constraint, which reads

$$\Sigma_D^2 + \frac{1}{12}N_{1D}^2 - \frac{1}{3}N_{1D}N_{2D} + \Omega_D + \cancel{(1 - H_D^2)} = 1; \quad (12)$$

the 3rd and the 5th term cancel because of (8).

Remark. An important remark is the following: Equation (11a) implies that H_D is decreasing if $H_D = 0$, i.e., $H'_D|_{H_D=0} = -q_D < 0$; therefore, a cosmological model with $H_D(\tau_0) > 0$ at some time τ_0 satisfies $H_D(\tau) > 0 \forall \tau \leq \tau_0$. Consequently, by proving the existence of models that satisfy $H_D(\tau) > 0$ for all sufficiently large τ , we prove the existence of models with $H_D > 0$ and thus positive expansion H for all times.

Remark. More common than (7) is the Hubble-normalized approach, where the variables are defined as in (7b), replacing D by H . Accordingly, $H_D \equiv 1$ and the system of equations is given by (10) and (11b)–(11d) with (12), where H_D is set to 1. Although the resulting equations are simpler than (17), we do not have a choice; we will see that the Hubble-normalized approach necessarily fails to uncover our results.

The quantity s appears as the argument of the function $u(s)$ in (11b); it is determined in terms of the variables N_{1D} and N_{2D} by

$$s = \left(2 + \frac{N_{2D}}{N_{1D}}\right)^{-1}. \quad (13)$$

The evolution equation for s is

$$s' = -12s\left(\frac{1}{2} - s\right)\Sigma_D. \quad (11e)$$

We are free to add this equation to the system (11).

Of the five equations (11a)–(11e) three are independent, because there exist two constraints between the variables: (8) and (13). Our aim is to use this system to construct a three-dimensional dynamical system with advantageous properties. A good start is to simply discard the equation for N_{2D} , keeping the equation for s . The remaining constraint,

$$\frac{1 - H_D^2}{N_{1D}^2} = \frac{2}{3}s^{-1}\left(\frac{1}{2} - s\right), \quad (14)$$

reveals that s uniquely determines the ratio of $1 - H_D^2$ and N_{1D}^2 . Hence, in order to reconstruct $1 - H_D^2$ and N_{1D}^2 we use s in combination with a ‘radial’ variable r ,

$$r = \sqrt{(1 - H_D^2)^2 + (N_{1D}^2)^2}. \quad (15)$$

Obviously, by restricting ourselves to $H_D > 0$ we obtain a one-to-one transformation whose inverse reads

$$H_D = H_D(r, s) = \sqrt{1 - r \frac{1 - 2s}{\sqrt{1 - 4s + 13s^2}}}, \quad (16a)$$

$$N_{1D} = N_{1D}(r, s) = \sqrt{\frac{3rs}{\sqrt{1 - 4s + 13s^2}}}. \quad (16b)$$

This is a well-defined and regular map on the preimage of the set $\mathbb{R}^+ \times \mathbb{R}^+$. (We refrain from going into details in this paper, since we merely use that (16) is well-behaved for sufficiently small r ; however, we may refer to [3].)

Finally, we express the Einstein evolution equations in the variables D and r, s, Σ_D . The equation for D is (10),

the system for r, s, Σ_D reads

$$r' = r \left(2H_D(q_D - H_D\Sigma_D) - \frac{54\Sigma_D s^2}{1 - 4s + 13s^2}\right), \quad (17a)$$

$$\Sigma_D' = -(2 - q_D)H_D\Sigma_D - (1 - H_D^2)(1 - \Sigma_D^2) + \frac{1}{3}N_{1D}^2 + 3\Omega_D(u(s) - w), \quad (17b)$$

$$s' = -12s\left(\frac{1}{2} - s\right)\Sigma_D, \quad (17c)$$

where $q_D = 2\Sigma_D^2 + \frac{1}{2}(1 + 3w)\Omega_D$, and H_D and N_{1D} are functions of r and s , see (16). The Gauss constraint reads

$$\Sigma_D^2 + \frac{1}{12}N_{1D}^2 + \Omega_D = 1; \quad (18)$$

it is used to solve for Ω_D .

The dynamical system (17) completely describes the dynamics of locally rotationally symmetric cosmological models of Bianchi type IX in their expanding phase ($H > 0$). In other words, each solution of (17) yields an LRS Bianchi type IX model in its expanding phase, and conversely, the expanding phase of each model is represented by a solution of (17). The main advantage of the system (17) over other representations of the Einstein equations lies in its extendibility: The system (17) possesses a regular extension to its boundaries $\Sigma_D = \pm 1$, $s = 0$, $s = \frac{1}{2}$, and $r = 0$. For details we refer to [3].

Anisotropic matter. The function $u(s)$ in (17b) encodes the properties of the matter model. For isotropic matter we have $u(s) \equiv w$; for anisotropic matter, $u(s)$ represents the rescaled anisotropic pressure in the plane of local rotational symmetry, cf. (6); recall that $s \in [0, 1/2]$. From $w_2(s_1, 0, s_3) = v_-$ we obtain $u(0) = v_-$, cf. (5) and (6). The value of u at $s = 1/2$ is not independent; to see this recall first that $w_1 + 2w_2 = 3w$; now, $s = 1/2$ corresponds to $(s_1, s_2, s_3) = (0, 1/2, 1/2)$, so $w_1(0, 1/2, 1/2) + 2w_2(0, 1/2, 1/2) = 3w$; hence, $w_1(0, s_2, s_3) = v_-$ results in $v_- + 2u(1/2) = 3w$. Summarizing,

$$u(0) = v_-, \quad u(1/2) = w + (1/2)(w - v_-). \quad (19)$$

For reasonable matter models like collisionless matter, elastic matter, or magnetic fields, $u(s)$ is a monotone function on $[0, 1/2]$ interpolating between the values (19); we refer to [2] for examples. It is often beneficial to use the anisotropy parameter β instead of the constant v_- ; it is defined as

$$\beta = \frac{2(w - v_-)}{1 - w}. \quad (20)$$

Finally, let us state the assumptions on the matter model we consider in this paper. We assume that

$$-\frac{1}{3} < w < \frac{1 - \sqrt{3}}{3} \approx -0.244, \quad (21)$$




FIG. 1: The admissible values of the rescaled isotropic pressure w and v_{\pm} , $v_{\pm} = 3w - 2v_{\mp}$, which represent (the extremes of) the rescaled anisotropic pressures in the plane of symmetry and orthogonal to it, respectively. Both the strong and the dominant energy condition are satisfied. $(1 - \sqrt{3})/3 \approx -0.24$.

hence the strong energy condition is satisfied. Furthermore, v_{-} is assumed to satisfy

$$\frac{1}{6} \left(1 + 6w - \sqrt{-3 + (1 - 3w)^2} \right) < v_{-} \quad (22)$$

$$v_{-} < \frac{1}{6} \left(1 + 6w + \sqrt{-3 + (1 - 3w)^2} \right).$$

The admissible domain of the parameters w and v_{\pm} is depicted in Fig. 1. Clearly, the dominant energy condition is satisfied since $|v_{\pm}| < 1$.

Results. Since the dynamical system (17) extends regularly to $r = 0$ it is suggestive to analyze the system induced on that surface. Setting $r = 0$ in (17) is straightforward. From (16) we obtain $H_D|_{r=0} = 1$, $N_{1D}^2|_{r=0} = 0$, so that (18) becomes $\Omega_D = 1 - \Sigma_D^2$. This in turn implies $q_D = \frac{1}{2}(1 + 3w) + \frac{3}{2}(1 - w)\Sigma_D^2$. Insertion into (17) yields

$$\Sigma_D' = -3(1 - \Sigma_D^2) \left(\frac{1}{2}(1 - w)\Sigma_D - (u(s) - w) \right), \quad (23a)$$

$$s' = -12\Sigma_D s \left(\frac{1}{2} - s \right). \quad (23b)$$

The state space for this two-dimensional dynamical system is $[-1, 1] \times [0, 1/2]$. The dynamical systems analysis is straightforward. We use that $u(s)$ is a function such that $u(0) = v_{-}$ and $u(1/2) = w + (w - v_{-})/2$; here, w and v_{-} are assumed to satisfy (21) and (22), respectively.

We focus our attention on the fixed point R of (23), which is given by

$$R: r = 0, s = 0, \Sigma_D = -\beta. \quad (24)$$

It is straightforward to prove that R is a sink for the flow of the system (23), because

$$s^{-1}s'|_R = 6\beta < 0, \quad (25a)$$

$$(\Sigma_D + \beta)^{-1}(\Sigma_D + \beta)'|_R = -\frac{3}{2}(1 - \beta^2)(1 - w) < 0,$$

hence the eigenvalues of the linearization of (23) at R are negative.

The crucial property of the fixed point R is revealed by considering the full system (17): R is a sink not only on the boundary $r = 0$, but also for the full system (17). To see this we simply compute

$$r^{-1}r'|_R = 2 \left(\frac{3}{2}(1 - w)\beta^2 + \frac{1}{2}(1 + 3w) + \beta \right) < 0 \quad (25c)$$

and use (22) to establish that the r.h. side is negative.

Since the fixed point R is a sink for the flow of the system (17), i.e., for LRS Bianchi type IX models with anisotropic matter satisfying (21) and (22), there exists an open subset of LRS type IX initial data such that the corresponding solutions converge to R. Because $H_D = 1$ at R we find that H is positive for these solutions for late times; by the remark following (12) we obtain that H is positive, i.e., these models are expanding, for all times.

In the following we analyze the asymptotic behavior of these forever expanding LRS type IX solutions in terms of the metric variables: For a solution of (17) that converges to the point R we find $r \rightarrow 0$, $s \rightarrow 0$, $\Sigma_D \rightarrow -\beta$, hence $H_D \rightarrow 1$, $N_{1D} \rightarrow 0$ by (16). Furthermore, $N_{2D} \rightarrow \infty$; to see this we first note that $s \rightarrow 0$ implies $s \sim N_{1D}N_{2D}^{-1}$ by (13). Then, from $N_{1D} \sim \sqrt{rs}$, see (16b), we conclude that $N_{2D} \sim \sqrt{r/s}$. Using that $s^{-1}s|_R = 6\beta$ and $r^{-1}r|_R = 2(q + \beta)$, cf. (25a), where

$$q := q_D|_R = \frac{3}{2}(1 - w)\beta^2 + \frac{1}{2}(1 + 3w),$$

we finally get $N_{2D}^{-1}N_{2D}|_R = q - 2\beta > 0$, and the claim follows. Note that the fact that $N_{2D} \rightarrow \infty$ is unproblematic here, but makes the treatment of the problem extremely difficult in the standard Hubble-normalized approach. For large τ , the shear variables are $\sigma_1 = -2\sigma = -2D\Sigma_D \simeq 2D\beta$ and $\sigma_2 = \sigma_3 = \sigma = D\Sigma_D \simeq -D\beta$. To obtain the metric we integrate $\partial_t g_{kk} = 2g_{kk}(H + \sigma_k)$ (which is a consequence of $\partial_t g_{ij} = -2g_{il}k^l_j$). In the first step we note that $H = H_D D = D$ so that the equation reads $\partial_t g_{kk} = 2g_{kk}(D + \sigma_k)$, i.e.,

$$\partial_\tau g_{11} = 2g_{11}(1 + 2\beta), \quad \partial_\tau g_{22} = 2g_{22}(1 - \beta). \quad (26)$$

Second, we integrate $D^{-1}D'|_R = -(1 + q)$ and (9) to get $\tau = (1 + q)^{-1} \log(t - t_0) + \tau_0$ with constants τ_0 and t_0 . Therefore, (26) leads to an asymptotic behavior of the metric represented by

$$g_{11} \propto (t - t_0)^{\frac{2(1+2\beta)}{1+q}}, \quad g_{22} \propto (t - t_0)^{\frac{2(1-\beta)}{1+q}} \quad (27)$$

as $t \rightarrow \infty$. Note that β is negative, but $1 + 2\beta$ is positive, which is a consequence of (22). An interesting observation is the occurrence of partial (directional) accelerated expansion. A straightforward calculation reveals that $(1 - \beta)/(1 + q) > 1$, hence lengths in the plane of

local rotational symmetry expand at an accelerated rate. The maximal rate of acceleration is obtained by maximizing $(1 - \beta)/(1 + q)$ over the domain depicted in Fig. 1; this yields a maximal value of $(1 - \beta)/(1 + q) \approx 1.112$, is attained for w close to $-1/3$.

We conclude by restating the main result: The behavior (27) is generic, i.e., there exists an open set of LRS type IX initial data such that the associated solutions of the Einstein equations with anisotropic matter behave as (27) as $t \rightarrow \infty$ and thus expand forever. (These solutions do not behave extraordinarily in other respects; for instance, there exist forever expanding solutions that isotropize toward the singularity.) In this sense, eternal expansion is as likely as recollapse in the case of LRS Bianchi type IX with anisotropic matter that satisfies the conditions (21) and (22), and thus in particular the strong energy condition.

Acknowledgments. We gratefully acknowledge the hospitality of the Mittag-Leffler Institute.

[†] Electronic address: Mark.Heinzle@univie.ac.at

- [1] J.D. Barrow, G.J. Galloway, and F.J. Tipler. The closed-universe recollapse conjecture. *Mon. Not. R. Astron. Soc.*, 223 : 835–844, 1986.
- [2] S. Calogero and J.M. Heinzle. Dynamics of Bianchi type I solutions of the Einstein equations with anisotropic matter. *Ann. Henri Poincaré*, 2009. to appear.
- [3] S. Calogero and J.M. Heinzle. Dynamics of locally rotationally symmetric and spatially homogeneous cosmological models with anisotropic matter. Preprint, 2009.
- [4] J.M. Heinzle, N. Röhr, and C. Uggla. Matter and dynamics in closed cosmologies. *Phys. Rev. D*, 71 : 083506, 2005.
- [5] X.f. Lin and R.M. Wald. Proof of the closed-universe-recollapse conjecture for diagonal Bianchi type IX cosmologies. *Phys. Rev. D*, 40 : 3280–3286, 1989.
- [6] J. Wainwright and G.F.R. Ellis, editors. *Dynamical systems in cosmology*. Cambridge University Press, Cambridge, 1997.
- [7] R.M. Wald. *General Relativity*. The University of Chicago Press, 1984.

* Electronic address: calogero@ugr.es