

THE ROYAL
SWEDISH
ACADEMY OF
SCIENCES



**INSTITUT
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89
info@mittag-leffler.se www.mittag-leffler.se

On Milliken-Taylor Ultrafilters

H. Mildenberger

REPORT No. 28, 2009/2010, fall

ISSN 1103-467X

ISRN IML-R- -28-09/10- -SE+fall

ON MILLIKEN-TAYLOR ULTRAFILTERS

HEIKE MILDENBERGER

ABSTRACT. We show that there may be a Milliken-Taylor ultrafilter with infinitely many near coherence classes of ultrafilters in its projection to ω , answering a question by López-Abad. We show that k -coloured Milliken-Taylor ultrafilters have at least $k + 1$ near coherence classes of ultrafilters in its projection to ω . We show that the Mathias forcing with a Milliken-Taylor ultrafilters destroys all Milliken-Taylor ultrafilters from the ground model.

1. MILLIKEN-TAYLOR ULTRAFILTERS AND THEIR PROJECTIONS

We answer a question of López-Abad whether there can be more than two near coherence classes of ultrafilters in the core of a Milliken-Taylor ultrafilter. Then we show that in Milliken Taylor ultrafilter with k colours there are $k + 1$ near coherence classes in its projection to ω , generalising a result of Blass [7]. Then we investigate whether a Milliken-Taylor ultrafilter is preserved by forcing with another Milliken-Taylor ultrafilter. The somewhat surprising answer is no, independently of the distance of the two ultrafilters.

In the rest of this introductory section we review part of the relevant background.

Our nomenclature follows [10] and [5]. We let \mathbb{F} be the collection of all *non-empty* finite subsets of ω . For $a, b \in \mathbb{F}$ we write $a < b$ if $(\forall n \in a)(\forall m \in b)(n < m)$. We will work with proper filters on \mathbb{F} , i.e. non-empty subsets of $\mathcal{P}(\mathbb{F})$ that are closed under binary intersections and supersets and do not contain the empty set. A filter on \mathbb{F} is called non-principal if it does contain all sets of the form $\mathbb{F} \setminus E$, E finite. A sequence $\bar{c} = \langle c_n : n \in \omega \rangle$ of members of \mathbb{F} is called *unmeshed* if for all n , $c_n < c_{n+1}$. Henceforth, barred lower case variables stand for such sequences. For $n \leq \omega$, the set $(\mathbb{F})^n$ denotes the collection of all unmeshed sequences in \mathbb{F} of length n . If \bar{c} is a sequence in $(\mathbb{F})^\omega$, we write $(\text{FU})^\omega(\bar{c})$ for the set of all unmeshed sequences whose members are finite unions of some of the c_n 's and we write $\text{FU}(\bar{c})$ for the set of all finite unions of members of \bar{c} .

Date: March 5, 2010.

2010 Mathematics Subject Classification: 03E05, 03E17, 03E35.

Key words and phrases: Milliken-Taylor ultrafilters, P -points, near coherence classes of ultrafilters, forcing, preserving ultrafilters.

The author is entirely supported by Marie Curie grant PIEF-2008-219292 of the European Union.

Definition 1.1. Given \bar{c} and \bar{d} in $(\mathbb{F})^\omega$, we say that \bar{d} is a condensation or a block-subsequence of \bar{c} and we write $\bar{d} \sqsubseteq \bar{c}$ if $\bar{d} \in (\text{FU})^\omega(\bar{c})$. We say \bar{d} is almost a condensation of \bar{c} and we write $\bar{d} \sqsubseteq^* \bar{c}$ iff there is an n such that $\langle d_t : t \geq n \rangle$ is a condensation of \bar{c} .

Definition 1.2. A non-principal filter \mathcal{F} on \mathbb{F} is said to be an union filter if it has a basis of sets of the form $\text{FU}(D)$ for $D \subseteq \mathbb{F}$. A non-principal filter \mathcal{F} on \mathbb{F} is said to be an ordered-union filter if it has a basis of sets of the form $\text{FU}(\bar{d})$ for $\bar{d} \in (\mathbb{F})^\omega$. Let μ be an uncountable regular cardinal. An ordered-union filter is said to be μ -stable if, whenever it contains $\text{FU}(\bar{d}_\alpha)$ for $\bar{d}_\alpha \in (\mathbb{F})^\omega$, $\alpha < \mu$, then it also contains some $\text{FU}(\bar{e})$ for some \bar{e} that is almost a condensation of \bar{d}_α for $\alpha < \mu$. $< \kappa$ -stable means μ -stable for all $\mu < \kappa$. For “ \aleph_0 -stable” we say “stable”. Stable ordered-union ultrafilters are also called Milliken-Taylor ultrafilters.

Ordered-union ultrafilters need not exist, as their existence implies the existence of Q -points [5, Prop. 3.9], namely the minimum and the maximum projections (see Def. 1.8) are Q -points. An ultrafilter \mathcal{V} on ω is called a Q -point or a rare ultrafilter, if given a strictly increasing sequence π_n there is $A \in \mathcal{V}$ such that for all n , $|A \cap [\pi_n, \pi_{n+1})| = 1$. There are models without Q -points e.g., the Laver model [18] and all models of NCF [8]. NCF implies that any filter is nearly coherent to a filter with $\mathfrak{u} < \mathfrak{d}$ generators and this filter cannot be rapid. Rapidness says that the set of the enumerating functions of the members of the ultrafilter is a dominating family. Since rapidness is preserved under finite-to-one functions, under NCF there is no rapid filter. Since all Q points are rapid, there is no Q -point under NCF. With the help of Hindman’s theorem one shows that under $\text{MA}(\sigma\text{-centred})$ stable (even $< 2^\omega$ -stable) ordered-union ultrafilters exist [5]. We recall Hindman’s theorem:

Theorem 1.3. (Hindman, [13, Corollary 3.3]) *If the set \mathbb{F} is partitioned into finitely many pieces then there is a set $\bar{d} \in (\mathbb{F})^\omega$ such that $\text{FU}(\bar{d})$ is included in one piece.*

Indeed, for constructing a forcing $\mathbb{M}(\mathcal{U})$ we use only Hindman’s theorem. However, for analysing its behaviour we also derive from the following finite-dimensional version:

Theorem 1.4. (Milliken [19] and Taylor [21]) *If the set $(\mathbb{F})^n$ is partitioned into finitely many pieces then there is a set $\bar{d} \in (\mathbb{F})^\omega$ such that $(\text{FU}(\bar{d}))^n$ is included in one piece.*

As a motivation for the question we answer in this paper we consider also the following theorem for colourings with arbitrary many colours:

Theorem 1.5. *The canonical partition theorem. (Taylor [21]) If f is a function defined on \mathbb{F} then there is a set $\bar{d} \in (\mathbb{F})^\omega$ such that one of the following five statements holds for all $s, t \in \text{FU}(\bar{d})$.*

- (1) $f(s) = f(t)$

- (2) $f(s) = f(t)$ iff $\min(s) = \min(t)$,
- (3) $f(s) = f(t)$ iff $\max(s) = \max(t)$,
- (4) $f(s) = f(t)$ iff $(\min(s), \max(s)) = (\min(t), \max(t))$
- (5) $f(s) = f(t)$ iff $s = t$.

Instead of $(\mathbb{F})^\omega$ smaller domains for the colourings and for finding the homogeneous set can be considered:

Definition 1.6. *A ordered union ultrafilter \mathcal{U} on \mathbb{F} is said to have the Ramsey property, if for any $\bar{c} \in \mathcal{U}$ the set $(\text{FU}(\bar{c}))^n$ is partitioned into finitely many pieces then there is a set $\bar{d} \in \mathcal{U}$ such that $(\text{FU}(\bar{d}))^n$ is included in one piece.*

A ordered union ultrafilter \mathcal{U} on \mathbb{F} is said to have the canonical partition property if for any $\bar{c} \in \mathcal{U}$ for any function defined on $\text{FU}(\bar{c})$ there is $\bar{d} \in \mathcal{U}$, such that $f \upharpoonright \text{FU}(\bar{d})$ is canonised as above.

Now Milliken-Taylor ultrafilters are the reservoirs for the homogeneous sets from the previous theorems by the following theorem:

Theorem 1.7. *(Blass [5, Theorem 4.2 (a) to (c)]) For an ordered union ultrafilter \mathcal{U} the following are equivalent:*

- (a) *Stability,*
- (b) *the canonical partition property,*
- (c) *the Ramsey property.*

The following notions relate Milliken-Taylor ultrafilters to ultrafilters on ω

Definition 1.8. *Let \mathcal{U} be a filter on \mathbb{F} .*

- (1) *The core of \mathcal{U} is the filter $\Phi(\mathcal{U})$ such that*

$$X \in \Phi(\mathcal{U}) \text{ iff } (\exists \text{FU}(\bar{c}) \in \mathcal{U}) \left(\bigcup_{n \in \omega} c_n \subseteq X \right).$$

- (2) *The minimum projection of \mathcal{U} is the filter $\min(\mathcal{U})$ such that*

$$X \in \min(\mathcal{U}) \text{ iff } (\exists \text{FU}(\bar{c}) \in \mathcal{U}) (\{\min(c_n) : n \in \omega\} \subseteq X).$$

- (3) *Analogously we define the maximum projection of \mathcal{U} , $\max(\mathcal{U})$.*

So it follows from their definitions that $\min(\mathcal{U})$ and $\max(\mathcal{U})$ are in $[\Phi(\mathcal{U})] = \{\mathcal{V} \in \beta\omega : \mathcal{V} \supseteq \Phi(\mathcal{U})\}$.

We recall near coherence classes. For $B \subseteq \omega$ and $h: \omega \rightarrow \omega$, we let $h''B = \{h(b) : b \in B\}$ and $h^{-1}B = \{n : h(n) \in B\}$. By a *filter* we mean a proper filter on ω . We call a filter *non-principal* if it contains all cofinite sets. Let \mathcal{F} be a non-principal filter on ω and let $h: \omega \rightarrow \omega$ be finite-to-one (that means that the preimage of each natural number is finite). Then also $h(\mathcal{F}) = \{X : h^{-1}X \in \mathcal{F}\}$ is a non-principal filter. It is the filter generated

by $\{h''X : X \in \mathcal{F}\}$. Two filters \mathcal{F} and \mathcal{G} are *nearly coherent*, if there is some finite-to-one $h: \omega \rightarrow \omega$ such that $h(\mathcal{F}) \cup h(\mathcal{G})$ generates a filter. On the set $\beta\omega^*$ of non-principal ultrafilters on ω , the near coherence relation is an equivalence relation whose classes are called near coherence classes. Models with just one near coherence classes are known [8, 9, 17], and a model with just two classes is in preparation by Blass and Shelah. Apropos the possible infinite numbers of near coherence classes, Banach and Blass [1] showed: If there are infinitely many near coherence classes, then there are 2^c classes. Under CH or $\mathfrak{u} \geq \mathfrak{d}$, there are $2^{\mathfrak{d}}$ classes, and this and similar forms are early results on the spectrum of possible numbers of near coherence classes [4], for more history, see [1].

This paper is concerned with the following question. Let \mathcal{U} be a Milliken-Taylor ultrafilter. How many near coherence classes of ultrafilters are in $[\Phi(\mathcal{U})]$? For the wider class of union ultrafilters there is a recent result by Blass:

Theorem 1.9. [7, Theorem 38] *Let \mathcal{U} be a union ultrafilter. Then $\min(\mathcal{U})$ and $\max(\mathcal{U})$ are non-nearly coherent P -points.*

We remark that Blass [5] showed that the minimum and the maximum projections of Milliken-Taylor ultrafilters are selective ultrafilters (also called Ramsey ultrafilters), i.e., P -points and Q -points.

In the light of the canonical partition property Theorem 1.7 for \mathcal{U} it may appear difficult to find more near coherence classes beyond the class of $\min(\mathcal{U})$ and $\max(\mathcal{U})$. So are there more near coherence classes in $[\Phi(\mathcal{U})]$? We will see that canonisation for functions does not mean canonisation for ultrafilters on ω , after all an ultrafilter \mathcal{V} amounts to $\chi(\mathcal{V})$ many functions.

We recall some definitions: We say “ A is almost a subset of B ” and write $A \subseteq^* B$ iff $A \setminus B$ is finite. Similarly, the symbol $=^*$ denotes equality up to finitely many exceptions in $[\omega]^\omega$ or in ω^ω .

Let κ be a regular cardinal. An ultrafilter \mathcal{U} is called a P_κ -point if for every $\gamma < \kappa$, for every $A_i \in \mathcal{U}$, $i < \gamma$, there is some $A \in \mathcal{U}$ such that for all $i < \gamma$, $A \subseteq^* A_i$; such an A is called a *pseudo-intersection* or a *diagonalisation* of the A_i , $i < \gamma$. A P_{\aleph_1} -point is just called P -point. A notion of forcing \mathbb{P} preserves an ultrafilter \mathcal{U} iff $\Vdash_{\mathbb{P}} “(\forall X \in [\omega]^\omega)(\exists Y \in \mathcal{U})(Y \subseteq X \vee Y \subseteq \omega \setminus X)”$ and in the contrary case we say “ \mathbb{P} destroys \mathcal{U} ”. If \mathbb{P} is proper and preserves \mathcal{U} and \mathcal{U} is a P -point, then \mathcal{U} stays a P -point [8, Lemma 3.2].

Let \mathcal{F} be a filter. $\mathcal{B} \subseteq [\omega]^\omega$ is a *pseudobase* for \mathcal{F} if for every $X \in \mathcal{F}$ there is some $Y \in \mathcal{B}$ such that $Y \subseteq X$. A pseudobase $\mathcal{B} \subseteq \mathcal{F}$ is called a *base*. The character/ π -character of \mathcal{F} , $\chi(\mathcal{F})/ \pi\chi(\mathcal{F})$, is the smallest cardinality of a base/ pseudobase of \mathcal{F} . The ultrafilter characteristic, \mathfrak{u} , is the smallest character of a non-principal ultrafilter.

A set $M \subseteq {}^\omega\omega$ is called a *meagre set* if it is a countable union of nowhere dense sets. The covering number for the ideal \mathcal{M} of meagre sets, $\text{cov}(\mathcal{M})$, is the smallest number of meagre sets that cover together the real line.

In Section 2 we give two types of construction, one based on $\text{MA}(\sigma\text{-centred})$ and one on $\text{cov}(\mathcal{M}) = \mathfrak{c}$, of a Milliken-Taylor ultrafilter with infinitely many near coherence classes of ultrafilters in its projection. In Section 3 we generalise Blass' result Theorem 1.9 to $k + 1$ classes. In Section 4 we show that Milliken-Taylor ultrafilters are not preserved under any forcing with another or the same Milliken-Taylor ultrafilter.

2. A MILLIKEN-TAYLOR ULTRAFILTER WITH INFINITELY MANY PAIRWISE NON-NEARLY COHERENT ULTRAFILTERS IN ITS CORE

It might be possible that there are just two near coherence classes in the universe (Blass and Shelah have work in preparation towards this) and that there is a Milliken-Taylor ultrafilter such that just the minimum and the maximum projection are representatives of these two classes (this is much more vague). We do not know to exclude this. However, we know that the contrary is consistent:

Theorem 2.1. *Let κ be a regular cardinal. Under $\text{MA}(\sigma\text{-centred})$ and $\mathfrak{c} = \kappa$ there is a \mathfrak{c} -stable Milliken-Taylor ultrafilter with $2^{\mathfrak{c}}$ near coherence classes of ultrafilters represented in $[\Phi(\mathcal{U})]$.*

Remark: We carry out the construction only for countably many near coherence classes represented by \mathcal{V}^n , $n \in \omega$. We can get \mathfrak{c} classes in $[\Phi(\mathcal{U})]$ by an easy modification of the construction. In order to get the maximal number $2^{\mathfrak{c}}$ near coherence classes represented in $[\Phi(\mathcal{U})]$ we use that we construct the \mathcal{V}^n such that the sequence $\langle \mathcal{V}^n : n < \omega \rangle$ is discrete, that is for every n , there is $A \in \mathcal{V}^n$, $A \notin \mathcal{V}^m$ for $m \neq n$. This is automatically fulfilled in our construction. Then Banach's and Blass' technique [1] to generate $2^{\mathfrak{c}}$ classes in the closure of $\{\mathcal{V}^n : n \in \omega\}$ in $\beta\omega \setminus \omega$ works. A self-contained proof would mean to repeat good parts of their work, and therefore we refer the reader to [1].

For the proof we need the following definition:

Definition 2.2. *For $\bar{c} = (\mathbb{F})^{\leq \omega}$ and $A \subseteq \omega$ with $A \cap \text{set}(\bar{c}) \neq \emptyset$, we define $\bar{c} \upharpoonright A = \langle c_k \cap A : k \in \omega, A \cap c_k \neq \emptyset \rangle$ and $(\bar{c}; \text{past } n) = \langle c_k : c_k \cap [n, \infty) \neq \emptyset \rangle$. For $\bar{c} \in (\mathbb{F})^{\leq \omega}$ we let $\text{set}(\bar{c}) = \bigcup \{c_k : k \in |\bar{c}|\}$. The number of blocks in \bar{c} is denoted by $|\bar{c}|$. Unmeshed sets of blocks $\{c_n : n \in \omega\}$ are identified with their increasing enumerations $\bar{c} = \langle c_n : n \in \omega \rangle$. We do this also for finite numbers instead of ω .*

Proof. Let B_ε , $\varepsilon < \kappa$, $\varepsilon \equiv 0 \pmod{3}$ enumerate $\mathcal{P}(\mathbb{F})$, Y_ε , $\varepsilon < \kappa$, $\varepsilon \equiv 1 \pmod{3}$ enumerate $\mathcal{P}(\omega)$ and let f_ε , $\varepsilon < \kappa$, $\varepsilon \equiv 2 \pmod{3}$, enumerate $\omega^{\omega, \text{fto}}$. We modify the usual construction of a Milliken-Taylor ultrafilter by having three kinds of successor steps: Hinman steps, ultrafilter steps, and "making non-nearly coherent" steps. Also in the limit steps, we have to be careful to take a somewhat fat almost condensation. By induction on $\varepsilon < \mathfrak{c}$ we choose $\bar{c}_\varepsilon \in (\mathbb{F})^\omega$ and X_ε^n , $n < \omega$, with the following rules:

- (1) $\bar{c}_0 = \langle \{k\} : k < \omega \rangle$, $\bar{c}_\varepsilon \sqsubseteq^* \bar{c}_\delta$ for $\delta < \varepsilon < \mathfrak{c}$, we write $\bar{c}_\varepsilon = \langle c_{\varepsilon,k} : k < \omega \rangle$.
- (2) $X_\varepsilon^n \subseteq \text{set}(\bar{c}_\varepsilon)$ for $n < \omega$.
- (3) $X_\varepsilon^n \subseteq^* X_\delta^n$ for $\delta < \varepsilon < \bar{c}$ for every $n < \omega$.
- (4) If $\varepsilon = 0 \pmod 3$ then we let $\bar{c}'_{\varepsilon+1}$ be gotten by merging block from \bar{c}_ε and not dropping anything such that for each $k \in \omega$, $\bar{c}'_{\varepsilon+1,k}$ contains an element of $X_{\varepsilon+1}^i$ for all $i \leq k$. Then we take Hindman's theorem to get $\bar{c}_{\varepsilon+1} \sqsubseteq \bar{c}'_{\varepsilon+1}$ such that $\text{FU}(\bar{c}_{\varepsilon+1}) \subseteq B_\varepsilon$ or $\subseteq \mathbb{F} \setminus B_\varepsilon$, and still $\text{set}(\bar{c}_{\varepsilon+1}) \cap X_\varepsilon^n$ is infinite for all n , let $X_{\varepsilon+1}^n = X_\varepsilon^n \cap \text{set}(\bar{c}_{\varepsilon+1})$.
- (5) If $\varepsilon = 1 \pmod 3$ we choose $X_{\varepsilon+1}^n \subseteq Y_\varepsilon \cap X_\varepsilon^n$ or $\subseteq (\omega \setminus Y_\varepsilon) \cap X_\varepsilon^n$. We let $\bar{c}_{\varepsilon+1} = \bar{c}_\varepsilon$.
- (6) If $\varepsilon = 2 \pmod 3$ we choose $X_{\varepsilon+1}^n \subseteq X_\varepsilon^n$ such that for all $n \neq m$, $f_\varepsilon'' X_{\varepsilon+1}^n \cap f_\varepsilon'' X_{\varepsilon+1}^m = \emptyset$. We let $\bar{c}_{\varepsilon+1} = \bar{c}_\varepsilon$.
- (7) In the limit steps, we take parallel almost condensations and pseudointersection: Let $\varepsilon = \lim_{k \rightarrow \omega} \varepsilon_k$, ε_k , $k \in \omega$ strictly increasing, $\varepsilon_k = 0 \pmod 3$. Take $n(k) > k, n(k-1)$ so that $(\bar{c}_{\varepsilon_k} ; \text{past } n(k)) \sqsubseteq \bar{c}_{\varepsilon_{k+1}}$ and such that $\bar{c}_{\varepsilon_k, n(k)}$ contains an element of each of $X_{\varepsilon_k}^j$, $j \leq k$. Then let $\bar{c}_\varepsilon = \bar{c}_{\varepsilon_0} \upharpoonright [0, n(0)) \frown \bar{c}_{\varepsilon_1} \upharpoonright [n(0), n(1)) \dots$. Here, \frown means concatenation. Let $X_\varepsilon^n = \bigcup_{i < \omega} X_{\varepsilon_i}^n \cap [n(i-1), n(i))$. Then for every n , for every $k \geq n$ we took a block into the condensation that contains points of $X_{\varepsilon_k}^n \cap [n(k-1), n(k))$, so $X_\varepsilon^n \cap \text{set}(\bar{c}_\varepsilon)$ is infinite. In the general of a limit $\varepsilon < \mathfrak{c}$ we choose $n(k)$, $k \in \omega$ with the help of a σ -centred forcing as follows $\mathbb{P} = \{(c, F) : c \in (\mathbb{F})^{<\omega}, F \subseteq \varepsilon \text{ finite}\}$ with $(c, F) \leq_{\mathbb{P}} (d, F')$ iff $c \sqsubseteq d$, $F \subseteq F'$ and $d \upharpoonright (\max(\text{set}(c), \max(\text{set}(d))) \sqsubseteq \bar{c}_\zeta$ for all $\zeta \in F$ and $(\text{set}(d) \setminus \text{set}(c)) \cap X_\zeta^n \neq \emptyset$ for $n \leq |d|$ and $\zeta \in F$. By $\text{MA}(\sigma\text{-centred})$, there is a generic filter G and there is a generic real $\bar{c}_\varepsilon = \bigcup \{s : (s, F) \in G\}$.

Now let $\mathcal{V}^n = \{X \in [\omega]^\omega : \exists \varepsilon < \mathfrak{c}, X \supseteq X_\varepsilon^n\}$. Let $\mathcal{U} = \{B \subseteq \mathbb{F} : \exists \varepsilon < \mathfrak{c}, \text{FU}(\bar{c}_\varepsilon) \subseteq B\}$. This is Milliken-Taylor ultrafilter. \mathcal{V}^n is not nearly coherent to \mathcal{V}^m for $m \neq n$ and $\mathcal{V}^n \in \Phi(\mathcal{U})$. \dashv

Now we improve the theorem from $\text{MA}(\sigma\text{-centered})$ (which is equivalent to $\mathfrak{p} = \mathfrak{c}$ [3]) to the weaker hypothesis $\text{cov}(\mathcal{M}) = \mathfrak{c}$. However, there is one price: We get less stability. There will be an inductive construction of length \mathfrak{c} again, however, it is not \sqsubseteq^* -descending anymore. We call an arbitrary initial segment of the construction $(\mathcal{F}, (\mathcal{G}^n)_n)$ and do not bother about indexing the stage. By enumerating all descending ω -sequences and adding almost condensations to them to the filter we ensure stability.

Theorem 2.3. *Let $\text{cov}(\mathcal{M}) = \mathfrak{c}$. Then there is a Milliken-Taylor ultrafilter with $2^{\mathfrak{c}}$ near coherence classes of ultrafilters represented in $[\Phi(\mathcal{U})]$.*

Proof. We will do an inductive construction along \mathfrak{c} .

On the way we need a form of “generic existence” over initial segments of the construction. There will be steps $\alpha < \mathfrak{c}$ where we have to take an almost condensation of an ω -sequence \bar{d}^n of members of the first component \mathcal{F} , an ordered-union filter, of the initial segment $(\mathcal{F}, (\mathcal{G}^n)_n)$ of the construction. We will use a Cohen real (which we have by $\text{cov}(\mathcal{M}) > |\alpha|$) to find a fat enough almost condensation so that none of our initial segments to the countably many pairwise non-nearly coherent ultrafilter will get lost. In addition there will be steps where we have to take a pseudointersection over an ω -sequence $(X_k)_k$ of members of \mathcal{G}^n of the initial segment $(\mathcal{F}, (\mathcal{G}^n)_n)$ of the construction. Again a Cohen real is used to show that there is a pseudointersection that has infinite intersection with each element of \mathcal{G}^n . (This time an unbounded real would suffice.) Moreover, we have Hindman steps. We need a form of Hindman’s theorem that takes care of the initial segments of the ultrafilters on ω .

We will modify the following:

Theorem 2.4. [10, Theorem 5] *Let \mathcal{F} be an ordered union filter generated by $< \text{cov}(\mathcal{M})$ sets. Suppose that \mathbb{F} is partitioned into finitely many pieces. Then there is $D \in (\mathbb{F})^\omega$ such that $\text{FU}(D)$ is contained in one piece of the partition and $D \cap X$ is infinite for each $X \in \mathcal{F}$.*

We remark that by [16, Proposition 6.2] and [10, Theorem 6] the cardinal $\text{cov}(\mathcal{M})$ cannot be replaced by anything smaller.

Definition 2.5. *Let \mathcal{F} be an ordered union filter generated by $< \text{cov}(\mathcal{M})$ sets, and let \mathcal{G}^n , $n \in \omega$ be filters on ω , generated by $< \text{cov}(\mathcal{M})$ sets and $\mathcal{G}^n \subseteq \Phi(\mathcal{F})$. $\bar{d} \in \mathbf{V}_\alpha$ is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ if: For all $\bar{c} \in \mathcal{F}$, the following holds:*

$$(2.1) \quad \begin{aligned} & (\forall k \in \omega)(\text{FU}(\bar{c}) \cap \text{FU}(\bar{d}; \text{past } k) \text{ is infinite}), \text{ and} \\ & (\forall n)(\forall X \in \mathcal{G}^n)(X \cap \text{set}(\bar{c}) \cap \text{set}(\bar{d}) \text{ is infinite}). \end{aligned}$$

At many stages in the construction we add sets to the filters \mathcal{G}^n to get non-nearly coherent ultrafilters. Using the Cohen reals again, we get that the \mathcal{G}^n will grow into P -points \mathcal{V}^n . Then the sequence $\langle \mathcal{V}^n : n \in \omega \rangle$ is automatically discrete and hence ensures by Banach’s and Blass’ result that there are $2^\mathfrak{c}$ near coherence classes in $[\Phi(\mathcal{U})]$. Since every filter can be completed to an ultrafilter, these are the easy steps, corresponding to item (5) in the previous proof. For getting the non-near coherence we use that $\text{cov}(\mathcal{M}) \leq \mathfrak{u}$ and Blass’ construction from [4] and thus we can have item (6) from the previous proof also in our current construction. The Hindman steps and the steps to get stability require more care.

The following ensures that the Hindman tasks in the construction can be performed:

Theorem 2.6. *Let \mathcal{F} be an ordered union filter generated by $< \text{cov}(\mathcal{M})$ sets, and let \mathcal{G}^n , $n \in \omega$ be filters on ω , generated by $\kappa < \text{cov}(\mathcal{M})$ sets and $\mathcal{G}^n \subseteq \Phi(\mathcal{F})$. Suppose that \mathbb{F} is partitioned into finitely many pieces. Then*

there is $\bar{d} \in (\mathbb{F})^\omega$ such that $\text{FU}(\bar{d})$ is contained in one piece of the partition and $\text{FU}(\bar{d}; \text{past } k) \cap X$ is infinite for each $X \in \mathcal{F}$ and $\text{set}(\bar{d}) \cap Y$ is infinite for every $Y \in \bigcup_n \mathcal{G}^n$.

Proof. We could go for a modification of Eisworth's proof, using the Ellis-Numakura Theorem [11, 20] and Galvin and Glazer's proof of Hindman's theorem (see [14] and strengthen them in order to show that no filter \mathcal{G}^n gets lost. We use a more direct way, with Baumgartner's short proof of Hindman's theorem. In the course of the proof we argue thrice with the inequality $|\mathcal{F}|, |\mathcal{G}^n| < \text{cov}(\mathcal{M})$.

Given \bar{d} is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ and $B \subseteq \mathbb{F}$, we produce a B -homogeneous \bar{e} and a \bar{Z} , such that $\bar{e} \sqsubseteq^* \bar{d}$ and \bar{e} is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$. We look at Baumgartner's short proof of Hindman's theorem [2] and rework it step for step in order to see that the proof can be carried out within the set of $\bar{e} \in \mathbb{F}^\omega$ such that \bar{e} is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$.

Following Baumgartner, say $X \subseteq \mathbb{F}$ is large' for $\bar{d} \in (\mathbb{F})^\omega$ iff for every $\bar{d}' \sqsubseteq \bar{d}$, $\text{FU}(\bar{d}') \cap X \neq \emptyset$ or \bar{d}' is not good for $(\mathcal{F}, (\mathcal{G}^n)_n)$. We consider this modified largeness only for \bar{d} that are good for $(\mathcal{F}, (\mathcal{G}^n)_n)$.

Lemma 1(a) of Baumgartner: If X is large' for \bar{d} and $X = Y \cup Z$, then there is $\bar{d}' \sqsubseteq \bar{d}$ such that \bar{d}' is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ and either Y is large' for \bar{d}' or Z is large' for \bar{d}' .

We fill in Baumgartner's proof: Suppose it is false. Since Y is not large' for \bar{d} , there is $\bar{d}' \sqsubseteq \bar{d}$, \bar{d}' is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ and $\text{FU}(\bar{d}') \cap Y = \emptyset$. Since by the assumption that the lemma is false, Z is not large' for \bar{d}' there is $\bar{d}'' \sqsubseteq \bar{d}'$ such that $\text{FU}(\bar{d}'') \cap Z = \emptyset$ and \bar{d}'' is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$. But now $\text{FU}(\bar{d}'') \cap X = \emptyset$, contradicting the assumption that X is large' for \bar{d} .

Lemma 1(b) of Baumgartner: If X is large' for \bar{d} then for every $n \geq 0$, $\{x \in X : \min(x) > n\}$ is large' for \bar{d} . Clear, we take off only finitely many blocks and only finitely many points.

Lemma 2 of Baumgartner: Suppose X is large' for \bar{d} . Then there is a finite set $E \subseteq \text{FU}(\bar{d})$ such that for every $x \in \text{FU}(\bar{d})$ if $x \cap \bigcup E = \emptyset$, then there exists $d \in \text{FU}(E)$ such that $x \cup d \subseteq X$.

We recall and modify Baumgartner's proof: We let M be a model of ZFC^* a sufficiently rich finite fragment of ZFC that has cardinality κ such that \mathcal{F} and each of its generators, \mathcal{G}^n and each of its generators, B , and an enumeration of \mathbb{F} are elements of M . Since $\kappa < \text{cov}(\mathcal{M})$ there is a Cohen real c over M .

Suppose that the lemma is false. Then we may choose an arbitrary large $x_0 \in \text{FU}(\bar{d})$, and we take as x_0 the first $c(0)$ blocks of \bar{d} . x_0 is also conceived as $E(x_0) \subset \text{FU}(\bar{d})$ before taking the unions. Since the lemma is false there is $x_1 \in \text{FU}(\bar{d})$, $E(x_1) \subset \text{FU}(\bar{d})$ $x_1 > x_0$ and there is an $d \in \text{FU}(E(x_1))$ such that $x_1 \cup d \notin X$. Again x_1 can be chosen arbitrarily large, since we assume that the lemma is false. We take for x_1 the next $c(1)$ blocks of \bar{d} . So we go on with the inductive choice of the x_i . Then we let $y_n = x_{2n} \cup x_{2n+1}$. We show: \bar{y} is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$.

Suppose that there is some $\bar{c} \in \mathcal{F}$ and there is some k such that $\text{FU}(\bar{y}; \text{past } k) \cap \text{FU}(\bar{c}) = \emptyset$ or there are some n and some $Z \in \mathcal{G}^n$ and a $\bar{c} \in \mathcal{F}$ such that $\text{set}(\bar{y}; \text{past } k) \cap \text{set}(\bar{c}) \cap Z = \emptyset$. Since the choice of \bar{y} was done with the generic c in the generic extension $M[c]$, there is a Cohen condition $p \Vdash \text{FU}(\bar{y}; \text{past } k) \cap \text{FU}(\bar{c}) = \emptyset$, or there is a condition forcing the second fact. We show how to derive a contradiction in the second case, the first case is similar. So fix p such that

$$p \Vdash \text{FU}(\bar{y}; \text{past } k) \cap \text{set}(\bar{c}) \cap Z = \emptyset.$$

First let $m = \max\{\text{dom}(p), r\} + 1$ and extend p to an arbitrary $p' : 2m + 1 \rightarrow \omega$. Then p' decides the first $2m + 1$ values of c and also the first $m + 1$ values of \bar{y} as we set $y_{m+1} = x_{2m} \cup x_{2m+1}$. $\text{set}(\bar{d}; \text{past } k) \cap \text{set}(\bar{c}) \cap Z$ is infinite, since \bar{d} is good for $(\mathcal{F}, (\mathcal{G}^n))$. So we fix an element t in this set, that appeared in construction stage $2m + 2$ in block k after x_{2m+1} and let $q(2m + 2) = k$ then $q \geq p$ and $q \Vdash \text{set}(\bar{y}; \text{past } k) \cap \text{set}(\bar{c}) \cap X \neq \emptyset$. Contradiction.

So we showed that \bar{y} is good for $(\mathcal{F}, (\mathcal{G}^n))$. But by our choice of y_i , $\text{FU}(\bar{y}) \cap X = \emptyset$, contradiction.

Lemma 3 of Baumgartner. Suppose X is large' for \bar{d} . Then there is $e' \in \text{FU}(\bar{d})$ and there is some $\bar{d}' \sqsubseteq \bar{d}$, \bar{d}' is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ such that $\{x \in X : x \cup e' \in X\}$ is large' for $\text{FU}(\bar{d}')$.

This is proved literally as in Baumgartner, with large' instead of large: Let E be as in Lemma 2 and let $\bar{d}^1 = \bar{d} \upharpoonright [\max(E) + 1, \infty)$. Then \bar{d}^1 is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$, since we took off only finitely many blocks. So X is large' for \bar{d}^1 . For each $e' \in \text{FU}(E)$ let $X_{e'} = \{x \in X : x \cup e' \in X\}$. So

$$X \cap \text{FU}(\bar{d}^1) \subseteq \bigcup \{X_{e'} : e' \in \text{FU}(E)\}.$$

By finitely many repeated applications of Lemma 1(a) there is $\bar{d}' \sqsubseteq \bar{d}^1$ good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ and there is $e' \in \text{FU}(E)$ such that $X_{e'}$ is large' for \bar{d}' .

Lemma 4 of Baumgartner: If X is large' for \bar{d} then there is $\bar{d}' \sqsubseteq \bar{d}$ good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ in such that $\text{FU}(\bar{d}') \subseteq X$.

Proof of Lemma 4: By Lemma 3 there are sequences \bar{d}_n , e'_n , X_n such that

- (1) $\bar{d}_0 = \bar{d}$, $X_0 = X$,
- (2) $e'_n \in \text{FU}(\bar{d}_n)$ and \bar{d}_n is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ and e'_n is the union of the first $c(n)$ elements of \bar{d}_n ,
- (3) $X_{n+1} \subseteq X_n$ and $\bar{d}_{n+1} \sqsubseteq \bar{d}_n$,
- (4) X_n is large' for \bar{d}_n ,
- (5) if $x \in X_{n+1}$, then $x \cup e'_n \in X_n$,
- (6) $e'_n \cap e'_m = \emptyset$ if $m \neq n$.

Again we use the properties of the Cohen real to show that \bar{e}' is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$.

Now we choose $\bar{e} \sqsubseteq \bar{e}'$ such that e_0 is the union of the first $c(0)$ elements of \bar{e}' and

$$k_n = \max\{k : e'_k \subseteq \bigcup_{0 \leq i < n} e_i\},$$

then $e_n \in X_{k_{n+1}}$ and e_n has $c(n)$ blocks of \bar{e}' . Again we use the properties of the Cohen real to give \bar{e} is good for $(\mathcal{F}, (\mathcal{G}^n)_n)$.

If $\mathbb{F} = B_0 \cup \dots \cup B_k$, then one of the B_i is large' for any \bar{d} , since the set of requirements for being good for $(\mathcal{F}, (\mathcal{G}^n)_n)$ is directed. In the above formulae we take $B' = B$ or $B' = \mathbb{F} \setminus B$, so that B' is large' for \bar{d} . \dashv

For the stability steps in the construction we need:

Theorem 2.7. *Let \mathcal{F} be an ordered union filter generated by $< \text{cov}(\mathcal{M})$ sets, and let \mathcal{G}^n , $n \in \omega$ be filters on ω , generated by $< \text{cov}(\mathcal{M})$ sets and $\mathcal{G}^n \subseteq \Phi(\mathcal{F})$. Suppose that there is a \sqsubseteq^* -descending sequence \bar{d}^n , $n < \omega$, of $(\mathcal{F}, \bar{\mathcal{G}}^n)$ -good sequences. Then it has a lower bound that is good for $(\mathcal{F}, \bar{\mathcal{G}}^n)$.*

Proof. This is similar to just the last step of the previous proof. Take the Cohen real to pick a sufficiently large almost condensation of the \bar{d}^n . $\dashv_{2.3}$

3. REPLACING (\mathbb{F}, \cup) BY k -COLOURED BLOCK SEQUENCES (\mathbb{F}^k, \frown)

A Milliken-Taylor ultrafilter \mathcal{U}^k for block sequences with k values to block sequences with just one value $k = 1$, corresponding to “in”. We explain this:

Definition 3.1. $\mathbb{F}^k = \{s : \text{dom}(s) \in F, s : \text{dom}(s) \rightarrow \{1, \dots, k\}, s \text{ is onto}\}$. $S \subseteq \mathbb{F}^k$ is called *unmeshed* if $S = \{s_n : n \in |S|\}$ and $\text{dom}(s_0) < \text{dom}(s_1) \dots$. For unmeshed infinite sets $S \subseteq \mathbb{F}^k$ we write use also our former barred lower case letters \bar{c} and so on. Let $(\mathbb{F}^k)^{\leq \omega} = \{T \subseteq \mathbb{F}^k : T \text{ unmeshed}\}$. For $\bar{c} \in (\mathbb{F}^k)^\omega$, let $\text{FU}(\bar{c}) = \{c_{i_0} \frown c_{i_1} \frown \dots \frown c_{i_n} : n \in \omega, \text{ for } j \leq n, c_{i_j} \in S, \text{dom}(c_{i_0}) < \text{dom}(c_{i_1}) < \dots < \text{dom}(c_{i_n})\}$.

Definition 3.2. \mathcal{U}^k is called a Milliken-Taylor ultrafilter (on \mathbb{F}^k) if it is stable and if it has a basis of sets of the form $\text{FU}(S)$ for unmeshed $S \subseteq \mathbb{F}^k$.

Under CH or MA(σ -centred) Milliken-Taylor ultrafilters on \mathbb{F}^k exist. This is proved with the following strengthening of Hindman’s theorem:

Theorem 3.3. (Hindman, [13, Corollary 3.3]) *If the set \mathbb{F}^k is partitioned into finitely many pieces then there is a set $S \in (\mathbb{F}^k)^\omega$ such that $\text{FU}(S)$ is included in one piece.*

\mathcal{U}^k has the Ramsey property for n -tupels:

Theorem 3.4. *Let \mathcal{U}^k be a Milliken-Taylor ultrafilter. For any unmeshed $S \in \mathcal{U}^k$ the set $(\text{FU}(S))^n$ is partitioned into finitely many pieces then there is an unmeshed set $T \in \mathcal{U}$ such that $(\text{FU}(T))^n$ is included in one piece.*

Definition 3.5. Let \mathcal{U}^k be a Milliken-Taylor ultrafilter on \mathbb{F}^k . Then we define its core by

$$X \in \Phi(\mathcal{U}^k) \text{ iff } (\exists S \in \mathcal{U}^k)(X \supseteq \bigcup \{\text{dom}(s) : s \in S\}),$$

and its colour j core for $1 \leq j \leq k$ by

$$X \in \Phi_j(\mathcal{U}^k) \text{ iff } (\exists S \in \mathcal{U}^k)(X \supseteq \bigcup \{s^{-1''}\{j\} : s \in S\}).$$

For more on Ramsey theoretic ultrafilters on richer spaces see [12].

Theorem 3.6. Every k -valued Milliken-Taylor has at least $k + 1$ -near coherence classes of ultrafilters in $[\Phi(\mathcal{U})]$.

Proof. Then there are

$$\min_{\text{value } i}(\mathcal{U}^k) = \{\{\min(s^{-1''}\{i\}) : s \in S\} : S \in \mathcal{U}^k\}$$

and $\max_{\text{value } i}(\mathcal{U}^k)$ for $1 \leq i \leq k$ and they assume $k + 1$ near coherence classes, since

$$(3.1) \quad \begin{aligned} & \text{for some order } i_1 < \dots < i_k \text{ of } \{1, \dots, k\}, \\ & \text{for } \mathcal{U}^k \text{ many block pieces } s, \\ & \max(s^{-1''}\{i_j\}) = \min(s^{-1''}\{i_{j+1}\}) \text{ for } 1 \leq j \leq k. \end{aligned}$$

Blass' parity argument in his proof of 1.9 works in this situation and shows that there are at least $k + 1$ near coherence classes among the minimum projections and one maximum projection. We show that $\min_{\text{value } i}(\mathcal{U}^k)$ and $\max_{\text{value } i}(\mathcal{U}^k)$ are not nearly coherent for $i = 1, \dots, k$. By the above identities this gives $k + 1$ near coherence classes. For completeness we repeat the proof here. Suppose that f is finite-to-one and monotone and onto, $f(\max_{\text{value } i}(\mathcal{U}^k)) = f(\max_{\text{value } i}(\mathcal{U}^k))$ and let $I_n = f^{-1''}\{n\}$, $n \in \omega$, be adjacent increasing intervals.

Now let

$$E = \{s \in \mathbb{F}^k : |I_k \cap s^{-1''}\{i\}| \text{ is even}\}.$$

Since \mathcal{U} is an ultrafilter, it must contain E or $\mathbb{F} \setminus E$. Since \mathcal{U}^k is a union (well, better now concatenation) ultrafilter, there must be an infinite family A of pairwise disjoint members of \mathbb{F}^k such that $\text{FU}(A) \subseteq E$ or $\text{FU}(A) \cap E = \emptyset$. Since $\text{FU}(A)$ is closed under concatenation, and the sum of two odd numbers is even, we have $\text{FU}(A) \subseteq E$.

$A \in \mathcal{U}$, therefore $\max''_{\text{value } i} A = \{\max(s^{-1''}\{i\}) : s \in A\} \in \max_{\text{value } i}(\mathcal{U}^k)$ and $\min''_{\text{value } i} A = \{\min(s^{-1''}\{i\}) : s \in A\} \in \min_{\text{value } i}(\mathcal{U}^k)$. Since the two ultrafilters are nearly coherent by our assumption, the two sets have a non-empty (indeed, an infinite) intersection. So there an interval I_n with $\min(s^{-1''}\{i\}), \max(t^{-1''}\{i\}) \in I_n$. t meets besides I_n only earlier intervals, before I_n , if at all, and s meets besides I_n only later intervals after I_n , if at all. So $s \cup t \notin E$, but $s \cup t \in A$, contradiction to $\text{FU}(A) \subseteq E$. \dashv

Since by identifying colours from a k -coloured Milliken-Taylor ultrafilter we get an ℓ -coloured for $\ell < k$, we see that this proof gives an alternative way to show that under $\text{cov}(\mathcal{M}) = \mathfrak{c}$ (otherwise k -valued Milliken-Taylor ultrafilters need not exist) there are ordinary Milliken-Taylor ultrafilters with $k + 1$ near coherence classes in their core for any $k \geq 1$.

4. DESTROYING MILLIKEN TAYLOR ULTRAFILTERS WITH $\mathbb{M}(\mathcal{U})$

We review Matet forcing \mathbb{M} [6, 15] and its σ -centred suborders $\mathbb{M}(\mathcal{U})$ for a stable ordered-union ultrafilter \mathcal{U} .

Definition 4.1. *In the Matet forcing, \mathbb{M} , the conditions are pairs (s, \bar{c}) such that $s \in \mathbb{F}$ and $\bar{c} \in (\mathbb{F})^\omega$ and $s < c_0$. The forcing order is (s', \bar{c}') is stronger than (s, \bar{c}) , in symbols $(s', \bar{c}') \geq (s, \bar{c})$, iff $s \subseteq s'$ and $s' \setminus s$ is a union of finitely many of the c_n and \bar{c}' is a condensation of \bar{c} . Recall, the stronger condition is the larger one. This is in contrast to the order of almost condensation.*

In [6] it is shown that \mathbb{M} is proper. In unpublished work, Blass and Laflamme independently have shown that \mathbb{M} preserves P -points. Eisworth's work ([10, Theorem 4] or Theorem 4.4 below) implies this result, as we shall explain below.

Definition 4.2. *Given an ordered-union ultrafilter \mathcal{U} on \mathbb{F} we let $\mathbb{M}(\mathcal{U})$ consist of all pairs $(s, \bar{c}) \in \mathbb{M}$, such that $s \in \mathbb{F}$ and $\text{FU}(\bar{c}) \in \mathcal{U}$ and $s < \min(c_0)$. The forcing order is the same as in the Matet forcing.*

It is well known [15, 6] that Matet forcing \mathbb{M} can be decomposed into two steps $\mathbb{P}' * \mathbb{M}(\mathcal{U})$, such that \mathbb{P}' is ω_1 -closed (that is, every descending sequence of conditions of countable length has a lower bound) and adds a stable ordered-union ultrafilter \mathcal{U} on the set \mathbb{F} , and that $\mathbb{M}(\mathcal{U})$ is the Matet forcing with sequences from the ultrafilter (and hence it is σ -centred).

If \mathcal{U} is ultra on \mathbb{F} , then $\Phi(\mathcal{U})$ is not diagonalised (see [10, Prop. 2.3]) and also all finite-to-one images of $\Phi(\mathcal{U})$ are not diagonalised (same proof). So $\Phi(\mathcal{U})$ is not meagre.

Definition 4.3. *The Rudin-Blass ordering on filters on ω is defined as follows: Let $\mathcal{F} \leq_{RB} \mathcal{G}$ iff there is a finite-to-one h such that $h(\mathcal{F}) \subseteq h(\mathcal{G})$.*

There is also a version of the Rudin-Blass ordering with $h(\mathcal{F}) \subseteq \mathcal{G}$ instead of $h(\mathcal{F}) \subseteq h(\mathcal{G})$, however, the above version is more suitable to describe which ultrafilters are preserved. The following property of stable ordered-union ultrafilters \mathcal{U} builds on the Ramsey property of \mathcal{U} (Theorem 1.7) and will be important for our proof:

Theorem 4.4. *(Eisworth [10, “ \rightarrow ” Theorem 4, “ \leftarrow ” Cor. 2.5, this direction works also with non- P ultrafilters]) Let \mathcal{U} be a stable ordered-union ultrafilter on \mathbb{F} and let \mathcal{V} be a P -point. Iff $\mathcal{V} \not\leq_{RB} \Phi(\mathcal{U})$, then \mathcal{V} continues to generate an ultrafilter after we force with $\mathbb{M}(\mathcal{U})$.*

Now we show that for Milliken-Taylor ultrafilters there is no analogon.

Theorem 4.5. *Forcing with $\mathbb{M}(\mathcal{U})$ destroys any Milliken-Taylor ultrafilter \mathcal{V} .*

Proof. First case $\min(\mathcal{V}) \geq_{RK} \Phi(\mathcal{U})$, then by [10, Corollary 2.5], $\min(\mathcal{V})$ is destroyed and hence \mathcal{V} is destroyed. Second case: Same with $\max(\mathcal{V})$.

Third case: $\min(\mathcal{V})$ and $\max(\mathcal{V})$ are P -points that are $\not\geq_{RK} \Phi(\mathcal{U})$. Then both are preserved by 4.4. However, we show: In $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$, $\min(\mathcal{V})$ and $\max(\mathcal{V})$ are nearly coherent. So by Blass' result, \mathcal{V} is not a union ultrafilter anymore. The generic real is

$$r = \bigcup \{s : \exists \bar{c}(s, \bar{c}) \in G\}.$$

From r we get a finite-to-one function r^- , by letting $r^-(n) = |r \cap n|$. Then

$$\Vdash_{\mathbb{Q}_\alpha} \text{“}r^-(\min(\mathcal{V})) = r^-(\max(\mathcal{V}))\text{”}.$$

Given $(s, \bar{c}) \in \mathbb{M}(\mathcal{U})$ and $E \in \min(\mathcal{V})$ and $F \in \max(\mathcal{V})$, there is some $\bar{d} \sqsubseteq^* \bar{c}$, $\bar{d} \in \mathcal{U}$, such that $E \cap \text{set}(\bar{d}) = \emptyset$ and $F \cap \text{set}(\bar{d}) = \emptyset$ (this is possible since $\min(\mathcal{V}), \max(\mathcal{V}) \not\geq_{RB} \Phi(\mathcal{U})$ by the hypothesis). Now, for two suitable $k < k'$, we have $[\max(d_k), \min(d_{k'})] \cap F \neq \emptyset$ and $[\max(d_k), \min(d_{k'})] \cap E \neq \emptyset$. So $(s \cup d_{k'}, \bar{d} \upharpoonright [k' + 1, \infty))$ is stronger than (s, \bar{c}) and it forces that $r^-(E) \cap r^-(F) \neq \emptyset$. Since this works for any two sets, we have $r^-(\min(\mathcal{V}))$ is coherent with $r^-(\max(\mathcal{V}))$. \dashv

Acknowledgements: I am grateful to the Hebrew University for its generous hospitality. I thank Saharon Shelah for encouragement and for sharing his never-ending confidence. Moreover I like to thank the Institut Mittag-Leffler in Djursholm for hosting me in November/December 2009. I thank Jordi-López Abad for his question about the number of near coherence classes in the core of a Milliken-Taylor ultrafilter at this special semester.

REFERENCES

- [1] Taras Banach and Andreas Blass. The number of near-coherence classes of ultrafilters is either finite or 2^c . In Joan Bagaria and Stevo Todorćević, editors, *Set Theory*, Trends in Mathematics, pages 257–273. Birkhäuser, 2006.
- [2] James Baumgartner. A short proof of Hindman's theorem. *J. Comb. Theory (A)*, 17:384–386, 1974.
- [3] Murray Bell. On the combinatorial principle $P(c)$. *Fund. Math.*, 114:149–157, 1981.
- [4] Andreas Blass. Near coherence of filters, I: Cofinal equivalence of models of arithmetic. *Notre Dame J. Formal Logic*, 27:579–591, 1986.
- [5] Andreas Blass. Ultrafilters related to Hindman's finite unions theorem and its extensions. In S. Simpson, editor, *Logic and Combinatorics*, volume 65 of *Contemp. Math.*, pages 89–124. Amer. Math. Soc., 1987.
- [6] Andreas Blass. Applications of superperfect forcing and its relatives. In Juris Steprāns and Steve Watson, editors, *Set Theory and its Applications*, volume 1401 of *Lecture Notes in Mathematics*, pages 18–40, 1989.
- [7] Andreas Blass. Homogeneous sets from several ultrafilters. *Topology Appl.*, 156:2581–2594, 2009.

- [8] Andreas Blass and Saharon Shelah. There may be simple P_{\aleph_1} - and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed. *Ann. Pure Appl. Logic*, 33:213–243, 1987.
- [9] Andreas Blass and Saharon Shelah. Near coherence of filters. iii. a simplified consistency proof. *Notre Dame Journal of Formal Logic*, 30:530–538, 1989.
- [10] Todd Eisworth. Forcing and stable ordered-union ultrafilters. *J. Symbolic Logic*, 67:449–464, 2002.
- [11] Robert Ellis. *Lectures on Topological Dynamics*. Benjamin, 1969.
- [12] W. Timothy Gowers. Ramsey methods in Banach spaces. In William B. Johnson and Joram Lindenstrauss, editors, *Handbook of Geometry of Banach Spaces*, volume 2, pages 1071–1097. Elsevier, 2003.
- [13] Neil Hindman. Finite sums from sequences within cells of a partition of N . *J. Combin. Theory Ser. A*, 17:1–11, 1974.
- [14] Neil Hindman and Dona Strauss. *Algebra in the Stone-Ćech Compactification*. De Gruyter, 1998.
- [15] Pierre Matet. Some filters of partitions. *J. Symbolic Logic*, 53:540–553, 1988.
- [16] Pierre Matet and Janusz Pawlikowski. Ideal over ω and cardinal invariants of the continuum. *J. Symbolic Logic*, 63:1040–1054, 1998.
- [17] Heike Mildenberger and Saharon Shelah. The principle of near coherence of filters does not imply the filter dichotomy principle [MdSh:894]. *Trans. Amer. Math. Soc.*, 361:2305–2317, 2009.
- [18] Arnold Miller. There are no Q -points in Laver’s model for the Borel conjecture. *Proc. Amer. Math. Soc.*, 78:103–106, 1980.
- [19] Keith Milliken. Ramsey’s theorem with sums or unions. *Journal of Combinatorial Theory Series A*, 18:276–280, 1975.
- [20] Katsui Numakura. On bicomact semigroups. *Math. J. Okayama Univ.*, 1:99–108, 1952.
- [21] Alan D. Taylor. A canonical partition relation for finite subsets of ω . *Journal of Combinatorial Theory Series A*, 21(2):137 – 146, 1976.

HEIKE MILDENBERGER, EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY, EDMOND SAFRA CAMPUS GIVAT RAM, JERUSALEM 91904, ISRAEL
E-mail address: heike@math.huji.ac.il