

THE ROYAL
SWEDISH
ACADEMY OF
SCIENCES



**INSTITUT
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89
info@mittag-leffler.se www.mittag-leffler.se

**The relation of rapid ultrafilters and
Q-points to van der Waerden ideal**

J. Flaskova

REPORT No. 35, 2009/2010, fall

ISSN 1103-467X

ISRN IML-R- -35-09/10- -SE+fall

The relation of rapid ultrafilters and Q -points to van der Waerden ideal

JANA FLAŠKOVÁ¹

Department of Mathematics, University of West Bohemia

Abstract

We point out one of the differences between rapid ultrafilters and Q -points: Rapid ultrafilters may have empty intersection with van der Waerden ideal, whereas every Q -point has a non-empty intersection with van der Waerden ideal. Assuming Martin's axiom for countable posets we also construct a \mathscr{W} -ultrafilter which is not a Q -point.

1 Preliminaries

Let us first recall the definitions of the two classes of ultrafilters in question: An ultrafilter \mathscr{U} is a Q -point if for every partition $\{Q_n : n \in \omega\}$ of ω into finite sets there exists $A \in \mathscr{U}$ such that $|A \cap Q_n| \leq 1$ for every $n \in \omega$.

An ultrafilter \mathscr{U} is a *rapid ultrafilter* if the enumeration functions of sets in \mathscr{U} form a dominating family in $({}^\omega\omega, \leq^*)$, where the enumeration function of a set A is the unique strictly increasing function e_A from ω onto A .

Every Q -point is a rapid ultrafilter, but the converse is not true (see [Mi]).

Remember that a set $A \subseteq \omega$ is called an *AP-set* if it contains arbitrary long arithmetic progressions. It follows from van der Waerden theorem that sets which are not AP-sets form a proper ideal on ω . We will refer to this ideal as *van der Waerden ideal* and denote it by \mathscr{W} . It is known that van der Waerden ideal is an F_σ -ideal.

An ideal \mathscr{I} on ω is *tall* if for every infinite $A \subseteq \omega$ there exists infinite $B \subseteq A$ such that $B \in \mathscr{I}$. Van der Waerden ideal is clearly a tall ideal.

An ideal \mathscr{I} on ω is a *P-ideal* if for every $A_n \in \mathscr{I}$, $n \in \omega$ there exists $A \in \mathscr{I}$ such that $A_n \setminus A$ is finite for all $n \in \omega$. It is easy to see that van der Waerden ideal is not a *P-ideal* (consider e.g. sets $A_n = \{2^i + n : i \in \omega\}$).

¹The current research was partially supported by the Institut Mittag-Leffler (Djursholm, Sweden) and by the European Science Foundation (in the realm of the activity entitled 'New Frontiers of Infinity: Mathematical, Philosophical and Computational Prospects').

Baumgartner introduced \mathcal{I} -ultrafilters in [B]. We will repeat the definition and introduce a weaker notion of weak \mathcal{I} -ultrafilters. So, assume \mathcal{I} is a tall ideal on ω :

An ultrafilter \mathcal{U} on ω is called an \mathcal{I} -ultrafilter if for every function $f : \omega \rightarrow \omega$ there exists $U \in \mathcal{U}$ such that $f[U] \in \mathcal{I}$.

An ultrafilter \mathcal{U} on ω is called a weak \mathcal{I} -ultrafilter if for every finite-to-one function $f : \omega \rightarrow \omega$ there exists $U \in \mathcal{U}$ such that $f[U] \in \mathcal{I}$.

It is obvious that every \mathcal{I} -ultrafilter is a weak \mathcal{I} -ultrafilter and every weak \mathcal{I} -ultrafilter has nonempty intersection with the ideal \mathcal{I} .

2 Q-points and van der Waerden ideal

In this section we examine the connection between Q -points and van der Waerden ideal. First of all we prove that each Q -point not only has a nonempty intersection with van der Waerden ideal, but it is in fact a weak \mathcal{W} -ultrafilter.

Proposition 2.1. *Every Q -point is a weak \mathcal{W} -ultrafilter.*

Proof. Consider partition of ω into finite sets $\omega = \bigcup_{n \in \omega} I_n$ where $I_0 = \{0, 1\}$ and $I_n = [2^n, 2^{n+1})$ for every $n > 1$. Assume $f : \omega \rightarrow \omega$ is an arbitrary finite-to-one function. If \mathcal{U} is a Q -point then there exists $U \in \mathcal{U}$ such that $|U \cap f^{-1}[I_n]| \leq 1$ for every $n \in \omega$. Now enumerate (increasingly) $f[U] = \{u_n : n \in \omega\}$. Either $U_0 = f^{-1}\{u_{2k} : k \in \omega\}$ or $U_1 = f^{-1}\{u_{2k+1} : k \in \omega\}$ belongs to the ultrafilter \mathcal{U} . Without loss of generality we may assume $U_0 \in \mathcal{U}$. It follows from the definition that $f[U_0]$ does not contain an arithmetic progression of length 3, thus $f[U_0] \in \mathcal{W}$. \square

Weak \mathcal{W} -ultrafilter in the previous proposition cannot be replaced with \mathcal{I} -ultrafilter because Q -points in general need not be \mathcal{I} -ultrafilters. It follows from Proposition 2.4.7 in [F] according to which assuming Martin's axiom for countable posets for every tall ideal \mathcal{I} there exists a Q -point which is not an \mathcal{I} -ultrafilter.

One can, of course, ask whether the implication in Proposition 2.1 can be reversed. The answer is negative. If Martin's axiom for countable posets holds then there exists even a \mathcal{W} -ultrafilter which is not a Q -point. We will construct such an ultrafilter in the rest of this section starting with the following rather technical statement:

Proposition 2.2. (MA_{ctble}) Assume \mathcal{F} is a filter base of cardinality less than \mathfrak{c} with the following property

$$(\spadesuit) \quad (\forall F \in \mathcal{F}) (\forall k \in \omega) (\exists n \in \omega) |F \cap [2^n, 2^{n+1})| \geq k.$$

Assume $f \in {}^\omega\omega$. Then there exists $G \in [\omega]^\omega$ such that $f[G] \in \mathcal{W}$ and the filter base generated by \mathcal{F} and G has property (\spadesuit) .

Proof. If there exists $F \in \mathcal{F}$ such that $f[F] \in \mathcal{W}$, let $G = F$. If there exists $K \in [\omega]^{<\omega}$ such that the filter base generated by \mathcal{F} and $f^{-1}[K]$ has property (\spadesuit) , then put $G = f^{-1}[K]$. In the following we will assume that no such K exists. So we will assume that for every $K \in [\omega]^{<\omega}$ and for every $F \in \mathcal{F}$ and for every $k \in \omega$ there exists $n \in \omega$ such that $|(F \setminus f^{-1}[K]) \cap [2^n, 2^{n+1})| \geq k$ and we construct a suitable G eventually making use of Martin's axiom.

Consider a set

$$P = \{K \in [\omega]^{<\omega} : f[K] \text{ contains no arithmetic progressions of length } 3\}$$

equipped with partial order \leq_P defined in the following way: $K \leq_P L$ if and only if $K = L$ or $K \supset L$ and $\min(K \setminus L) > \max L$. For $F \in \mathcal{F}$ and $k \in \omega$ define $D_{F,k} = \{K \in P : (\exists n \in \omega) |K \cap F \cap [2^n, 2^{n+1})| \geq k\}$.

Claim: $D_{F,k}$ is a dense subset of (P, \leq_P) for every $F \in \mathcal{F}$, $k \in \omega$.

Proof of the Claim. Let $L \in P$ be arbitrary. We want to find $K \in D_{F,k}$ such that $K \leq_P L$. Let $n_0 = \max\{n \in \omega : L \cap [2^n, 2^{n+1}) \neq \emptyset\}$.

Case I. $\sup_{n \in \omega} |f[F \cap [2^n, 2^{n+1})]| = m < \infty$

For $N = \{i \in \omega : i \leq 3 \cdot \max f[L]\}$ there exists $n(N) > n_0$ such that $|(F \setminus f^{-1}[N]) \cap [2^{n(N)}, 2^{n(N)+1})| \geq k \cdot (m+1)$. According to the assumption of Case I. $|f[F \cap [2^{n(N)}, 2^{n(N)+1})]| \leq m$. Now, it follows from Dirichlet's box principle that there exist

$$l \in f[F \cap [2^{n(N)}, 2^{n(N)+1})] \text{ and } L' \subseteq (F \setminus f^{-1}[N]) \cap [2^{n(N)}, 2^{n(N)+1})$$

such that $|L'| \geq k$, $f[L'] = \{l\}$ and $l > 3 \cdot \max f[L]$. Put $K = L \cup L'$. It remains to verify that K is as required:

1. Observe that $K \in P$. Any arithmetic progression of length 3 in $f[K]$ contains l (otherwise $f[L]$ contains an arithmetic progression of length 3 in contradiction to the assumption $L \in P$). However, for every $a, b \in f[L]$ we have

$$l - \max\{a, b\} > 2 \cdot \max f[L] > \max f[L] > |a - b|.$$

Thus a, b and l cannot form an arithmetic progression.

2. Obviously, $K \in D_{F,k}$ because

$$|K \cap F \cap [2^{n(N)}, 2^{n(N)+1})| \geq |L' \cap F \cap [2^{n(N)}, 2^{n(N)+1})| \geq k.$$

3. Notice that $K \leq_P L$ because $n(N) > n_0$ and consequently

$$\min(K \setminus L) = \min L' > \max L.$$

Case II. $\sup_{n \in \omega} |f[F \cap [2^n, 2^{n+1})]| = \infty$

According to the assumption of Case II. there exists $n_1 > n_0$ such that $|f[F \cap [2^{n_1}, 2^{n_1+1})]| \geq 3 \cdot \max f[L] + (|L| + k)^2$. Let

$$A_0 = \{m \in f[F \cap [2^{n_1}, 2^{n_1+1})] : m > 3 \cdot \max f[L]\}.$$

Obviously, $|A_0| \geq (|L| + k)^2$. Now, choose $l_i \in A_0$ for $i = 0, \dots, k-1$ so that $f[L] \cup \{l_i : i < k\}$ contains no arithmetic progressions of length 3. This can be done by induction on i :

For $i = 0$ let $l_0 = \min A_0$. The set $B_1 = f[L] \cup \{l_0\}$ does not contain arithmetic progressions of length 3 because $f[L]$ did not and for arbitrary $a, b \in f[L]$

$$l_0 - \max\{a, b\} > 2 \cdot \max f[L] > \max f[L] > |a - b|.$$

If $0 < i < k-1$ and $l_j \in A_0$ for $j < i$ are already known such that $B_i = f[L] \cup \{l_j : j < i\}$ contains no arithmetic progressions of length 3, define

$$A_i = \{m \in A_0 : (\exists a, b \in B_i) a, b, m \text{ form an arithmetic progression}\}.$$

Since $|B_i| = |L| + i$ the set A_i has at most $\frac{1}{2}(|L| + i)(|L| + i - 1)$ elements. So $A_0 \setminus A_i \neq \emptyset$ and we may define $l_i = \min(A_0 \setminus A_i)$. It follows from the construction that $B_{i+1} = B_i \cup \{l_i\}$ contains no arithmetic progressions of length 3.

Finally, let $L' = F \cap [2^{n_1}, 2^{n_1+1}) \cap f^{-1}[\{l_i : i < k\}]$ and put $K = L \cup L'$. It remains to verify that K is as required:

1. Obviously, $K \in P$ because $f[K] = f[L] \cup \{l_i : i < k\}$ and the latter set contains no arithmetic progressions of length 3.
2. Observe that $K \in D_{F,k}$ because

$$|K \cap F \cap [2^{n_1}, 2^{n_1+1})| \geq |L'| \geq k.$$

3. Notice that $K \leq_P L$ because $n_1 > n_0$ and consequently

$$\min(K \setminus L) = \min L' > \max L.$$

□ *Claim.*

Since family $\mathcal{D} = \{D_{F,k} : F \in \mathcal{F}, k \in \omega\}$ consists of dense subsets of countable poset P and $|\mathcal{D}| < \mathfrak{c}$, it follows from Martin's axiom for countable posets that there is a \mathcal{D} -generic filter \mathcal{G} .

Let $G = \bigcup \{K : K \in \mathcal{G}\}$. It remains to verify that:

- $f[G]$ contains no arithmetic progressions of length 3, i.e. $f[G] \in \mathcal{W}$

Consider $a, b, c \in f[G]$ arbitrary. There exist $K_a, K_b, K_c \in \mathcal{G}$ such that $a \in f[K_a]$, $b \in f[K_b]$ and $c \in f[K_c]$. Since \mathcal{G} is a filter, there exists $K_0 \in \mathcal{G}$ which is \leq_P -below all three sets K_a, K_b, K_c . Thus $a, b, c \in K_0$. Now, K_0 as an element of P contains no arithmetic progressions of length 3, in particular a, b and c do not form an arithmetic progression.

- $(\forall F \in \mathcal{F}) (\forall k \in \omega) (\exists n \in \omega) |F \cap G \cap [2^n, 2^{n+1}]| \geq k$.

Take $k \geq 1$ arbitrary. For every $K \in \mathcal{G} \cap D_{F,k}$ we have $G \supseteq K$ and $|F \cap G \cap [2^n, 2^{n+1}]| \geq |F \cap K \cap [2^n, 2^{n+1}]| \geq k$ for some n . □

Lemma 2.3. *Every filter base \mathcal{F} which has property (\spadesuit) introduced in Proposition 2.2 can be extended into an ultrafilter which is not a Q -point.*

Proof. The family $\{[2^n, 2^{n+1}] : n \in \omega\}$ is a partition of ω into finite sets witnessing the fact that an ultrafilter with property (\spadesuit) is not a Q -point. Therefore we will show that every filter base \mathcal{F} with property (\spadesuit) can be extended into an ultrafilter with property (\spadesuit) . This can be accomplished by transfinite induction on $\alpha < \mathfrak{c}$ where in each non-limit step one subset of ω is considered and the filter base is extended by either the set itself or its complement.

To this end, consider a filter base \mathcal{F} with property (\spadesuit) and $A \subseteq \omega$: Either $(\forall F \in \mathcal{F}) (\forall k \in \omega) (\exists n \in \omega) |F \cap A \cap [2^n, 2^{n+1}]| \geq k$ and then the filter base \mathcal{F}' generated by \mathcal{F} and A has property (\spadesuit) .

Or $(\exists F_0 \in \mathcal{F}) (\exists k_0 \in \omega) (\forall n \in \omega) |F_0 \cap A \cap [2^n, 2^{n+1}]| < k_0$ and then the filter base \mathcal{F}' generated by \mathcal{F} and $\omega \setminus A$ has property (\spadesuit) . Indeed, since \mathcal{F} has property (\spadesuit) for every $F \in \mathcal{F}$ and for every $k \in \omega$ there exists $n \in \omega$ such that $|F \cap F_0 \cap [2^n, 2^{n+1}]| \geq k + k_0$. However, $|F \cap F_0 \cap A \cap [2^n, 2^{n+1}]| < k_0$ and so $|F \cap (\omega \setminus A) \cap [2^n, 2^{n+1}]| \geq |F \cap F_0 \cap (\omega \setminus A) \cap [2^n, 2^{n+1}]| \geq k$. □

Theorem 2.4. *(MA_{ctble}) There is a \mathcal{W} -ultrafilter which is not a Q -point.*

Proof. Enumerate all functions ${}^\omega\omega = \{f_\alpha : \alpha < \mathfrak{c}\}$. By transfinite induction on $\alpha < \mathfrak{c}$ we will construct filter bases \mathcal{F}_α so that the following conditions are satisfied:

- (i) \mathcal{F}_0 is the Fréchet filter
- (ii) $\mathcal{F}_\alpha \subseteq \mathcal{F}_\beta$ whenever $\alpha \leq \beta$
- (iii) $\mathcal{F}_\gamma = \bigcup_{\alpha < \gamma} \mathcal{F}_\alpha$ for γ limit
- (iv) $(\forall \alpha) |\mathcal{F}_\alpha| \leq |\alpha| \cdot \omega$
- (v) $(\forall \alpha) \mathcal{F}_\alpha$ has property (\spadesuit)
- (vi) $(\forall \alpha) (\exists F \in \mathcal{F}_{\alpha+1}) f_\alpha[F] \in \mathcal{W}$

Conditions (i)–(iii) take care of the beginning the induction and keep it going. Because of (iii) conditions (iv) and (v) are satisfied at limit stages of the construction and one only has to ensure that conditions (iv)–(vi) hold at non-limit induction steps.

Induction step: Suppose we already know \mathcal{F}_α . If there is $A \in \mathcal{F}_\alpha$ such that $f_\alpha[A] \in \mathcal{W}$ then simply put $\mathcal{F}_{\alpha+1} = \mathcal{F}_\alpha$. If $f_\alpha[F] \notin \mathcal{W}$ for every $F \in \mathcal{F}_\alpha$ then apply Proposition 2.2 on filter base \mathcal{F}_α and function f_α . Let $\mathcal{F}_{\alpha+1}$ be the filter base generated by \mathcal{F}_α and G .

Finally, let $\mathcal{F} = \bigcup_{\alpha < \mathfrak{c}} \mathcal{F}_\alpha$. Since the filter base \mathcal{F} has property (\spadesuit) , it can be extended into an ultrafilter which is not a Q -point according to Lemma 2.3. It is a \mathcal{W} -ultrafilter which is not a Q -point because each ultrafilter which extends \mathcal{F} is a \mathcal{W} -ultrafilter due to condition (vi). \square

3 Rapid ultrafilters and van der Waerden ideal

In this section we prove that rapid ultrafilters, unlike Q -points, may have an empty intersection with van der Waerden ideal. We will construct such an ultrafilter assuming Martin's axiom for countable posets in Theorem 3.4, which is actually slightly stronger than required.

Let us start with the definition of summable ideals. They play an important role in an alternative characterization of rapid ultrafilters, which we use later in the proof.

For a function $g : \omega \rightarrow (0, \infty)$ such that $\sum_{n \in \omega} g(n) = +\infty$ the family

$$\mathcal{I}_g = \{A \subseteq \omega : \sum_{a \in A} g(a) < +\infty\}$$

is an ideal on ω , which we call a *summable ideal (determined by function g)*. A summable ideal is tall if and only if $\lim_{n \rightarrow \infty} g(n) = 0$.

A characterization of rapid ultrafilters involving summable ultrafilters can be found e.g. in [M-A] as Theorem 2.8.10. We restate it here in terms of weak \mathcal{I} -ultrafilters.

Proposition 3.1 ([M-A]). *The following statements are equivalent for an ultrafilter $\mathcal{U} \in \omega^*$:*

1. \mathcal{U} is a rapid ultrafilter
2. \mathcal{U} is a weak \mathcal{I}_g -ultrafilter for every tall summable ideal \mathcal{I}_g
3. \mathcal{U} is a weak \mathcal{I} -ultrafilter for every tall analytic P -ideal \mathcal{I}

Notice that analytic ideals in the third statement of the previous theorem are P -ideals. Since van der Waerden ideal is not a P -ideal, the theorem does not contradict the existence of rapid ultrafilters, whose intersection with van der Waerden ideal is empty.

If we assume Martin's axiom for countable posets then there are not only rapid ultrafilters containing merely AP-sets, but also hereditarily rapid ultrafilters, which we define as follows:

An ultrafilter \mathcal{U} is a *hereditarily rapid ultrafilter* if it is a rapid ultrafilter such that the ultrafilter $f(\mathcal{U})$ generated by the sets $\{f[U] : U \in \mathcal{U}\}$ is again a rapid ultrafilter for every $f \in {}^\omega\omega$.

The characterization of rapid ultrafilters in Proposition 3.1 reformulates for hereditarily rapid ultrafilters in a straightforward way:

Proposition 3.2. *The following statements are equivalent for $\mathcal{U} \in \omega^*$:*

1. \mathcal{U} is a hereditarily rapid ultrafilter
2. \mathcal{U} is an \mathcal{I}_g -ultrafilter for every tall summable ideal \mathcal{I}_g
3. \mathcal{U} is an \mathcal{I} -ultrafilter for every tall analytic P -ideal \mathcal{I}

We will concentrate on the first two statements of the previous theorem and we will construct our ultimate goal – a hereditarily rapid ultrafilter which does not intersect van der Waerden ideal – as an ultrafilter which is \mathcal{I}_g -ultrafilter for every tall summable ideal \mathcal{I}_g . The following lemma is crucial for the induction step of the construction:

Lemma 3.3. (MA_{ctble}) *Let \mathcal{I}_g be a tall summable ideal. Assume \mathcal{F} is a filter base, $|\mathcal{F}| < \mathfrak{c}$, $\mathcal{F} \cap \mathcal{U} = \emptyset$ and $f \in {}^\omega\omega$. Then there exists $G \in [\omega]^\omega$ such that $f[G] \in \mathcal{I}_g$ and $F \cap G$ is an AP-set for every $F \in \mathcal{F}$.*

Proof. If there exists $F \in \mathcal{F}$ such that $f[F] \in \mathcal{S}_g$, put $G = F$. If there exists $K \in [\omega]^{<\omega}$ such that $F \cap f^{-1}[K]$ is an AP-set for every $F \in \mathcal{F}$, then put $G = f^{-1}[K]$. In the following we will assume that no such set exists, i.e.

(♣) for every $K \in [\omega]^{<\omega}$ there is $F_K \in \mathcal{F}$ such that $F_K \cap f^{-1}[K] \in \mathcal{W}$.

This also means that $F \setminus f^{-1}[K]$ is an AP-set for every $F \in \mathcal{F}$.

Consider poset

$$P = \{K \in [\omega]^{<\omega} : \sum_{a \in f[K]} g(a) \leq \left(2 - \frac{1}{2^{|K|}}\right) \cdot \max_{a \in f[K]} g(a)\}$$

with partial order \leq_P defined by: $K \leq_P L$ if and only if $K = L$ or $K \supset L$ and $\min(K \setminus L) > \max L$. Now, for every $F \in \mathcal{F}$ and every $k \geq 1$ define $D_{F,k} = \{K \in P : K \cap F \text{ contains an arithmetic progression of length } k\}$.

Claim. $D_{F,k}$ is dense in P for every $F \in \mathcal{F}$ and $k \geq 1$.

Proof of the Claim. Take $L \in P$ arbitrary. Since $\lim_{n \rightarrow \infty} g(n) = 0$ there exists $n_L \in \omega$ such that for every $n > n_L$

$$g(n) < \frac{1}{2^{|L|+1} \cdot k} \cdot \max_{a \in f[L]} g(a).$$

According to the assumption (♣) there exists $F_{n_L} \in \mathcal{F}$ such that $F_{n_L} \cap f^{-1}[0, n_L] \in \mathcal{W}$. It follows that $A_{n_L} = (F \cap F_{n_L}) \setminus f^{-1}[0, n_L]$ is an infinite AP-set, thus one can choose an arithmetic progression $L' \subset A_{n_L}$ such that $|L'| = k$ and $\min L' > \max L$. Let $K = L \cup L'$. Observe that due to the choice of L' one has $\max_{a \in f[K]} g(a) = \max_{a \in f[L]} g(a)$. To see that $K \in P$ notice that

$$\begin{aligned} \sum_{a \in f[K]} g(a) &\leq \sum_{a \in f[L]} g(a) + \sum_{a \in f[L']} g(a) \leq \\ &\leq \left(2 - \frac{1}{2^{|L|}}\right) \cdot \max_{a \in f[L]} g(a) + |L'| \cdot \max_{a \in f[L']} g(a) \leq \\ &\leq \left(2 - \frac{1}{2^{|L|}}\right) \cdot \max_{a \in f[L]} g(a) + k \cdot \frac{1}{2^{|L|+1} \cdot k} \cdot \max_{a \in f[L]} g(a) \leq \\ &\leq \left(2 - \frac{1}{2^{|L|+1}}\right) \max_{a \in f[L]} g(a) \leq \left(2 - \frac{1}{2^{|K|}}\right) \max_{a \in f[K]} g(a) \end{aligned}$$

It is obvious that $K \leq_P L$. Also $K \in D_{F,k}$ because $K \cap F \supseteq L' \cap F$ contains an arithmetic progression of length k . \square *Claim.*

Since family $\mathcal{D} = \{D_{F,k} : F \in \mathcal{F}, k \geq 1\}$ consists of dense subsets of countable poset P and $|\mathcal{D}| < \mathfrak{c}$, it follows from Martin's axiom for countable posets that there is a \mathcal{D} -generic filter \mathcal{G} .

Let $G = \bigcup\{K : K \in \mathcal{G}\}$. It remains to verify that:

- $f[G] \in \mathcal{I}_g$, i.e. $\sum_{a \in f[G]} g(a) < +\infty$

Enumerate $f[G] = \{u_n : n \in \omega\}$. For every n there exists $K_n \in \mathcal{G}$ such that $u_n \in f[K_n]$. We may assume $K_{n+1} \leq_P K_n$ (and thus $f[K_{n+1}] \supseteq f[K_n]$) for every $n \in \omega$ because \mathcal{G} is a filter. Since $f[G] = \bigcup_{n \in \omega} f[K_n]$, we get

$$\sum_{a \in f[G]} g(a) = \lim_{n \rightarrow \infty} \sum_{a \in f[K_n]} g(a) \leq \lim_{n \rightarrow \infty} \left(2 - \frac{1}{2^{|K_n|}}\right) \max_{a \in f[K_n]} g(a) \leq 2 \cdot \max_{a \in \omega} g(a).$$

- $(\forall F \in \mathcal{F}) G \cap F$ is an AP-set

Take $k \geq 1$ arbitrary. For every $K \in \mathcal{G} \cap D_{F,k}$ we have $G \supseteq K$ and $K \cap F$ contains an arithmetic progression of length k . Hence $G \cap F$ contains arithmetic progressions of arbitrary length, i.e. $G \cap F$ is an AP-set. \square

Theorem 3.4. (*MA_{ctble}*) *There is a hereditarily rapid ultrafilter \mathcal{U} such that $\mathcal{U} \cap \mathcal{W} = \emptyset$.*

Proof. Enumerate all pairs $\{\langle f_\alpha, g_\alpha \rangle : \alpha < \mathfrak{c}\}$ where $f_\alpha \in {}^\omega\omega$ and \mathcal{I}_{g_α} is a tall summable ideal. By transfinite induction on $\alpha < \mathfrak{c}$ we will construct filter bases \mathcal{F}_α so that the following conditions are satisfied:

- (i) \mathcal{F}_0 is the Fréchet filter
- (ii) $\mathcal{F}_\alpha \subseteq \mathcal{F}_\beta$ whenever $\alpha \leq \beta$
- (iii) $\mathcal{F}_\gamma = \bigcup_{\alpha < \gamma} \mathcal{F}_\alpha$ for γ limit
- (iv) $(\forall \alpha) |\mathcal{F}_\alpha| \leq |\alpha| \cdot \omega$
- (v) $(\forall \alpha) (\forall F \in \mathcal{F}_\alpha) F$ is an AP-set
- (vi) $(\forall \alpha) (\exists F \in \mathcal{F}_{\alpha+1}) f_\alpha[F] \in \mathcal{I}_{g_\alpha}$

Conditions (i)–(iii) take care of the beginning of the induction and keep it going. Because of (iii) conditions (iv) and (v) are satisfied at limit stages of the construction and one only has to ensure that conditions (iv)–(vi) hold at non-limit induction steps.

Induction step: Suppose we already know \mathcal{F}_α . If there is $A \in \mathcal{F}_\alpha$ such that $f_\alpha[A] \in \mathcal{I}_{g_\alpha}$ then simply put $\mathcal{F}_{\alpha+1} = \mathcal{F}_\alpha$. If $f_\alpha[F] \notin \mathcal{I}_{g_\alpha}$ for every $F \in \mathcal{F}_\alpha$ then apply Lemma 3.3 on ideal \mathcal{I}_{g_α} , filter base \mathcal{F}_α and function f_α . Let $\mathcal{F}_{\alpha+1}$ be the filter base generated by \mathcal{F}_α and G .

Finally, let $\mathcal{F} = \bigcup_{\alpha < \mathfrak{c}} \mathcal{F}_\alpha$. Since F is an AP-set for every $F \in \mathcal{F}$ the filter base \mathcal{F} can be extended to an ultrafilter \mathcal{U} which does not intersect

the van der Waerden ideal. Every ultrafilter which extends \mathcal{F} , however, is an \mathcal{I}_{g_α} -ultrafilter for every tall summable ideal \mathcal{I}_{g_α} because of condition (vi). Thus \mathcal{U} is a hereditarily rapid ultrafilter satisfying $\mathcal{U} \cap \mathcal{W} = \emptyset$. \square

Corollary 3.5. (MA_{ctble}) *There exists a rapid ultrafilter \mathcal{U} such that \mathcal{U} contains only AP-sets.* \square

References

- [B] J. Baumgartner, Ultrafilters on ω , *J. Symbolic Logic* **60**, no. 2, 624–639, 1995.
- [F] J. Flašková, Ultrafilters and small sets, *Ph.D. thesis*, Charles University, Prague 2006.
- [M-A] D. Meza Alcántara, Ideals and filters on countable sets. *Ph.D. thesis*. UNAM México, 2009.
- [Mi] A. W. Miller, There are no Q -points in Laver’s model for the Borel conjecture, *Proc. Amer. Math. Soc.* **78**, no. 1, 103–106, 1980.