

THE ROYAL  
SWEDISH  
ACADEMY OF  
SCIENCES



**INSTITUT  
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden  
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89  
info@mittag-leffler.se www.mittag-leffler.se

## **Milnor-Wood inequalities for products**

M. Bucher and T. Gelander

REPORT No. 5, 2011/2012, spring

ISSN 1103-467X

ISRN IML-R- -5-11/12- -SE+spring

# MILNOR–WOOD INEQUALITIES FOR PRODUCTS

MICHELLE BUCHER AND TSACHIK GELANDER

ABSTRACT. We prove Milnor–wood inequalities for local products of manifolds. As a consequence, we establish the generalized Chern Conjecture for products  $M \times \Sigma^k$  for any manifold  $M$  and  $k$  copies of a surface  $\Sigma$  for  $k$  sufficiently large.

## 1. INTRODUCTION

Let  $M$  be an  $n$ -dimensional topological manifold. Consider the Euler class  $\varepsilon_n(\xi) \in H^n(M, \mathbb{R})$  and Euler number  $\chi(\xi) = \langle \varepsilon_n(\xi), [M] \rangle$  of oriented  $\mathbb{R}^n$ -vector bundles over  $M$ . We say that the manifold  $M$  satisfies a Milnor–wood inequality with constant  $c$  if for every flat oriented  $\mathbb{R}^n$ -vector bundles  $\xi$  over  $M$ , the inequality

$$|\chi(\xi)| \leq c \cdot |\chi(M)|$$

holds. Recall that a bundle is flat if it is induced by a representation of the fundamental group  $\pi_1(M)$ . We denote by

$$MW(M) \in \mathbb{R} \cup \{+\infty\}$$

the smallest such constant.

If  $X$  is a simply connected Riemannian manifold, we denote

$$\widetilde{MW}(X) := \sup\{MW(M) : M \text{ is closed quotients of } X\}.$$

Milnor’s seminal inequality [Mi58] amounts to showing that the Milnor–wood constant of the hyperbolic plane  $\mathcal{H}$  is  $\widetilde{MW}(\mathcal{H}) = 1/2$ , and in [BuGe11], we showed that  $\widetilde{MW}(\mathcal{H}^n) = 1/2^n$ .

In this note we prove a product formula for the Milnor–wood constants of general closed manifolds:

**Theorem 1.** *For any pair of compact manifolds  $M_1, M_2$*

$$MW(M_1 \times M_2) = MW(M_1) \cdot MW(M_2).$$

For the product formula for universal Milnor–wood constant, we restrict to Hadamard manifolds:

---

The authors gratefully acknowledge support by the Institut Mittag-Leffler (Djursholm, Sweden). Michelle Bucher acknowledges support by the Swiss National Science Foundation grant PP00P2-128309/1. Tseachik Gelander acknowledges support by the Israel Science Foundation and the European Research Council.

**Theorem 2.** *Let  $X_1, X_2$  be Hadamard manifolds. Then*

$$\widetilde{MW}(X_1 \times X_2) = \widetilde{MW}(X_1) \cdot \widetilde{MW}(X_2).$$

One important application of Milnor–wood inequalities is to make progress on the generalized Chern Conjecture.

**Conjecture 3** (Generalized Chern Conjecture). *Let  $M$  be a closed oriented aspherical manifold. If the tangent bundle  $TM$  of  $M$  admits a flat structure then  $\chi(M) = 0$ .*

This conjecture has been suggested by Milnor [Mi58]<sup>1</sup> and is a strong version of the famous Chern conjecture which merely predicts the vanishing of the Euler characteristic for affine manifolds, equivalently, for manifolds admitting a torsion free flat connection.

As pointed out in [Mi58], if  $MW(M) < 1$  then the Generalized Chern Conjecture holds for  $M$ . Indeed, if  $\chi(M) \neq 0$  the inequality

$$|\chi(M)| = |\chi(TM)| \leq MW(M) \cdot |\chi(M)| < |\chi(M)|$$

leads to a contradiction to the assumption that  $M$  has a flat structure.

One can use Theorem 1 to extend the family of manifolds satisfying the Generalized Chern Conjecture. For instance, we prove a stable variant of the Generalized Chern Conjecture:

**Corollary 4.** *Let  $M$  be a manifold with a finite Milnor–Wood constant. Then the product  $M \times \Sigma^k$ , where  $\Sigma$  is a surface of genus  $\geq 2$  and  $k > \log_2(MW(X))$  satisfies the Generalized Chern Conjecture. In particular, if  $\chi(M) \neq 0$ , then  $M \times \Sigma^k$  does not admit an affine structure.*

**Remark 5.** 1. *One can replace  $\Sigma^k$  in Corollary 4 by any  $\mathcal{H}^k$ -manifold.*

2. *The corollary is somehow dual to a question of Yves Benoist [Be00, Section 3, p. 19] asking whether for every closed manifold  $M$  there exists  $m$  such that  $M \times S^m$  admits an affine structure. For example, if  $M$  is a hyperbolic manifold or a sphere, the product  $M \times S^1$  admits an affine structure. On the other hand, if  $M$  admits a quaternionic hyperbolic structure then  $m = 1$  will not suffice, since the holonomy representation of  $\pi_1(M)$  is super-rigid in  $Sp(2,1)$  by Corlette’s theorem and the latter has no nontrivial 9-dimensional linear representations.*

Note that since there are only finitely many isomorphism classes of oriented  $\mathbb{R}^n$ -bundles which admit a flat structure, it is immediate

---

<sup>1</sup>In [Mi58] Milnor suggested the generalized conjecture without the assumption that  $M$  is aspherical, however Smillie [Sm77] gave counter examples, in any even dimension  $\neq 2$ , when this assumption is omitted.

that the set

$$\{|\chi(\xi)| \mid \xi \text{ is a flat oriented } \mathbb{R}^n\text{-bundle over } M\}$$

is finite for every  $M$ . In particular, if  $\chi(M) \neq 0$ , there exists a finite Milnor–Wood constant  $MW(M) < +\infty$ . However, in general, the Milnor–Wood constant can be infinite, since the implication

$$\chi(M) = 0 \Rightarrow \chi(\xi) = 0,$$

for  $\xi$  a flat oriented  $\mathbb{R}^n$ -bundle, does not hold in general as we will show in Section 5. Our example is inspired by Smillie’s counterexample of the Generalized Chern Conjecture [Sm77] for nonaspherical manifolds, and likewise this manifold is nonaspherical.

The following questions are quite natural:

- (1) Does there exist a finite constant  $c(n)$  depending on  $n$  only such that  $MW(M) \leq c(n)$  for every closed aspherical  $n$ -manifold?
- (2) Let  $X$  be a contractible Riemannian manifold such that there exists a closed  $X$ -manifold  $M$  with  $MW(M) < \infty$ . Is  $\widetilde{MW}(X)$  necessarily finite?
- (3) Does  $\chi(M) = 0 \Rightarrow \chi(\xi) = 0$  for flat oriented  $\mathbb{R}^n$ -bundles  $\xi$  over aspherical manifolds  $M$ ?

## 2. REPRESENTATIONS OF PRODUCTS

**Lemma 6.** *Let  $H_1, H_2$  be groups and  $\rho : H_1 \times H_2 \rightarrow GL_n(\mathbb{R})$  a representation of the direct product and suppose that  $\rho(H_i)$  is non-amenable for both  $i = 1, 2$ . Then, up to replacing the  $H_i$ ’s by finite index subgroups, either*

- $V = \mathbb{R}^n$  decomposes as an invariant direct sum  $V = V' \oplus V''$  where the restriction  $\rho|_{V'} = \rho'_1 \otimes \rho'_2$  is a nontrivial tensor representation, or
- $V = V_1 \oplus V_2$  where  $G_i$  is scalar on  $V_i$ .

*Proof.* This can be easily deduced from the proof of [BuGe11, Proposition 6.1].  $\square$

**Proposition 7.** *Let  $H = \prod_{i=1}^k H_i$  be a direct product of groups and let  $\rho : H \rightarrow GL_n^+(\mathbb{R})$  be an orientable representation, where  $n = \sum_{i=1}^k m_i$ . Suppose that  $\rho(H_i)$  is nonamenable for every  $i$ . Then, up to replacing the  $H_i$ ’s by finite index subgroups  $H'_i = \prod_{i=1}^k H'_i$ , either*

- (1) *there exists  $1 \leq i_0 < k$  such that  $V = \mathbb{R}^n$  decomposes non-trivially to an invariant direct sum  $V = V' \oplus V''$  and the restricted representation  $\rho|_{(H'_{i_0} \times \prod_{i>i_0} H'_i, V')}$*

$$H'_{i_0} \times \prod_{i>i_0} H'_i \longrightarrow GL(V')$$

is a nontrivial tensor, or

(2) the representation  $\rho'$  factors through

$$\rho' : \prod_{i=1}^k H'_i \longrightarrow \left( \prod_{i=1}^k GL_{m'_i}(\mathbb{R}) \right)^+ \longrightarrow GL_n^+(\mathbb{R}),$$

where the latter homomorphism is, up to conjugation, the canonical diagonal embedding, and  $\rho'(H'_i)$  restricts to a scalar representation on each  $GL_{m'_i}(\mathbb{R})$ , for  $i \neq j$ .

Moreover, if all  $m_i$  are even then either  $m'_i < m_i$  for some  $i$  or one can replace  $GL$  with  $GL^+$  everywhere.

The notation  $\left( \prod_{i=1}^k GL_{m'_i}(\mathbb{R}) \right)^+$  stands for the intersection of  $\prod_{i=1}^k GL_{m'_i}(\mathbb{R})$  with the positive determinant matrices.

*Proof.* We argue by induction on  $k$ . For  $k = 2$  the alternative is immediate from Lemma 6. Suppose  $k > 2$ . If Item (1) does not hold, it follows from Lemma 6 that, up to replacing the  $H_i$ 's by some finite index subgroups,  $V$  decomposes invariantly to  $V = V_1 \oplus V'_1$  where  $\rho(H_1)$  is scalar on  $V'_1$  and  $\rho(\prod_{i>1} H_i)$  is scalar on  $V_1$ . We now apply the induction hypothesis for  $\prod_{i>1} H_i$  restricted to  $V'_1$ .

Finally, in Case (2), since  $\sum m_i = n$ , either  $m'_i < m_i$  for some  $i$  or equality holds everywhere. In the later case, if all the  $m_i$ 's are even, given  $g \in H_i$ , since the restriction of  $\rho(g)$  each  $V_{j \neq i}$  is scalar, it has positive determinant. We deduce that also  $\rho(g)|_{V_i}$  has positive determinant.  $\square$

### 3. MULTIPLICATIVITY OF THE MILNOR-WOOD CONSTANT FOR PRODUCT MANIFOLDS – A PROOF OF THEOREM 1

Let  $M_1, M_2$  be two arbitrary manifolds. We prove that

$$MW(M_1 \times M_2) = MW(M_1) \cdot MW(M_2).$$

First note that the inequality  $MW(M_1 \times M_2) \geq MW(M_1) \cdot MW(M_2)$  is trivial. Indeed, let  $\xi_1, \xi_2$  be flat oriented bundles over  $M_1$  and  $M_2$  respectively of the right dimension such that  $|\chi(\xi_i)| = MW(M_i) \cdot |\chi(M_i)|$  for  $i = 1, 2$ . Then  $\xi_1 \times \xi_2$  is a flat bundle over  $M_1 \times M_2$  with

$$|\chi(\xi_1 \times \xi_2)| = |\chi(\xi_1)| |\chi(\xi_2)| = MW(M_1) \cdot MW(M_2) \cdot |\chi(M_1 \times M_2)|.$$

For the other inequality, let  $\xi$  be a flat oriented  $\mathbb{R}^n$ -bundle over  $M_1 \times M_2$ , where  $n = \text{Dim}(M_1) + \text{Dim}(M_2)$ . We need to show that

$$|\chi(\xi)| \leq MW(X_1) \cdot MW(X_2) \cdot |\chi(M)|.$$

Observe that if we replace  $M$  by a finite cover, and the bundle  $\xi$  by its pullback to the cover, then both sides of the previous inequality are multiplied by the degree of the covering.

The flat bundle  $\xi$  is induced by a representation

$$\rho : \pi_1(M_1 \times M_2) \cong \pi_1(M_1) \times \pi_1(M_2) \longrightarrow \mathrm{GL}_n^+(\mathbb{R}).$$

If  $\rho(\pi_1(M_i))$  is amenable for  $i = 1$  or  $2$ , then  $\rho^*(\varepsilon_n) = 0$  [BuGe11, Lemma 4.3] and hence  $\chi(\xi) = 0$  and there is nothing to prove. Thus, we can without loss of generality suppose that, upon replacing  $\Gamma$  by a finite index subgroup the representation  $\rho$  factors as in Proposition 7.

In case (1) of the proposition, we obtain that  $\rho^*(\varepsilon_n) = 0$  by Lemma 10 and [BuGe11, Lemma 4.2]. In case (2) we get that  $\rho$  factors through

$$\rho : \pi_1(M_1) \times \pi_1(M_2) \longrightarrow \left( \mathrm{GL}_{m'_1}(\mathbb{R}) \times \mathrm{GL}_{m'_2}(\mathbb{R}) \right)^+ \xrightarrow{i} \mathrm{GL}_n^+(\mathbb{R}),$$

where the latter embedding  $i$  is up to conjugation the canonical embedding. Furthermore, up to replacing  $\rho$  by a representation in the same connected component of

$$\mathrm{Rep}(\pi_1(M_1) \times \pi_1(M_2), \left( \mathrm{GL}_{m'_1}(\mathbb{R}) \times \mathrm{GL}_{m'_2}(\mathbb{R}) \right)^+)$$

which will have no influence on the pullback of the Euler class, we can without loss of generality suppose that the scalar representations of  $\pi_1(M_1)$  on  $\mathrm{GL}_{m'_2}$  and  $\pi_1(M_2)$  on  $\mathrm{GL}_{m'_1}$  are trivial, so that  $\rho$  is a product representation. If  $m'_1$  or  $m'_2$  is odd, then  $i^*(\varepsilon_n) = 0 \in H_c^n((\mathrm{GL}_{m'_1}(\mathbb{R}) \times \mathrm{GL}_{m'_2}(\mathbb{R}))^+)$ . If  $m'_1$  and  $m'_2$  are both even then Proposition 7 further tells us that either  $m'_i < m_i$  for  $i = 1$  or  $2$ , or the image of  $\rho$  lies in  $\mathrm{GL}_{m_1}^+(\mathbb{R}) \times \mathrm{GL}_{m_2}^+(\mathbb{R})$ . In the first case, the Euler class vanishes [BuGe11, Lemma 4.2], while in the second case, we immediately obtain the desired inequality. This finishes the proof of Theorem 1.

#### 4. MULTIPLICATIVITY OF THE UNIVERSAL MILNOR-WOOD CONSTANT FOR HADAMARD MANIFOLDS - A PROOF OF THEOREM 2

Theorem 2 can be reformulated as follows:

**Theorem 8.** *Let  $X$  be a Hadamard manifold with de-Rham decomposition  $X = \prod_{i=1}^k X_i$ , then  $\widetilde{MW}(X) = \prod_{i=1}^k \widetilde{MW}(X_i)$ .*

We shall now prove Theorem 8. Note that the inequality " $\geq$ " is obvious. Let  $M = \Gamma \backslash X$  be a compact  $X$ -manifold. We must show that  $\mathrm{MW}(M) \leq \prod_{i=1}^k \widetilde{MW}(X_i)$ . Note that  $\Gamma$  is torsion free. Let us also assume that  $k \geq 2$ . If  $M$  is reducible one can argue by induction using Theorem 1. Thus we may assume that  $M$  is irreducible. Observe that this implies that  $\mathrm{Isom}(X)$  is not discrete. If  $\Gamma$  admits a nontrivial normal abelian subgroup then by the flat torus theorem (see [BH99, Ch. 7])  $X$  admits an Euclidian factor

which implies the vanishing of the Euler class. Assuming that this is not the case we apply the Farb–Weinberger theorem [FaWe08, Theorem 1.3] to deduce that  $X$  is a symmetric space of non-compact type. Thus, up to replacing  $M$  by a finite cover (equivalently, replace  $\Gamma$  by a finite index subgroup), we may assume that  $\Gamma$  lies in  $G = \text{Isom}(X)^\circ = \prod_{i=1}^k \text{Isom}(X_i)^\circ = \prod_{i=1}^k G_i$  and  $G$  is an adjoint semisimple Lie group without compact factors and  $\Gamma \leq G$  is irreducible in the sense that its projection to each factor is dense. Denote by  $\tilde{G}_i$  the universal cover of  $G_i$ , and by  $\tilde{\Gamma} \leq \prod_{i=1}^k \tilde{G}_i$  the pullback of  $\Gamma$ .

Let  $\rho : \Gamma \rightarrow \text{GL}_n^+(\mathbb{R})$  be a representation inducing a flat oriented vector bundle  $\xi$  over  $M$ . Up to replacing  $\Gamma$  by a finite index subgroup, we may suppose that  $\rho(\Gamma)$  is Zariski connected. Let  $S \leq \text{GL}_n^+(\mathbb{R})$  be the semisimple part of the Zariski closure of  $\rho(\Gamma)$ , and let  $\rho' : \Gamma \rightarrow S$  be the quotient representation. By superrigidity, the map  $\text{Ad} \circ \rho' : \Gamma \rightarrow \text{Ad}(S)$  extends to  $\phi : \Gamma \leq \prod_{i=1}^k G_i \rightarrow \text{Ad}(S)$  (see [Ma91, Mo06, GKM08]). This map can be pulled to  $\tilde{\phi} : \tilde{\Gamma} \rightarrow S$ . Recall also that  $\prod_{i=1}^k \tilde{G}_i$  is a central discrete extension of  $\prod_{i=1}^k G_i$  and, likewise,  $\tilde{\Gamma}$  is a central extension of  $\Gamma$ . If  $n_i = \dim X_i$  and  $n = \sum_{i=1}^k n_i$  we deduce from Proposition 7 and Lemma 10 that either the Euler class vanishes or the image of  $\tilde{\phi}$  lies (up to decomposing the vector space  $\mathbb{R}^n$  properly) in  $(\prod_{i=1}^k \text{GL}_{n_i})^+$ .

Suppose that  $\text{MW}(X_i)$  is finite for all  $i = 1, \dots, k$  and let  $M_i$  be closed  $X_i$ -manifolds. Let  $\xi'$  be the flat vector bundle on  $\prod_{i=1}^k M_i$  coming from  $\tilde{\rho}$  reduced to  $\prod_{i=1}^k M_i$ , and let  $\xi'_i$  be the vector bundle on  $M_i$  induced by  $\tilde{\rho}_i$ ,  $i = 1, \dots, k$ . By Lemma 9, we have

$$\frac{\chi(\xi)}{\text{vol}(M)} = \frac{\chi(\xi')}{\text{vol}(\prod_{i=1}^k M_i)} = \prod_{i=1}^k \frac{\chi(\xi'_i)}{\text{vol}(M_i)} \leq \prod_{i=1}^k \text{MW}(X_i),$$

which finishes the proof of Theorem 8.  $\square$

##### 5. EXAMPLE: A FLAT BUNDLE WITH NONZERO EULER NUMBER OVER A MANIFOLD WITH ZERO EULER CHARACTERISTIC

Recall that given two closed manifolds of even dimension, the Euler characteristic of connected sums behaves as

$$\chi(M_1 \sharp M_2) = \chi(M_1) + \chi(M_2) - 2.$$

The idea is to find  $M = M_1 \sharp M_2$  such that  $M_1$  admits a flat bundle with nontrivial Euler number in turn inducing such a bundle on the connected sum, and to choose then  $M_2$  in such a way that the Euler characteristic of the connected sum vanishes. Take thus

$$M_1 = \Sigma_2 \times \Sigma_2, \quad M_2 = (S^1 \times S^3) \sharp (S^1 \times S^3) \quad \text{and} \quad M = M_1 \sharp M_2.$$

These manifolds have the following Euler characteristics:

$$\begin{aligned}\chi(M_1) &= 4, \\ \chi(M_2) &= 2\chi(S^1 \times S^3) - 2 = -2, \\ \chi(M) &= 0.\end{aligned}$$

Let  $\eta$  be a flat bundle over  $\Sigma_2$  with Euler number  $\chi(\eta) = 1$ . (Note that we know that such a bundle exists by [Mi58].) Let  $f : M \rightarrow M_1$  be a degree 1 map obtained by sending  $M_2$  to a point, and consider

$$\xi = f^*(\eta \times \eta).$$

Obviously, since  $\eta$  is flat, so is the product  $\eta \times \eta$  and its pullback by  $f$ . Moreover, the Euler number of  $\xi$  is

$$\chi(\xi) = \chi(\eta \times \eta) = 1.$$

Indeed, the Euler number of  $\eta \times \eta$  is the index of a generic section of the bundle, which we can choose to be nonzero on  $f(M_2)$ , so that we can pull it back to a generic section of  $\xi$  which will clearly have the same index as the initial section on  $\eta \times \eta$ .

## 6. PROPORTIONALITY PRINCIPLES AND VANISHING OF THE EULER CLASS OF TENSOR PRODUCTS

**Lemma 9.** *Let  $X$  be a simply connected Riemannian manifold,  $G = \text{Isom}(M)$  and  $\rho : G \rightarrow \text{GL}_n^+(\mathbb{R})$  a representation. Then  $\frac{\chi(\xi_\rho)}{\text{vol}(M)}$ , where  $M = \Gamma \backslash X$  is a closed  $X$ -manifold and  $\xi_\rho$  is the flat vector bundle induced on  $M$  by  $\rho$  restricted to  $\Gamma$ , is a constant independent of  $M$ .*

*Proof.* There is a canonical isomorphism  $H_c^*(G) \cong H^*(\Omega^*(X)^G)$  between the continuous cohomology of  $G$  and the cohomology of the cocomplex of  $G$ -invariant differential forms  $\Omega^*(X)^G$  on  $X$  equipped with its standard differential. (For  $G$  a semisimple Lie group, every  $G$ -invariant form is closed, hence one further has  $H^*(\Omega^*(X)^G) \cong \Omega^*(X)^G$ .) In particular, in top dimension  $n = \dim(X)$ , the cohomology groups are 1-dimensional  $H_c^n(G) \cong H^n(\Omega^*(X)^G) \cong \mathbb{R}$  and contain the cohomology class given by the volume form  $\omega_X$ .

Since the bundle  $\xi_\rho$  over  $M$  is induced by  $\rho$ , its Euler class  $\varepsilon_n(\xi_\rho)$  is the image of  $\varepsilon_n \in H_c^n(\text{GL}^+(\mathbb{R}, n))$  under

$$H_c^n(\text{GL}^+(\mathbb{R}, n)) \xrightarrow{\rho^*} H_c^n(G) \longrightarrow H^n(\Gamma) \cong H^n(M),$$

where the middle map is induced by the inclusion  $\Gamma \hookrightarrow G$ . In particular,  $\rho^*(\varepsilon_n) = \lambda \cdot [\omega_X] \in H_c^n(G)$  for some  $\lambda \in \mathbb{R}$  independent of  $M$ . It follows that  $\chi(\xi_\rho)/\text{Vol}(M) = \lambda$ .  $\square$



**Lemma 10.** *Let  $\rho_{\otimes} : GL^+(n, \mathbb{R}) \times GL^+(m, \mathbb{R}) \rightarrow GL^+(nm, \mathbb{R})$  denote the tensor representation. If  $n, m \geq 2$ , then*

$$\rho_{\otimes}^*(\varepsilon_{nm}) = 0 \in H_c^{nm}(GL(n, \mathbb{R}) \times GL(m, \mathbb{R})).$$

*Proof.* The case  $n = m = 2$  was proven in [BuGe11, Lemma 4.1], based on the simple observation that interchanging the two  $GL^+(2, \mathbb{R})$  factors does not change the sign of the top dimensional cohomology class in  $H_c^4(GL(2, \mathbb{R}) \times GL(2, \mathbb{R})) \cong \mathbb{R}$ , but it changes the orientation on the tensor product, and hence the sign of the Euler class in  $H_c^4(GL^+(4, \mathbb{R}))$ .

Let us now suppose that at least one of  $n, m$  is strictly greater than 2, or equivalently, that  $n + m < nm$ . The Euler class is in the image of the natural map

$$H^{nm}(BGL(nm, \mathbb{R})) \longrightarrow H_c^{nm}(GL(nm, \mathbb{R})).$$

By naturality, we have a commutative diagram

$$\begin{array}{ccc} H^{nm}(BGL^+(nm, \mathbb{R})) & \longrightarrow & H_c^{nm}(GL^+(nm, \mathbb{R})) \\ \downarrow \rho_{\otimes}^* & & \downarrow \rho_{\otimes}^* \\ H^{nm}(B(GL^+(n, \mathbb{R}) \times GL^+(m, \mathbb{R}))) & \longrightarrow & H_c^{nm}(GL^+(n, \mathbb{R}) \times GL^+(m, \mathbb{R}))). \end{array}$$

Since the image of the lower horizontal arrow is contained in degree  $\leq n + m$ , it follows that  $\rho_{\otimes}^*(\varepsilon_{nm}) = 0$ .  $\square$

#### REFERENCES

- [Be00] Y. Benoist, *Tores affines in Crystallographic groups and their generalizations (Kortrijk, 1999)*, Contemp. Math. 262 (2000) 1–37.
- [BH99] R. Bridson, A. Haefliger, *Metric Spaces of Non-Positive Curvature*, Springer, 1999.
- [BuGe08] M. Bucher, T. Gelander. *Milnor–wood inequalities for locally  $(\mathbb{H}^2)^n$ -manifolds*. C. R. Acad. Sci. Paris 346 (2008), no. 11-12, 661-666.
- [BuGe11] M. Bucher, T. Gelander. *The generalized Chern conjecture for manifolds that are locally a product of surfaces*. Advances of Math., in press.
- [FaWe08] B. Farb, S. Weinberger, *Isometries, rigidity and universal covers*. Ann. of Math. (2) 168 (2008), no. 3, 915–940.
- [GKM08] T. Gelander, A. Karlsson, G.A. Margulis, *Superrigidity, generalized harmonic maps and uniformly convex spaces*. Geom. Funct. Anal. 17 (2008), no. 5, 1524–1550.
- [Ma91] G.A. Margulis, *Discrete Subgroups of Semisimple Lie Groups*, Ergeb. Math., Springer-Verlag, 1991.
- [Mi58] J. Milnor, *On the existence of a connection with curvature zero*. Comment. Math. Helv. 32, 1958, 215-223.
- [Mo06] N. Monod, *Superrigidity for irreducible lattices and geometric splitting*. J. Amer. Math. Soc. 19 (2006), no. 4, 781–814.
- [Sm77] J. Smillie, *Flat manifolds with non-zero Euler characteristics*, Comment. Math. Helv. 52 (1977) 453–355.