

THE ROYAL  
SWEDISH  
ACADEMY OF  
SCIENCES



**INSTITUT  
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden  
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89  
info@mittag-leffler.se www.mittag-leffler.se

**Sub-Riemannian geometry of Stiefel  
manifolds**

C. Autenried and I. Markina

REPORT No. 23, 2012/2013, fall

ISSN 1103-467X

ISRN IML-R- -23-12/13- -SE+fall

# SUB-RIEMANNIAN GEOMETRY OF STIEFEL MANIFOLDS

CHRISTIAN AUTENRIED, IRINA MARKINA

ABSTRACT. In the paper we consider the Stiefel manifold  $V_{n,k}$  as a principal  $U(k)$ -bundle over the Grassmann manifold and study the cut locus from the unit element. We gave the complete description of this cut locus on  $V_{n,1}$  and presented the sufficient condition on the general case.

## 1. INTRODUCTION

A sub-Riemannian geometry is an abstract setting for study geometry with non-holonomic constraints. A sub-Riemannian manifold is a triplet  $(Q, D, g_D)$ , where  $Q$  is a  $C^\infty$ -smooth manifold,  $D$  is a smooth sub-bundle of the tangent bundle  $TQ$  of the manifold  $Q$  (or smooth distribution) and  $g_D$  is a smoothly varying with respect to  $q \in Q$  inner product  $g_D(q): D_q \times D_q \rightarrow \mathbb{R}$ . The topic is actively developed last decades and as, now classical, sources we refer to [1, 11, 20, 24, 28].

One of the main objects for interest are normal and abnormal geodesics that are two different but not mutually disjoint families. The exponential map is not a local diffeomorphism anymore. Nevertheless, the singularities of the exponential map, as in the Riemannian geometry are closely related to the cut locus and failure of the optimality for geodesics. The cut locus in sub-Riemannian geometry is a quite few studied object. There exist very few results concerning the global and local structure of it and the results are mostly restricted to low dimensional manifolds. The work [25] studies the one dimensional Heisenberg group, and the results easily can be extended to higher dimensions. A full description of the global structure of the cut locus for the groups  $SU(2)$ ,  $SO(3)$ ,  $SL(2)$ , and lens spaces is given in [10]. For the groups  $SO(3)$ ,  $SL(2)$ , and lens spaces the cut locus is a stratified set, whereas in  $SU(2)$  it is a maximal circle without one point. The reader will find similar structures to those that obtained in the present work. The global structure of the exponential map and the cut locus of the identity on the group  $SE(2)$  is completely presented in [27].

The nature of normal and abnormal geodesics and complexity of the cut locus structure in sub-Riemannian geometry on the example of the Martinet manifold is pointed out in the work [4]. The Martinet manifold is the smooth manifold  $\mathbb{R}^3$  with smooth distribution spanned by vector fields

$$X = \frac{\partial}{\partial x} + \frac{1}{2}y^2 \frac{\partial}{\partial t}, \quad Y = \frac{\partial}{\partial y}$$

---

2010 *Mathematics Subject Classification.* 53C17, 52C30, 53C22.

*Key words and phrases.* Sub-Riemannian geometry, normal geodesic, cut locus, Stiefel manifolds, Grassmann manifold.

The authors are partially supported by the NFR-FRINAT grants #204726/V30 and #213440/BG..

and an inner product, making  $X, Y$  orthonormal. The cut locus is the Martinet surface  $y = 0$  minus the abnormal geodesic  $z = 0$  inside of the surface [4][Theorem 1.2]. The cut locus for contact manifolds were also studied in [5].

A progress in study of the cut locus of the identity on the sub-Lorentzian counterpart of one dimensional Heisenberg group can be found in [17].

In the present work we consider the Stiefel manifold  $V_{n,k}$  as a principal  $U(k)$ -bundle with the Grassmann manifold as a base space. We completely describe the cut locus from the unit element for the case  $V_{n,1}$ . The technical difficulties and possible presents of abnormal geodesics did not allowed to extend this result to the general case  $V_{n,k}$ . Nevertheless, we present a partial description of the cut locus, that is to our knowledge almost unique example for manifolds of higher dimensions.

The structure of the work is the following. Section 2 collects the basic definitions that nowadays are standard in sub-Riemannian geometry, but sometimes fussy. In Section 3 we define Grassmann and Stiefel manifolds, metric of constant bi-invariant type and normal geodesics based on the general theorem that can be found in [24]. In Section 4 we describe the cut locus for the equivalence class of the unit element on the principal  $U(1)$ -bundle structure on the Stiefel manifold  $V_{n,1}$ . Since the considered manifold is homogeneous it gives the structure of the cut locus for any point. The last Section 5 is dedicate to the cut locus for the general case of the Stiefel manifold  $V_{n,k}$ .

## 2. BASIC DEFINITIONS FROM SUB-RIEMANNIAN GEOMETRY

We remind the necessary definitions and propositions based on [24].

**Definition 1.** *A sub-Riemannian manifold is a triplet  $(Q, \mathcal{H}, \langle \cdot, \cdot \rangle)$ , where  $Q$  is a  $C^\infty$ -manifold,  $\mathcal{H}$  is a vector subbundle of the tangent bundle  $TQ$ , and  $\langle \cdot, \cdot \rangle$  is a fibre inner-product. The subbundle  $\mathcal{H}$  is called horizontal and  $\mathcal{H}_q$  is a horizontal space at a point  $q \in Q$ . The metric  $\langle \cdot, \cdot \rangle_q: \mathcal{H}_q \times \mathcal{H}_q \rightarrow \mathbb{R}$ ,  $q \in Q$  is called a sub-Riemannian metric, and the couple  $(\mathcal{H}, \langle \cdot, \cdot \rangle)$  is a sub-Riemannian structure on  $Q$ .*

**Definition 2.** *The horizontal subbundle  $\mathcal{H}$  is called bracket generating if for every  $q \in Q$  there exists  $r(q) \in \mathbb{Z}^+$  s.t.*

$$\mathcal{H}^{r(q)} = T_q Q,$$

where  $\mathcal{H}^1 := \mathcal{H}$  and  $\mathcal{H}^{r+1} := [\mathcal{H}^r, \mathcal{H}] + \mathcal{H}^r$ ,  $r \geq 1$ .

**Definition 3.** *An absolutely continuous curve  $\gamma: [0, T] \rightarrow Q$  is called horizontal if  $\dot{\gamma}(t) \in \mathcal{H}_{\gamma(t)}$  almost everywhere.*

**Definition 4.** *We define the length  $l := l(\gamma)$  of an absolutely continuous horizontal curve  $\gamma: [0, T] \rightarrow Q$  as in the Riemannian geometry:*

$$l(\gamma) := \int_0^T \|\dot{\gamma}\| dt = \int_0^T \sqrt{\langle \dot{\gamma}(t), \dot{\gamma}(t) \rangle} dt.$$

Introduce the function  $d(q_0, q)$  for  $q_0, q \in Q$  by

$$d(q_0, q) := \inf_{\gamma} \{l(\gamma)\},$$

where the infimum is taken over all absolutely continuous horizontal curves that connect  $q_0$  and  $q$ . If there is no horizontal curve joining  $q_0$  to  $q$ , then we declare  $d(q_0, q) = \infty$ .

Recall the Chow-Rashevskii theorem [12, 26] that gives a sufficient condition of the existence of horizontal curves.

**Theorem 1.** *Let  $Q$  be a connected manifold. If the horizontal subbundle  $\mathcal{H} \subset TQ$  is bracket generating, then any two points in  $Q$  can be joined by a horizontal curve.*

It follows that if  $\mathcal{H}$  is bracket generating on a connected manifold, then the function  $d$  introduced in Definition 3 is finite and defines the distance between two points on the manifold, called Carnot-Carathéodory distance.

**Definition 5.** *An absolutely continuous horizontal curve that realizes the distance between two points is called a minimizing geodesic.*

Let  $Q$  be  $n$ -dimensional smooth manifold and  $\mathcal{H}$  be a smooth horizontal subbundle such that  $\dim \mathcal{H}_q = k \leq n$  for all  $q \in Q$ . Considering a neighborhood  $U_q$  around  $q \in Q$  such that the subbundle  $\mathcal{H}$  is trivialized in  $U_q$ , one can find a local orthonormal basis  $X_1, \dots, X_k$  with respect to the sub-Riemannian metric  $\langle \cdot, \cdot \rangle$ . The associated sub-Riemannian metric Hamiltonian is given by

$$H(p, \lambda) = \frac{1}{2} \sum_{m=1}^k \lambda(X_m(p))^2,$$

where  $(p, \lambda) \in T^*U_q$ . A *normal geodesic* is defined as the projection to  $U_q \subset Q$  of the solution to the Hamiltonian system

$$\dot{p}_i = \frac{\partial H}{\partial \lambda_i} \quad \dot{\lambda}_i = -\frac{\partial H}{\partial p_i},$$

where  $(p_i, \lambda_i)$  are the coordinates in  $T^*U_q$ . We note that the word “normal” appears due to the fact that in the sub-Riemannian geometry there is another type of geodesics, calling “abnormal” arising from different type of Hamiltonian function. For a more detailed examination of abnormal geodesics we refer to [2, 3, 9, 23, 20]. The present work is mostly concerned with the normal geodesics, therefore we omit the detailed definition for abnormal ones.

Suppose two differentiable manifolds  $Q, M$ , and the submersion  $\pi: Q \rightarrow M$  are given. The fibre through  $q \in Q$  is the set  $Q_m := \pi^{-1}(m)$ ,  $m = \pi(q)$ , which is a submanifold according to the implicit function theorem. The differential  $d_q\pi: T_qQ \rightarrow T_{\pi(q)}M$  of  $\pi$  defines the vertical space  $\mathcal{V}_q \subset T_qQ$ . The vertical space at  $q$  is the tangent space to the fibre  $Q_{\pi(q)}$  and it can be written as  $\mathcal{V}_q := \ker(d_q\pi) = T_q(Q_m)$ , where  $\ker(d_q\pi)$  denotes the kernel of the linear map  $d_q\pi$ . It can be shown that  $\mathcal{V} = \bigcup_{q \in Q} \mathcal{V}_q$  is a smooth subbundle of  $TQ$  that is called vertical subbundle [24].

**Definition 6.** *An Ehresmann connection (or connection) for a submersion  $\pi: Q \rightarrow M$  is a subbundle  $\mathcal{H} \subset TQ$  that is everywhere transverse and of complementary dimension to the vertical:  $\mathcal{V}_q \oplus \mathcal{H}_q = T_qQ$ . The space  $\mathcal{H}_q$  is called horizontal subspace of  $T_qQ$ .*

**Definition 7.** *Let  $\pi: Q \rightarrow M$  be a submersion with connection  $\mathcal{H}$  and let  $c: I \rightarrow M$  be a curve starting at  $m \in M$ . A curve  $\gamma: I \rightarrow Q$  is called a horizontal lift of the curve  $c$  if  $\gamma$  is tangent to  $\mathcal{H}$  and projects to  $c$ , i.e.  $\dot{\gamma}(t) \in \mathcal{H}_{\gamma(t)}$  and  $\pi \circ \gamma(t) = c(t)$  for all  $t \in I$ .*

There are different ways to introduce a subriemannian structure on  $Q$ . In the sequel we describe two of them and indicate when they coincide.

Assuming that  $Q$  is a Riemannian manifold in the submersion  $\pi: Q \rightarrow M$ , we can use its Riemannian metric to define the orthogonal complement  $\mathcal{H}_q$  of the vertical space  $\mathcal{V}_q$  at each point  $q \in Q$ . Then  $\mathcal{H}$  is a connection and the restriction of the Riemannian metric to  $\mathcal{H}$  defines a sub-Riemannian metric on  $Q$ .

Assume that the manifold  $M$  is endowed with a Riemannian metric and the submersion  $\pi: Q \rightarrow M$  has a connection  $\mathcal{H}$ . Since  $\mathcal{V}_q = \ker(d_q\pi)$  and  $\text{Im}(d_q\pi|_{\mathcal{H}_q}) = \text{Im}(d_q\pi) = T_{\pi(q)}M$ , it follows that  $d_q\pi|_{\mathcal{H}_q}$  is a linear isomorphism from  $\mathcal{H}_q$  to  $T_{\pi(q)}M$ . By pulling back the Riemannian metric on  $M$  to  $Q$ , we obtain a sub-Riemannian metric on  $Q$  with underlying subbundle  $\mathcal{H}$ . This sub-Riemannian metric is said to be induced by the connection  $\mathcal{H}$  on  $Q$  and the Riemannian metric on  $M$ .

Suppose  $Q$  and  $M$  are smooth Riemannian manifolds and a submersion  $\pi: Q \rightarrow M$  is given. Let  $\mathcal{H}_q$  be orthogonal complement to the vertical  $\mathcal{V}_q$  at every  $q \in Q$ . Two ways of inducing a sub-Riemannian metric on  $Q$ , by restricting the Riemannian metric of  $Q$  or by pulling back the Riemannian metric on  $M$  using  $d\pi$ , coincide if  $d_q\pi$  restricts to a linear isometry  $\mathcal{H}_q \rightarrow T_{\pi(q)}M$  for all  $q \in Q$ .

**Definition 8.** *Let  $Q$  and  $M$  be Riemannian manifolds and let  $\pi: Q \rightarrow M$  be a submersion. Let  $\mathcal{V}_q \subset T_qQ$  denote the vertical subspace at  $q \in Q$  and  $\mathcal{H}_q := \mathcal{V}_q^\perp$  be its orthogonal complement. If  $d\pi: TQ \rightarrow TM$  restricts to a linear isometry  $\mathcal{H}_q \rightarrow T_{\pi(q)}M$  for each  $q \in Q$ , then  $\pi$  is called a Riemannian submersion.*

Thus, Riemannian metrics on  $Q$  and  $M$  induce the same subriemannian structure on  $Q$  if the submersion is Riemannian.

**Definition 9.** *A fibre bundle  $\pi: Q \rightarrow M$  is a principal  $G$ -bundle if its fibre is a Lie group  $G$  that acts freely and transitively on each fibre.*

As a consequence we can identify  $M$  with the quotient  $Q/G$  of  $Q$  by the group action of  $G$ . Furthermore,  $\pi$  corresponds to the canonical projection to the quotient.

**Definition 10.** *A connection on  $\pi: Q \rightarrow M$  is a principal bundle connection if the action of  $G$  preserves the connection.*

We assume that the group acts on itself on the right  $q \mapsto qg$ ,  $q \in Q$ ,  $g \in G$ .

**Definition 11.** *Let  $Q \rightarrow M$  be a principal  $G$ -bundle with connection  $\mathcal{H}$ . A sub-Riemannian metric on  $(Q, \mathcal{H})$  which is invariant under the action of  $G$  is called a metric of bundle type.*

A sub-Riemannian metric which is induced from a  $G$ -invariant metric on  $Q$  is an example of a metric of bundle type.

**Definition 12.** *A bi-invariant Riemannian metric  $\langle \cdot, \cdot \rangle$  on a differentiable manifold  $Q$  with the Lie group  $G$  acting on it is said to be of constant bi-invariant type if its inertia tensor  $\mathbb{I}_q: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$  defined by  $\mathbb{I}_q(\xi, \eta) := \langle \sigma_q\xi, \sigma_q\eta \rangle$  is independent of  $q \in Q$ . Here*

$$\begin{aligned} \sigma_q: \mathfrak{g} &\rightarrow T_qQ \\ \xi &\mapsto \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} q \exp(\epsilon\xi). \end{aligned}$$

**Definition 13.** *Let  $\pi: Q \rightarrow M$  be a principal  $G$ -bundle with a Riemannian metric of constant bi-invariant type and  $\mathcal{H}$  be the induced connection. We define the  $\mathfrak{g}$ -valued*

connection one-form  $A_q$  uniquely as the linear operator  $A_q: T_qQ \rightarrow \mathfrak{g}$  which satisfies following properties:

$$\ker(A_q) = \mathcal{H}_q, \quad A_q \circ \sigma_q = Id_{\mathfrak{g}},$$

where  $Id_{\mathfrak{g}}$  is the identity map on  $\mathfrak{g}$ .

The map  $A: TQ \rightarrow \mathfrak{g}$  defines a  $\mathfrak{g}$ -valued connection tensor on  $Q$ .

**Theorem 2.** [24] *Let  $\pi: Q \rightarrow M$  be a principal  $G$ -bundle with a Riemannian metric of constant bi-invariant type. Let  $\mathcal{H}$  be the induced connection, with  $\mathfrak{g}$ -valued connection tensor  $A$ . Let  $\exp_R$  be the Riemannian exponential map, so that  $\gamma_R(t) = \exp_R(tv)$  is the Riemannian geodesic through  $q$  with initial velocity  $v \in T_qQ$ . Then any horizontal lift  $\gamma$  of the projection  $\pi \circ \gamma_R$  is a normal subriemannian geodesic and is given by*

$$\gamma(t) = \exp_R(tv) \exp_G(-tA(v)),$$

where  $\exp_G: \mathfrak{g} \rightarrow G$  is the group  $G$  exponential map. Moreover, all normal subriemannian geodesics can be obtained in this way.

### 3. STIEFEL AND GRASSMANN MANIFOLDS

We use the following notations in the present section. Let  $\mathbb{C}^n$  denote a  $n$ -dimensional complex vector space and  $\mathbb{C}^{m \times n}$  the set of  $(m \times n)$ -matrices with complex entries. We want to apply Theorem 2 for the submersion  $\pi: V_{n,k}(\mathbb{C}^n) \rightarrow G_{n,k}(\mathbb{C}^n)$ , where  $V_{n,k}(\mathbb{C}^n) = V_{n,k}$  is the Stiefel manifold and  $G_{n,k}(\mathbb{C}^n) = G_{n,k}$  is the Grassmann manifold for  $n \in \mathbb{N}$  and  $k \in \{1, \dots, n\}$ .

We start from the description of a general construction. Given a group  $G$  with an invariant inner product on its Lie algebra  $\mathfrak{g}$  and two subgroups  $H, K \subset G$ , we form the quotient spaces  $G/H$  and  $G/(H \times K)$ . The submersion  $G/H \rightarrow G/(H \times K)$  is a principal  $K$ -bundle, with Riemannian metrics on  $G/H$  and  $G/(H \times K)$  induced from the bi-invariant Riemannian metric on  $G$  generated by an invariant inner product. The Riemannian metrics are induced by the projections  $G \rightarrow G/H$  and  $G \rightarrow G/(H \times K)$ . Both manifolds in the submersion  $G/H \rightarrow G/(H \times K)$  are homogeneous manifolds, where the group  $G$  acts. The geodesics on  $G/H$  are the projections from  $G$  of one-parameter subgroups  $\exp(t\xi)$  with  $\xi$  orthogonal to the Lie algebra  $\mathfrak{h} \subset \mathfrak{g}$  of  $H$ . We set  $G = U(n)$ ,  $H = U_n(n-k)$ ,  $K = U_n(k)$ , where

$$U_n(k) := \left\{ \begin{pmatrix} U_k & 0 \\ 0 & I_{n-k} \end{pmatrix} \mid U_k \in U(k) \right\} \subset U(n) \quad \text{and}$$

$$U_n(n-k) := \left\{ \begin{pmatrix} I_k & 0 \\ 0 & U_{n-k} \end{pmatrix} \mid U_{n-k} \in U(n-k) \right\} \subset U(n).$$

Note that we use the notations  $U_n(k)$  and  $U_n(n-k)$  in the current section to emphasize that the elements of these groups are written as  $(n \times n)$ -matrices. Then the quotient  $G/H$  is isomorphic to the Stiefel manifold  $V_{n,k}$  and  $G/(H \times K)$  is isomorphic to the Grassmann manifold  $G_{n,k}$ .

**3.1. Unitary group and bi-invariant metric.** Before we give detailed definition of Stiefel and Grassmann manifolds we remind that the unitary group  $U(n)$  is a matrix Lie group, whose elements  $X$  satisfy the condition

$$U(n) = \{X \in \mathbb{C}^{n \times n} \mid \bar{X}^T X = X \bar{X}^T = I_n\}.$$

Here  $I_n$  is the unite  $(n \times n)$ -matrix and  $X^T$  is the transposed of the matrix  $X$ . The Lie algebra  $\mathfrak{u}(n)$  consists of all skew-Hermitian matrices:

$$\mathfrak{u}(n) = \{\mathcal{X} \in \mathbb{C}^{n \times n} \mid \mathcal{X} = -\bar{\mathcal{X}}^T\}.$$

We remind that a matrix  $X \in U(n)$  is of full rank, its columns and rows are orthonormal to each other with respect to the standard Hermitian product in  $\mathbb{C}^n$  and that the diagonal of the skew-Hermitian matrices are purely imaginary. Moreover, the Hermitian product in  $\mathbb{C}^n$  is invariant under the action of  $U(n)$ , that particularly means that the orthogonality is preserved under this action. The Lie algebra  $\mathfrak{u}(n)$  can be endowed with the inner product  $-2n \operatorname{tr}(\mathcal{X}\mathcal{Y})$ ,  $\mathcal{X}, \mathcal{Y} \in \mathfrak{u}(n)$ . Considering  $U(n)$  as a smooth manifold, we denote its points by  $q$  and the metric at this point by  $\langle \cdot, \cdot \rangle_{U(n)}(q)$  or if it is clear from the context simply by  $g_q$ . Then a left-invariant metric on  $U(n)$  with respect to the group action of  $U(n)$  on its Lie algebra is given by

$$\begin{aligned} \langle \cdot, \cdot \rangle_{U(n)}(q): \quad T_q U(n) \times T_q U(n) &\cong \mathfrak{qu}(n) \times \mathfrak{qu}(n) &\rightarrow \mathbb{R} \\ (q\mathcal{X}, q\mathcal{Y}) &\mapsto \langle q\mathcal{X}, q\mathcal{Y} \rangle_{U(n)}(q) := -2n \operatorname{tr}(\mathcal{X}\mathcal{Y}) \end{aligned}$$

$q \in U(n)$ ,  $\mathcal{X}, \mathcal{Y} \in \mathfrak{u}(n)$ . This metric is actually bi-invariant, that follows from the observation that can be found, for instance, in [15] and [22]. Let  $\mathfrak{g}$  be a Lie algebra of a Lie group  $G$  endowed with an inner product  $(\cdot, \cdot)_{\mathfrak{g}}$ . An inner product  $(\cdot, \cdot)_{\mathfrak{g}}$  is called invariant if it is invariant under the adjoint action of  $G$ , i.e.  $(q^{-1}\eta q, q^{-1}\xi q)_{\mathfrak{g}} = (\eta, \xi)_{\mathfrak{g}}$  for all  $\eta, \xi \in \mathfrak{g}$  and  $q \in G$ . Then it is well known, see for instance [19], that an invariant inner product  $(\cdot, \cdot)_{\mathfrak{g}}$  on a Lie algebra  $\mathfrak{g}$  determines a bi-invariant metric on the group  $G$  via

$$\langle \eta, \xi \rangle_G(q) := (q^{-1}\eta, q^{-1}\xi)_{\mathfrak{g}} = (\eta q^{-1}, \xi q^{-1})_{\mathfrak{g}}$$

for all  $\eta, \xi \in T_q G$ .

We only need to check that the inner product  $(\mathcal{X}, \mathcal{Y})_{\mathfrak{u}(n)} = -2n \operatorname{tr}(\mathcal{X}\mathcal{Y})$  on  $\mathfrak{u}(n)$  is invariant. Indeed,

$$\begin{aligned} (q^{-1}\mathcal{X}q, q^{-1}\mathcal{Y}q)_{\mathfrak{u}(n)} &= -2n \operatorname{tr}(q^{-1}\mathcal{X}q q^{-1}\mathcal{Y}q) = -2n \operatorname{tr}(q^{-1}\mathcal{X}\mathcal{Y}q) \\ &= -2n \operatorname{tr}(\mathcal{Y}q q^{-1}\mathcal{X}) = -2n \operatorname{tr}(\mathcal{X}\mathcal{Y}) = (\mathcal{X}, \mathcal{Y})_{\mathfrak{u}(n)} \end{aligned}$$

for all  $\mathcal{X}, \mathcal{Y} \in \mathfrak{u}(n)$  and  $q \in U(n)$ .

**Remark 1.** *The left and right action of any subgroup  $U_n(k)$ ,  $1 \leq k \leq n$  on the group  $U(n)$  and on the Lie algebra  $\mathfrak{u}(n)$  are defined as a matrix multiplication from the left or from the right. The inner product  $(\cdot, \cdot)_{\mathfrak{g}} = -2n \operatorname{tr}(\cdot, \cdot)$  on the Lie algebra  $\mathfrak{u}(n)$  is invariant under the adjoint action of  $U_n(k)$  and therefore the metric  $\langle \cdot, \cdot \rangle_{U(n)}$ , defined by left or right translations by the action of  $U_n(k)$ , is bi-invariant under this action.*

**3.2. Stiefel manifold and metric of constant bi-invariant type.** The Stiefel manifold  $V_{n,k}$  is the set of all  $k$ -tuples  $(q_1, \dots, q_k)$  of vectors  $q_i \in \mathbb{C}^n$ ,  $i \in \{1, \dots, k\}$ , which are orthonormal with respect to the standard Hermitian metric. This is a compact manifold which can be equivalently defined as

$$V_{n,k} := \{X \in \mathbb{C}^{n \times k} \mid \bar{X}^T X = I_k\}.$$

The condition  $\bar{X}^T X = I_k$  is equivalent to the orthonormality of columns. This  $k$  orthonormal columns can be considered as arbitrary  $k$  columns in a matrix  $X \in U(n)$ .

This allows us to realize the Stiefel manifold as a quotient set of  $U(n)$  by the group  $U_n(n-k)$ . To do this we introduce the equivalence relation  $\sim_1$  on  $U(n)$  by

$$q \sim_1 p \Leftrightarrow q = p \begin{pmatrix} I_k & 0 \\ 0 & U_{n-k} \end{pmatrix}, \quad q, p \in U(n), \quad U_{n-k} \in U(n-k).$$

This results to the equivalence class for  $q \in U(n)$

$$[q]^{\sim_1} = \left\{ q \begin{pmatrix} I_k & 0 \\ 0 & U_{n-k} \end{pmatrix}, \mid U_{n-k} \in U(n-k) \right\} \in U(n)/U_n(n-k), \quad q \in U(n).$$

The quotient  $U(n)/U_n(n-k)$  is a smooth manifold with the quotient topology and we denote the natural projection from  $U(n)$  to the quotient  $U(n)/U_n(n-k)$  by  $\pi_1$ . We identify the equivalence class  $[q]^{\sim_1}$  with a point in the Stiefel manifold and write  $[q]_{V_{n,k}} \in V_{n,k}$  instead of  $[q]^{\sim_1}$  to emphasize that point belongs to the Stiefel manifold. So, practically, an element of  $V_{n,k}$  is thought of an element in  $U(n)$  whose first  $k$  columns from the left are of interest and the last  $n-k$  columns are not. The real dimension of  $V_{n,k}$  is  $2nk - k^2$ .

The tangent space to the Stiefel manifold is a quotient of the tangent space to  $U(n)$  by tangent space of the equivalence classes. To obtain it we differentiate curves  $\gamma(t) \in [q]^{\sim_1}$  at  $t = 0$  for a fixed  $q \in U(n)$  and get the space  $\mathcal{R} = \left\{ q \begin{pmatrix} 0 & 0 \\ 0 & \mathcal{C} \end{pmatrix} \mid \mathcal{C} \in \mathfrak{u}(n-k) \right\}$ . Intuitively, movements in the direction  $\mathcal{R}$  make no change in the quotient space. It follows that the tangent space  $T_{[q]_{V_{n,k}}} V_{n,k}$  to the Stiefel manifold at  $[q]_{V_{n,k}} \in V_{n,k}$  is given by the quotient of the tangent space  $T_q U(n)$ , that is isomorphic to  $q\mathfrak{u}(n)$ , by  $\mathcal{R}$ :

$$T_{[q]_{V_{n,k}}} V_{n,k} = \left\{ [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix} \mid \mathcal{X}_1 \in \mathfrak{u}(k), \mathcal{X}_2 \in \mathbb{C}^{(n-k) \times k} \right\}.$$

Similar results can be find in [6] or [21].

Now we define a metric  $\langle \cdot, \cdot \rangle_{V_{n,k}}$  on  $V_{n,k}$  by

$$\begin{aligned} & \left\langle [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, [q]_{V_{n,k}} \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{V_{n,k}} \left( [q]_{V_{n,k}} \right) \\ := & \left\langle q \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, q \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{U(n)} (q) = \left( \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right)_{\mathfrak{u}(n)}, \end{aligned}$$

where  $q \in [q]_{V_{n,k}}$  is any representative of the equivalence class  $[q]_{V_{n,k}}$ .

Since  $U_k[q]_{V_{n,k}} = [U_k q]_{V_{n,k}}$  and  $[q]_{V_{n,k}} U_k = [q U_k]_{V_{n,k}}$ ,  $U_k \in U_n(k)$ , it follows directly from the definition of the metric on  $T_{[q]_{V_{n,k}}} V_{n,k}$  and the bi-invariance of the metric  $\langle \cdot, \cdot \rangle_{U(n)}$  with respect to  $U_n(k)$  that

$$\begin{aligned} & \left\langle [U_k q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, [U_k q]_{V_{n,k}} \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{V_{n,k}} \\ = & \left( \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right)_{\mathfrak{u}(n)} \\ = & \left\langle [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, [q]_{V_{n,k}} \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{V_{n,k}} \end{aligned}$$



and

$$\begin{aligned}
& \left\langle [qU_k]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, [qU_k]_{V_{n,k}} \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{V_{n,k}} \\
&= \left( \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right)_{\mathfrak{u}(n)} \\
&= \left\langle [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & -\bar{\mathcal{X}}_2^T \\ \mathcal{X}_2 & 0 \end{pmatrix}, [q]_{V_{n,k}} \begin{pmatrix} \mathcal{Y}_1 & -\bar{\mathcal{Y}}_2^T \\ \mathcal{Y}_2 & 0 \end{pmatrix} \right\rangle_{V_{n,k}},
\end{aligned}$$

where  $U_k$  is any element in  $U_n(k) \subset U(n)$ .

Now we show that the metric  $\langle \cdot, \cdot \rangle_{V_{n,k}}$  on  $V_{n,k}$  is of constant bi-invariant type with respect to the right group action of  $U_n(k)$ . To prove it we recall that the infinitesimal generator  $\sigma_{[q]_{V_{n,k}}} : \mathfrak{u}_n(k) \rightarrow T_{[q]_{V_{n,k}}} V_{n,k}$  is given by  $\sigma_{[q]_{V_{n,k}}}(\xi) = [q]_{V_{n,k}} \xi$ , where  $\mathfrak{u}_n(k) = \left\{ \begin{pmatrix} u_k & 0 \\ 0 & 0 \end{pmatrix} \mid u_k \in \mathfrak{u}(k) \right\}$  is the Lie algebra of  $U_n(k)$ . It follows that

$$\mathbb{I}_{[q]_{V_{n,k}}}(\xi, \eta) = \langle [q]_{V_{n,k}} \xi, [q]_{V_{n,k}} \eta \rangle_{V_{n,k}} = -2n \operatorname{tr}(\xi \eta), \quad \text{where } [q]_{V_{n,k}} \in V_{n,k}.$$

This implies that  $\mathbb{I}_{[q]_{V_{n,k}}}(\xi, \eta)$  is independent of  $[q]_{V_{n,k}}$ . We will use the notation  $g$  for the metric  $\langle \cdot, \cdot \rangle_{V_{n,k}}$  whenever it does not produce confusions.

**3.3. Grassmann manifold.** The Grassmann manifold  $G_{n,k}$  is defined as a collection of all  $k$ -dimensional subspaces  $\Lambda$  of  $\mathbb{C}^n$ . Equivalently, an element  $\Lambda$  of  $G_{n,k}$  can be written as a  $(n \times k)$  matrix with columns  $e_1, \dots, e_k$ , such that  $\operatorname{span}(e_1, \dots, e_k) = \Lambda$ . We are interested in the representation of  $G_{n,k}$  as a quotient of  $U(n)$  by some subgroup. As in the previous case of the Stiefel manifold, we divide  $U(n)$  by  $U_n(n-k)$ , but moreover, since the definition of  $G_{n,k}$  does not depend on the choice of the orthonormal basis  $e_1, \dots, e_k$  for  $\Lambda$ , but only on its span, we also quotient  $U(n)$  by the group  $U_n(k)$  that leaves  $\operatorname{span}\{e_1, \dots, e_k\}$  invariant. Therefore, we define the equivalence relation  $\sim_2$  in  $U(n)$  by

$$m_1 \sim_2 m_2 \quad :\iff \quad m_1 = m_2 \begin{pmatrix} U_k & 0 \\ 0 & U_{n-k} \end{pmatrix}, \quad m_1, m_2 \in U(n),$$

where  $U_k \in U(k)$ ,  $U_{n-k} \in U(n-k)$ . This leads to the equivalence class

$$[m]^{\sim_2} = \left\{ m \begin{pmatrix} U_k & 0 \\ 0 & U_{n-k} \end{pmatrix} \mid U_k \in U(k), U_{n-k} \in U(n-k) \right\} \subset U(n), \quad m \in U(n),$$

which is isomorphic to  $U(k) \times U(n-k) \cong U_n(k) \times U_n(n-k)$ . We identify  $G_{n,k}$  with the quotient space  $U(n)/(U_n(k) \times U_n(n-k))$  and use the notation  $[m]_{G_{n,k}}$  for  $[m]^{\sim_2}$  in the present Section 3.

Furthermore, we obtain that the tangent space to the equivalence class  $[m]^{\sim_2}$  is

$$\left\{ m \begin{pmatrix} \mathcal{X}_1 & 0 \\ 0 & \mathcal{X}_4 \end{pmatrix} \mid \mathcal{X}_1 \in \mathfrak{u}(k), \mathcal{X}_4 \in \mathfrak{u}(n-k) \right\}, \quad m \in U(n),$$

and it implies that the tangent space of  $G_{n,k}$  at the point  $[m]_{G_{n,k}}$  is given by

$$T_{[m]_{G_{n,k}}} G_{n,k} = \left\{ [m]_{G_{n,k}} \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \mid \mathcal{X}_2 \in \mathbb{C}^{k \times (n-k)} \right\}.$$

It has real dimension  $2k(n-k)$  that gives the real dimension of  $G_{n,k}$ , see also [6, 21].

We define a metric  $\langle \cdot, \cdot \rangle_{G_{n,k}}$  on  $G_{n,k}$  by

$$\begin{aligned} & \left\langle [m]_{G_{n,k}} \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix}, [m]_{G_{n,k}} \begin{pmatrix} 0 & \mathcal{Y}_2 \\ -\bar{\mathcal{Y}}_2^T & 0 \end{pmatrix} \right\rangle_{G_{n,k}} \left( [m]_{G_{n,k}} \right) \\ & := \left\langle m \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix}, m \begin{pmatrix} 0 & \mathcal{Y}_2 \\ -\bar{\mathcal{Y}}_2^T & 0 \end{pmatrix} \right\rangle_{U(n)} (m) \\ & = \left( \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix}, \begin{pmatrix} 0 & \mathcal{Y}_2 \\ -\bar{\mathcal{Y}}_2^T & 0 \end{pmatrix} \right)_{\mathfrak{u}(k)}, \end{aligned}$$

where  $m \in U(n)$  is any representative of  $[m]_{G_{n,k}}$ .

**3.4. Submersion  $\pi: V_{n,k} \rightarrow G_{n,k}$  and sub-Riemannian geodesics.** Now we have all necessary tools to define the submersion

$$\begin{aligned} \pi: V_{n,k} & \rightarrow G_{n,k}, \\ [q]_{V_{n,k}} & \mapsto [m]_{G_{n,k}}. \end{aligned}$$

The projection  $\pi$  sends the equivalence class  $[q]^{\sim 1}$  to the equivalence class  $[m]^{\sim 2}$ , where  $m \in U(n)$  can be any matrix from the set

$$\left\{ q \begin{pmatrix} U_k & 0 \\ 0 & U_{n-k} \end{pmatrix} \mid U_k \in U(k), U_{n-k} \in U(n-k) \right\}.$$

Note that the latter set consists of all unitary matrices whose first  $k$  columns from the left span the same space as the first left  $k$  columns of  $q$ . This implies that a fibre over a point  $[m]_{G_{n,k}} \in G_{n,k}$  is given by

$$\begin{aligned} \pi^{-1}([m]_{G_{n,k}}) & = \left\{ \left[ m \begin{pmatrix} U_k & 0 \\ 0 & I_{n-k} \end{pmatrix} \right]_{V_{n,k}} \mid U_k \in U(k) \right\} \\ & = \left\{ [m]_{V_{n,k}} \begin{pmatrix} U_k & 0 \\ 0 & I_{n-k} \end{pmatrix} \mid U_k \in U(k) \right\}, \quad m \in U(n), \end{aligned}$$

which is homeomorphic to  $U_n(k) \cong U(k)$ .

The submersion  $\pi$  is also a principal  $U_n(k)$ -bundle, where the right group action is defined by the multiplication from the right by an element from  $U_n(k)$ . It remains to show that the right action of  $U_n(k)$  is continuous, preserves the fibre, acts freely and transitively on the fibre.

The multiplication of  $[q]_{V_{n,k}} \in V_{n,k}$  from the right by an element  $U_k^0 \in U(k)$  is given by

$$q \begin{pmatrix} I_k & 0 \\ 0 & U_{n-k} \end{pmatrix} \begin{pmatrix} U_k^0 & 0 \\ 0 & I_{n-k} \end{pmatrix} = q \begin{pmatrix} U_k^0 & 0 \\ 0 & U_{n-k} \end{pmatrix}, \quad q \in U(n),$$

where  $U_{n-k}$  is an arbitrary element of  $U(n-k)$  and  $U_k^0$  is a fixed element of  $U(k)$ . It follows that the right multiplication is well defined and continuous. It can also be seen, that it preserves the fibre of  $\pi^{-1}(\pi([q]_{V_{n,k}}))$ . By the definition of the fibre it is clear that  $[q]_{V_{n,k}} U(k) = \pi^{-1}(\pi([q]_{V_{n,k}}))$  and this implies the transitivity of the  $U_n(k)$  action.

To show that  $U_n(k)$  acts freely, we assume that  $\tilde{U}_1 = \begin{pmatrix} U_1 & 0 \\ 0 & I_{n-k} \end{pmatrix} \in U_n(k)$ ,  $\tilde{U}_2 = \begin{pmatrix} U_2 & 0 \\ 0 & I_{n-k} \end{pmatrix} \in U_n(k)$  and  $[q]_{V_{n,k}} \tilde{U}_1 = [q]_{V_{n,k}} \tilde{U}_2$  with  $[q]_{V_{n,k}} = \begin{pmatrix} q_1 & q_2 \\ q_3 & q_4 \end{pmatrix}$ . Then we get the equations

$$\begin{aligned} q_1 U_1 = q_1 U_2 &\iff q_1 = q_1 U_2 U_1^{-1} = q_1 U_1 U_2^{-1} \\ q_3 U_1 = q_3 U_2 &\iff q_3 = q_3 U_2 U_1^{-1} = q_3 U_1 U_2^{-1}, \end{aligned}$$

that leads to  $U_1 = U_2$  and so  $\tilde{U}_1 = \tilde{U}_2$ . Thus, we showed that  $\pi: V_{n,k} \rightarrow G_{n,k}$  is a principal  $U_n(k)$ -bundle.

The differential of  $\pi$  defines the vertical and horizontal spaces. The differential  $d_{[q]_{V_{n,k}}} \pi$  at  $[q]_{V_{n,k}}$  acts as

$$[q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \mapsto [m]_{G_{n,k}} \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix}.$$

So, the kernel of  $d_{[q]_{V_{n,k}}} \pi$  or the vertical space  $\mathcal{V}_{[q]_{V_{n,k}}}$  is given by

$$\mathcal{V}_{[q]_{V_{n,k}}} = \left\{ [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & 0 \\ 0 & 0 \end{pmatrix} \mid \mathcal{X}_1 \in \mathfrak{u}(k) \right\}, \quad q \in U(n).$$

We choose the horizontal space of  $V_{n,k}$  by setting

$$(1) \quad \mathcal{H}_{[q]_{V_{n,k}}} = \left\{ [q]_{V_{n,k}} \begin{pmatrix} 0 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \mid \mathcal{X}_2 \in \mathbb{C}^{k \times (n-k)} \right\}, \quad q \in U(n).$$

It is clear that  $d\pi: TV_{n,k} \rightarrow TG_{n,k}$  is a linear isometry if we restrict it to the horizontal space,  $\mathcal{H}_{[q]_{V_{n,k}}} \rightarrow T_{\pi([q]_{V_{n,k}})} G_{n,k}$  for each  $[q]_{V_{n,k}} \in V_{n,k}$ , therefore  $\pi$  is a Riemannian submersion.

The  $\mathfrak{u}_n(k)$ -valued connection one-form  $A_{[q]_{V_{n,k}}}: T_{[q]_{V_{n,k}}} V_{n,k} \rightarrow \mathfrak{u}_n(k)$  is given by

$$A_{[q]_{V_{n,k}}} \left( [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \right) := \begin{pmatrix} \mathcal{X}_1 & 0 \\ 0 & 0 \end{pmatrix} \in \mathfrak{u}_n(k), \quad \mathcal{X}_2 \in \mathbb{C}^{k \times (n-k)}.$$

Now we can write precisely the normal sub-Riemannian geodesic on  $V_{n,k}$  starting from  $[q]_{V_{n,k}}$ . It is given by

$$\begin{aligned} (2) \quad \gamma(t) &= \exp_R(tv) \exp_{U_n(k)}(-tA(v)) \\ &= \pi_1 \left( q \exp_{U(n)} \left( t \begin{pmatrix} \mathcal{X}_1 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \right) \right) \exp_{U_n(k)} \left( -t \begin{pmatrix} \mathcal{X}_1 & 0 \\ 0 & 0 \end{pmatrix} \right) \end{aligned}$$

where  $q \in U(n)$ ,  $v = [q]_{V_{n,k}} \begin{pmatrix} \mathcal{X}_1 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \in T_{[q]_{V_{n,k}}} V_{n,k}$  with  $\begin{pmatrix} \mathcal{X}_1 & \mathcal{X}_2 \\ -\bar{\mathcal{X}}_2^T & 0 \end{pmatrix} \in \mathfrak{u}(n)$ .

We simplify the notation and from now on write  $q \in V_{n,k}$ ,  $m \in G_{n,k}$ ,  $U(k)$  for  $U_n(k)$ ,  $U(n-k)$  for  $U_n(n-k)$ , and  $g$  for the Riemannian metric of constant bi-invariant type.

#### 4. THE CUT-LOCUS OF $V_{n,1}$

In this section we study the cut locus of the Stiefel manifold  $V_{n,1}$  considered as a sub-Riemannian manifold by making use of the normal sub-Riemannian geodesics (2). Recall the definition of the cut locus.

**Definition 14.** *The cut locus of  $q_0 \in Q$  is the set  $K_{q_0}$  of points reached optimally by more than one geodesic, i. e. the cut locus is*

$$K_{q_0} := \left\{ q \in Q \mid \text{there exist } T \in \mathbb{R}^+, v_1, v_2 \in T_{q_0}Q, v_1 \neq v_2, \text{ and} \right. \\ \left. \text{minimizing geodesics } \gamma_{v_1}(t), \gamma_{v_2}(t), \text{ starting from } q_0, \text{ and} \right. \\ \left. \gamma_{v_1}(T) = \gamma_{v_2}(T) = q \right\}.$$

The main theorem is stated as following.

**Theorem 3.** *The cut locus  $K_{Id}$  on  $V_{n,1}$  is given by*

$$L := \left\{ \left[ \begin{pmatrix} C & 0 \\ 0 & D \end{pmatrix} \right]_{V_{n,k}} \mid C \in U(1), D \in U(n-1) \right\} \setminus \left\{ [Id]_{V_{n,k}} \right\}.$$

Before we present the proof of Theorem 3 we consider in details the particular case for  $n = 2, k = 1$ . It allows to understand the general idea of the proof without using tough technical calculations.

**4.1. The cut locus of  $V_{2,1}$ .** Observe that  $V_{2,1}$  is three dimensional and the distribution (1) is strongly bracket generating. Recall the definition.

**Definition 15.** *A smooth distribution  $\mathcal{H}$  on  $M$  is strongly bracket generating if and only if for any non-zero section  $\mathcal{Z}$  of  $\mathcal{H}$ , the tangent bundle  $TM$  is generated by  $\mathcal{H}$  and  $[\mathcal{Z}, \mathcal{H}]$ .*

For the manifold  $V_{2,1}$  it is reduced to the statement that there exist two sections  $\mathcal{Z}_1$  and  $\mathcal{Z}_2$  of  $\mathcal{H}$  such that  $\text{span}\{\mathcal{Z}_1(q), \mathcal{Z}_2(q), [\mathcal{Z}_1, \mathcal{Z}_2](q)\} = T_q V_{2,1}$  for all  $q \in V_{2,1}$ . Namely,  $\mathcal{Z}_1(q) := q \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  and  $\mathcal{Z}_2(q) := q \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ . It is known, see for instance [10, 24], that on sub-Riemannian manifolds with strongly bracket generating distributions all minimizing geodesics are normal.

Recall that the identity  $\text{Id}$  on  $V_{2,1}$  and  $G_{2,1}$  is an equivalence class containing the unit element of  $U(n)$  and taken with respect to the corresponding equivalence relation. The tangent spaces at the identity are given by

$$T_{\text{Id}}V_{2,1} = \left\{ \text{Id} \begin{pmatrix} x_1 & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \mid x_1 = \lambda i, \lambda \in \mathbb{R}, x_2 \in \mathbb{C} \right\}$$

and

$$T_{\text{Id}}Gr_{2,1} = \left\{ \text{Id} \begin{pmatrix} 0 & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \mid x_2 \in \mathbb{C} \right\}.$$

For a given initial vector  $v = \text{Id} \begin{pmatrix} x_1 & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \in T_{\text{Id}}V_{2,1}$  a normal sub-Riemannian geodesic is written as

$$\begin{aligned} \gamma_v(t) &= \pi_1(\exp_{U(2)}(tv)) \exp_{U(1)} \left( -t \begin{pmatrix} \lambda i & 0 \\ 0 & 0 \end{pmatrix} \right) \\ &= \pi_1 \left( \exp_{U(2)} \left( t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right) \right) \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \\ &= \pi_1 \left( \begin{pmatrix} \gamma_v^1(t) & \gamma_v^2(t) \\ \gamma_v^3(t) & \gamma_v^4(t) \end{pmatrix} \right) = \left[ \begin{pmatrix} \gamma_v^1(t) & \gamma_v^2(t) \\ \gamma_v^3(t) & \gamma_v^4(t) \end{pmatrix} \right]_{V_{n,k}} \end{aligned}$$

with

$$\begin{aligned}
\gamma_v^1(t) &= \left( \frac{\lambda}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} + \frac{1}{2} \right) \mu_1(-\lambda, x_2, t) + \left( -\frac{\lambda}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} + \frac{1}{2} \right) \mu_2(-\lambda, x_2, t), \\
\gamma_v^2(t) &= \frac{x_2 i}{\sqrt{\lambda^2 + 4x_2\bar{x}_2}} (\mu_2(\lambda, x_2, t) - \mu_1(\lambda, x_2, t)), \\
\gamma_v^3(t) &= \frac{i}{4x_2} \left( \frac{\lambda^2}{\sqrt{\lambda^2 + 4x_2\bar{x}_2}} - \sqrt{\lambda^2 + 4x_2\bar{x}_2} \right) (\mu_2(-\lambda, x_2, t) - \mu_1(-\lambda, x_2, t)) \\
&= -\frac{\bar{x}_2 i}{\sqrt{\lambda^2 + 4x_2\bar{x}_2}} (\mu_2(-\lambda, x_2, t) - \mu_1(-\lambda, x_2, t)), \\
\gamma_v^4(t) &= -\frac{\mu_1(\lambda, x_2, t)}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} (\lambda - \sqrt{\lambda^2 + 4x_2\bar{x}_2}) + \frac{\mu_2(\lambda, x_2, t)}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} (\lambda + \sqrt{\lambda^2 + 4x_2\bar{x}_2}),
\end{aligned}$$

where

$$\mu_1(\lambda, x_2, t) = e^{\frac{ti}{2}(\lambda + \sqrt{\lambda^2 + 4x_2\bar{x}_2})} \text{ and } \mu_2(\lambda, x_2, t) = e^{\frac{ti}{2}(\lambda - \sqrt{\lambda^2 + 4x_2\bar{x}_2})}.$$

In calculations we used the diagonalization of the matrix  $t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} = SDS^{-1}$  with

$$\begin{aligned}
S &= \begin{pmatrix} 1 & 1 \\ -\frac{i}{2x_2}(\lambda - \sqrt{\lambda^2 + 4x_2\bar{x}_2}) & -\frac{i}{2x_2}(\lambda + \sqrt{\lambda^2 + 4x_2\bar{x}_2}) \end{pmatrix}, \\
S^{-1} &= \begin{pmatrix} \frac{\lambda}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} + \frac{1}{2} & -\frac{x_2 i}{\sqrt{\lambda^2 + 4x_2\bar{x}_2}} \\ \frac{1}{2} - \frac{\lambda}{2\sqrt{\lambda^2 + 4x_2\bar{x}_2}} & \frac{x_2 i}{\sqrt{\lambda^2 + 4x_2\bar{x}_2}} \end{pmatrix}, \\
D &= \begin{pmatrix} it\left(\frac{\lambda + \sqrt{\lambda^2 + 4x_2\bar{x}_2}}{2}\right) & 0 \\ 0 & it\left(\frac{\lambda - \sqrt{\lambda^2 + 4x_2\bar{x}_2}}{2}\right) \end{pmatrix},
\end{aligned}$$

in order to express  $\exp_{U(2)} \left( t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right) = S \exp_{U(2)}(D) S^{-1}$ .

**Lemma 1.** *The set*

$$L := \left\{ \left[ \exp_{U(2)} \begin{pmatrix} c_1 i & 0 \\ 0 & c_2 i \end{pmatrix} \right]_{V_{2,1}} \mid c_1, c_2 \in \mathbb{R} \right\} \setminus \{\text{Id}\}$$

is the cut locus  $K_{\text{Id}}$  of  $V_{2,1}$ .

*Proof.* It is clear that it is enough to concentrate on the calculation of the first column  $\begin{pmatrix} \gamma_v^1(t) \\ \gamma_v^3(t) \end{pmatrix}$  in the equivalence class  $\left[ \begin{pmatrix} \gamma_v^1(t) & \gamma_v^2(t) \\ \gamma_v^3(t) & \gamma_v^4(t) \end{pmatrix} \right]_{V_{2,1}}$ . We show first that if  $q \in L$ , then there are several minimizing geodesics reaching  $q$  in the same time.

Suppose there exists an initial vector  $v^* = \begin{pmatrix} \lambda i & x_2^* \\ -\bar{x}_2^* & 0 \end{pmatrix}$  with  $x_2^* \neq 0$ , and  $T \in \mathbb{R}^+$  such that the minimizing geodesic  $\gamma_{v^*}$  connects  $\text{Id} = \left[ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right]_{V_{2,1}}$  with

$$q = \gamma_{v^*}(T^*) = \left[ \begin{pmatrix} e^{c_1 i} & 0 \\ 0 & e^{c_2 i} \end{pmatrix} \right]_{V_{2,1}} \in L.$$

We see that  $\gamma_{v^*}^2(T^*) = 0$ . It implies the following equivalences

$$(3) \quad \begin{aligned} & \mu_1(\lambda^*, x_2^*, T^*) = \mu_2(\lambda^*, x_2^*, T^*) \\ \iff & e^{\frac{T^* i}{2}(\lambda^* + \sqrt{(\lambda^*)^2 + 4|x_2^*|^2})} = e^{\frac{T^* i}{2}(\lambda^* - \sqrt{(\lambda^*)^2 + 4|x_2^*|^2})} \\ \iff & e^{\frac{T^* i}{2}\sqrt{(\lambda^*)^2 + 4|x_2^*|^2}} = e^{-\frac{T^* i}{2}\sqrt{(\lambda^*)^2 + 4|x_2^*|^2}} \\ \iff & \frac{T^*}{2}\sqrt{(\lambda^*)^2 + 4x_2^* \bar{x}_2^*} = k\pi, \quad \text{for some } k \in \mathbb{Z}. \end{aligned}$$

Let us fix such  $k \in \mathbb{Z}$  and note

$$\mu_1(\lambda^*, x_2^*, T^*) = e^{\frac{T^* i}{2}(\lambda^* + \sqrt{(\lambda^*)^2 + 4|x_2^*|^2})} = \pm e^{\frac{T^* i}{2}\lambda^*} = \mu_2(\lambda^*, x_2^*, T^*).$$

We conclude that functions  $\mu_1(\lambda^*, x_2^*, T^*)$  and  $\mu_2(\lambda^*, x_2^*, T^*)$  are independent of  $x_2$  itself, but depend on the norm  $|x_2^*|^2 = x_2^* \bar{x}_2^*$ . Let us pick up another initial velocity vector  $v_1 = \begin{pmatrix} \lambda^* i & y_2 \\ -\bar{y}_2 & 0 \end{pmatrix}$  with  $|y_2|^2 = |x_2^*|^2$  and  $x_2^* \neq y_2$ . Then  $\gamma_{v_1}(T^*) = q$ .

In the next step we show that the length of the geodesic  $\gamma_{v_1}$  coincides with the length of the minimizing geodesic  $\gamma_{v^*}$ . We actually claim that the length of any geodesic  $\gamma_v$  with  $v = \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix}$  depends on the fixed final time  $T$  and the norm  $|x_2|$ .

We recall that the square of the length of the velocity vector  $\dot{\gamma}_v(t)$  is given by

$$g(\dot{\gamma}_v(t), \dot{\gamma}_v(t)) = -2n \operatorname{tr}((\gamma_v(t)^{-1} \dot{\gamma}_v(t))^2).$$

Fix a point  $f(t) = \exp_{U(2)} \left( t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right) \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \in U(2)$  such that  $\gamma_v(t) = \pi_1(f(t))$ . To calculate  $\dot{\gamma}_v(t) = d_{f(t)} \pi_1 f'(t)$  we use the chain rule

$$\begin{aligned} \dot{\gamma}_v(t) &= d_{f(t)} \pi_1 \left[ \exp_{U(2)} \left\{ t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right\} \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \right. \\ &+ \left. \exp_{U(2)} \left\{ t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right\} \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -\lambda i & 0 \\ 0 & 0 \end{pmatrix} \right] \\ &= d_{f(t)} \pi_1 \left[ \exp_{U(2)} \left\{ t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right\} \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda i & x_2 e^{\lambda i t} \\ -\bar{x}_2 e^{-\lambda i t} & 0 \end{pmatrix} \right. \\ &+ \left. \exp_{U(2)} \left\{ t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right\} \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -\lambda t i & 0 \\ 0 & 0 \end{pmatrix} \right] \\ &= d_{f(t)} \pi_1 \left[ \exp_{U(2)} \left\{ t \begin{pmatrix} \lambda i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix} \right\} \begin{pmatrix} e^{-\lambda t i} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & x_2 e^{\lambda i t} \\ -\bar{x}_2 e^{-\lambda i t} & 0 \end{pmatrix} \right] \\ &= \gamma_v(t) \begin{pmatrix} 0 & x_2 e^{\lambda i t} \\ -\bar{x}_2 e^{-\lambda i t} & 0 \end{pmatrix}. \end{aligned}$$

It follows that  $g(\dot{\gamma}_v(t), \dot{\gamma}_v(t)) = -4 \operatorname{tr} \begin{pmatrix} -|x_2|^2 & 0 \\ 0 & -|x_2|^2 \end{pmatrix} = 8|x_2|^2$ . Since the length of  $\gamma_v$  depends only on  $T$  and the norm  $|x_2|$  we conclude that  $\gamma_v$  is a minimizing geodesic from the identity to  $q$ . With this we finished to show the inclusion  $L \subset K_{\operatorname{Id}}$ .

To prove the converse inclusion  $K_{\operatorname{Id}} \subset L$  we use a contradiction. Suppose  $q \in V_{2,1} \setminus L$ , but  $q \in K_{\operatorname{Id}}$ , i. e. there exist  $v_1 = \begin{pmatrix} \lambda_1 i & x_2 \\ -\bar{x}_2 & 0 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} \lambda_2 i & y_2 \\ -\bar{y}_2 & 0 \end{pmatrix} \in \mathfrak{u}(n)$  with  $v_1 \neq v_2$  such that  $\gamma_{v_1}$  and  $\gamma_{v_2}$  are minimizing geodesics from the identity to  $q$ , which reach  $q$  optimally at time  $T \in \mathbb{R}^+$ . Note that values  $x_2$  and  $y_2$  do not vanish as otherwise  $\gamma_{v_1}(t) = \gamma_{v_2}(t) = \operatorname{Id}$  for all  $t \in \mathbb{R}$ .

For any unitary matrix  $q = \begin{pmatrix} q_1 & q_2 \\ q_3 & q_4 \end{pmatrix}$  one obtains  $q_2 \neq 0 \Leftrightarrow q_3 \neq 0$ . It follows that if  $q \in V_{2,1} \setminus L$ , then

$$\gamma_{v_1}^2 \neq 0, \gamma_{v_1}^3 \neq 0 \iff \mu_2(\lambda, x_2, T) \neq \mu_1(\lambda, x_2, T) \iff \frac{T}{2} \sqrt{\lambda^2 + 4|x_2|^2} \notin \pi\mathbb{Z}$$

by (3). It immediately implies  $T < \min \left\{ \frac{2\pi}{\sqrt{\lambda_1^2 + 4|x_2|^2}}, \frac{2\pi}{\sqrt{\lambda_2^2 + 4|y_2|^2}} \right\}$ . In the next step we show that both inequalities can not be realized under our assumptions.

**Case 1.** Assume that  $|x_2|^2 \neq |y_2|^2$  and  $\lambda_1, \lambda_2$  are arbitrary. Then  $g(v_1, v_1) = 8|x_2|^2 \neq 8|y_2|^2 = g(v_2, v_2)$ , that implies that the length of both minimizing geodesics  $\gamma_{v_1}$  and  $\gamma_{v_2}$  is different, which is a contradiction to the assumption that they are both minimizing at the same time.

**Case 2.** Let  $|x_2|^2 = |y_2|^2$  and  $\lambda_1 = \lambda_2$ . As  $\gamma_{v_1}^3(T) = \gamma_{v_2}^3(T) \neq 0$  it follows that

$$\begin{aligned} \gamma_{v_1}^3(T) &= - \frac{\bar{x}_2 i}{\sqrt{\lambda_1^2 + 4|x_2|^2}} (\mu_2(\lambda_1, x_2, T) - \mu_1(\lambda_1, x_2, T)) \\ &= - \frac{\bar{y}_2 i}{\sqrt{\lambda_2^2 + 4|y_2|^2}} (\mu_2(\lambda_2, y_2, T) - \mu_1(\lambda_2, y_2, T)) = \gamma_{v_2}^3(T) \end{aligned}$$

if and only if  $x_2 = y_2$ ,

because  $\sqrt{\lambda_1^2 + 4|x_2|^2} = \sqrt{\lambda_2^2 + 4|y_2|^2}$ ,  $\mu_1(\lambda_1, x_2, T) = \mu_1(\lambda_2, y_2, T)$ , and  $\mu_2(\lambda_1, x_2, T) = \mu_2(\lambda_2, y_2, T)$  by definition. But the equality  $x_2 = y_2$  implies  $v_1 = v_2$ , that leads to a contradiction.

**Case 3.** Finally we suppose that  $\lambda_1 \neq \lambda_2$  and  $|x_2|^2 = |y_2|^2$ . As in the previous case the equality  $\gamma_{v_1}^3(T) = \gamma_{v_2}^3(T)$  implies  $|\gamma_{v_1}^3(T)| = |\gamma_{v_2}^3(T)|$  and

$$\frac{1}{\sqrt{\lambda_1^2 + 4|x_2|^2}} |\mu_2(\lambda_1, x_2, T) - \mu_1(\lambda_1, x_2, T)| = \frac{1}{\sqrt{\lambda_2^2 + 4|y_2|^2}} |\mu_2(\lambda_2, y_2, T) - \mu_1(\lambda_2, y_2, T)|.$$

Taking into account  $|\exp\left(\frac{T i \lambda_j}{2}\right)| = 1$  for  $j = 1, 2$ , we obtain

$$\begin{aligned} |\mu_2(\lambda_j, x_2, T) - \mu_1(\lambda_j, x_2, T)| &= |e^{-\frac{T i}{2} \sqrt{\lambda_j^2 + 4|x_2|^2}} - e^{\frac{T i}{2} \sqrt{\lambda_j^2 + 4|x_2|^2}}| \\ &= 2 \sin\left(\frac{T}{2} \sqrt{\lambda_j^2 + 4|x_2|^2}\right), \quad j = 1, 2, \end{aligned}$$

as  $\sin x > 0$  for  $x \in (0, \pi)$ . These both equations lead to

$$\begin{aligned} \frac{2 \sin\left(\frac{T}{2} \sqrt{\lambda_1^2 + 4|x_2|^2}\right)}{\sqrt{\lambda_1^2 + 4|x_2|^2}} &= \frac{2 \sin\left(\frac{T}{2} \sqrt{\lambda_2^2 + 4|y_2|^2}\right)}{\sqrt{\lambda_2^2 + 4|y_2|^2}} \\ \iff \frac{2 \sin\left(\frac{T}{2} \sqrt{\lambda_1^2 + 4|x_2|^2}\right)}{T \sqrt{\lambda_1^2 + 4|x_2|^2}} &= \frac{2 \sin\left(\frac{T}{2} \sqrt{\lambda_2^2 + 4|y_2|^2}\right)}{T \sqrt{\lambda_2^2 + 4|y_2|^2}}. \end{aligned}$$

Since the function  $\frac{\sin x}{x}$  is injective on the interval  $(0, \pi)$  we obtain  $\frac{T}{2} \sqrt{\lambda_1^2 + 4|x_2|^2} = \frac{T}{2} \sqrt{\lambda_2^2 + 4|y_2|^2}$  which is equivalent to  $\lambda_1 = \pm \lambda_2$ .

We only need to consider the case  $\lambda_1 = -\lambda_2$ . Note that  $\mu_j(\lambda_2, x_2, T) = \mu_j(\lambda_2, y_2, T)$ ,  $j = 1, 2$ , and

$$\mu_1(-\lambda_2, x_2, T) = \frac{1}{\mu_2(\lambda_2, y_2, T)}, \quad \mu_2(-\lambda_2, x_2, T) = \frac{1}{\mu_1(\lambda_2, y_2, T)}.$$

From this it follows that

$$(4) \quad \gamma_{v_1}^1(T) = \gamma_{v_2}^1(T) \iff \frac{\tan\left(\frac{T}{2} \sqrt{\lambda_2^2 + 4|y_2|^2}\right)}{\sqrt{\lambda_2^2 + 4|y_2|^2}} = \frac{\tan\left(\frac{T\lambda_2}{2}\right)}{\lambda_2}.$$

Since  $0 < \frac{T\lambda_2}{2} < \frac{T}{2} \sqrt{\lambda_2^2 + 4y_2^2} < \pi$  the equality (4) is not true.




FIGURE 1.  $\frac{\tan(x)}{x}$  on the interval  $[0, \pi]$

Figure 4.1 shows that  $y_1 < y_2$  implies  $\frac{\tan y_1}{y_1} \neq \frac{\tan y_2}{y_2}$ . The similar calculations can be found in [10, p. 1871].

These 3 cases finish the proof of the theorem.  $\square$

**4.2. Isomorphism between  $V_{2,1}$  and  $SU(2)$ .** In this subsection we show that the results obtained above recover the results obtained in [10]. An element  $q$  of  $V_{2,1}$  is an equivalence class which can be written as

$$[q]_{V_{2,1}} = \left\{ \begin{pmatrix} \alpha & \exp(\lambda i)\bar{\beta} \\ \beta & -\exp(\lambda i)\bar{\alpha} \end{pmatrix} \mid \lambda \in (0, 2\pi) \right\}.$$



Since  $\begin{pmatrix} \alpha & \exp(\lambda i)\bar{\beta} \\ \beta & -\exp(\lambda i)\bar{\alpha} \end{pmatrix}$  is a unitary matrix we know that the norm  $\|\alpha\|^2 + \|\beta\|^2$  of the vector  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$  is equal one. Furthermore, we note that points  $q \in V_{2,1}$  can be parametrized by the vector  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ . Recall the definition of the group  $SU(2) = \left\{ \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \mid \|\alpha\|^2 + \|\beta\|^2 = 1 \right\}$ . So, it is clear that every element of  $SU(2)$  can be represented by the vector  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ . It follows that the both manifolds are isomorphic by the mapping  $f: V_{2,1} \rightarrow SU(2)$ ,  $\begin{pmatrix} \alpha \\ \beta \end{pmatrix} \mapsto \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ .

**Remark 1.** *The set  $L$  as a subset of  $V_{2,1}$  depends only on  $c_1 \in (0, 2\pi)$ , since the part depending on  $c_2$  is quotient out. This implies that the cut locus of  $SU(2)$ , which is given by  $\left\{ \begin{pmatrix} \exp(c_1 i) & 0 \\ 0 & \exp(-c_1 i) \end{pmatrix} \mid c_1 \in (0, 2\pi) \right\}$  [10], has a bijective relation under the map  $f$  to the cut locus of  $V_{2,1}$ , given in Lemma 1.*

Now we proceed to the proof of Theorem 3 that describes the cut locus from the identity on  $V_{n,1}$ .

*Proof.* We only need to show the inclusion  $K_{Id} \subset L$  since the converse inclusion  $L \subset K_{Id}$  will be proved in Theorem 4 for any  $V_{n,k}$ .

First of all we claim that in the case of  $V_{n,1}$  there are no abnormal minimizing geodesics because the distribution is the strongly bracket generating. To show that the horizontal distribution is strongly bracket generating we consider an arbitrary element  $\begin{pmatrix} 0 & B \\ -\bar{B}^T & 0 \end{pmatrix} \in \mathcal{H}_{Id}$  with  $B = (b_1, \dots, b_{n-1}) \in \mathbb{C}^{1 \times (n-1)}$ . Take basis elements  $\begin{pmatrix} 0 & E_{mj} \\ -\bar{E}_{mj}^T & 0 \end{pmatrix}$ ,  $m \in \{0, 1\}$ , and  $j \in \{1, \dots, n-1\}$ . Here  $E_{mj} \in \mathbb{C}^{1 \times (n-1)}$  is the row with entry  $i^m$  at the place  $j$  and zeros everywhere else. Then, the commutator is written as

$$\left[ \begin{pmatrix} 0 & B \\ -\bar{B}^T & 0 \end{pmatrix}, \begin{pmatrix} 0 & E_{mj} \\ -\bar{E}_{mj}^T & 0 \end{pmatrix} \right] = \begin{pmatrix} -B\bar{E}_{mj}^T + E_{mj}\bar{B}^T & 0 \\ 0 & 0 \end{pmatrix},$$

where

$$-B\bar{E}_{mj}^T + E_{mj}\bar{B}^T = \begin{cases} -2i \operatorname{Im}(b_j), & m = 0 \\ -2i \operatorname{Re}(b_j), & m = 1 \end{cases}.$$

An arbitrary choice of  $B$  and linearity of Lie bracket imply that any  $B$  generates the whole vertical space which allows to conclude that the distribution  $\mathcal{H}$  is strongly bracket generating.

Now we calculate the precise form of a geodesic  $\gamma_v$ , concentrating on components  $\gamma_v^1$  and  $\gamma_v^3$ . Let the velocity vector be given by  $v = \begin{pmatrix} xi & B \\ -\bar{B}^T & 0 \end{pmatrix}$ , where  $x \in \mathbb{R}$  and  $B \in \mathbb{C}^{1 \times (n-1)}$ . Recall  $\exp(tv) = \sum_{n=0}^{\infty} \frac{t^n}{n!} v^n$ . First we will calculate the two parts of

$v^n := v(n)$ , namely  $v^1(n)$  and  $v^2(n)$ . From the recursion formula  $v^n = v^{n-1}v$  it follows that

$$v^1(n) = v^1(n-1)xi - v^2(n-1)\bar{B}^T = v^1(n-1)xi - v^1(n-2)B\bar{B}^T$$

as  $v^2(n) = v^1(n-1)B$ . Furthermore, as  $v^n = vv^{n-1}$  we deduce

$$v^3(n) = -\bar{B}^T v^1(n-1).$$

Having the initial values  $v^1(0) = 1$ ,  $v^1(1) = xi$ , and  $v^3(0) = 0$  we obtain by using the program **wolframalpha** that

$$\begin{aligned} v^1(n) &= \frac{2^{-n-1}}{i\sqrt{x^2+4B\bar{B}^T}} \left( ix((i\sqrt{x^2+4B\bar{B}^T}+ix)^n - (ix-i\sqrt{x^2+4B\bar{B}^T})^n) \right. \\ &\quad \left. + i\sqrt{x^2+4B\bar{B}^T}((ix-i\sqrt{x^2+4B\bar{B}^T})^n + (i\sqrt{x^2+4B\bar{B}^T}+ix)^n) \right) \end{aligned}$$

and

$$\begin{aligned} \exp(tv)^1 &= \sum_{n=0}^{\infty} \frac{t^n}{n!} v^1(n) \\ &= \frac{1}{2i\sqrt{x^2+4B\bar{B}^T}} \left( e^{-\frac{it}{2}(\sqrt{x^2+4B\bar{B}^T}-x)} \right. \\ &\quad \left. \times \left( i\sqrt{x^2+4B\bar{B}^T}(e^{it\sqrt{x^2+4B\bar{B}^T}}+1) + ix(e^{it\sqrt{x^2+4B\bar{B}^T}}-1) \right) \right) \\ &= \frac{1}{2\sqrt{x^2+4B\bar{B}^T}} \left( e^{-\frac{it}{2}(\sqrt{x^2+4B\bar{B}^T}-x)} \right. \\ &\quad \left. \times \left( \sqrt{x^2+4B\bar{B}^T}(e^{it\sqrt{x^2+4B\bar{B}^T}}+1) + x(e^{it\sqrt{x^2+4B\bar{B}^T}}-1) \right) \right). \end{aligned}$$

The first part of the normal geodesic  $\gamma_v^1(t)$  is written as

$$\begin{aligned} \gamma_v^1(t) &= \exp_{U(n)}(tv)^1 \exp_{U(1)}(-tix) \\ &= \frac{1}{2\sqrt{x^2+4B\bar{B}^T}} e^{-\frac{it}{2}(\sqrt{x^2+4B\bar{B}^T}+x)} \\ &\quad \times \left( \sqrt{x^2+4B\bar{B}^T}(e^{it\sqrt{x^2+4B\bar{B}^T}}+1) + x(e^{it\sqrt{x^2+4B\bar{B}^T}}-1) \right), \end{aligned}$$

that coincides with calculations in the case  $n = 2$ .

The second important part of the geodesic is

$$\begin{aligned} \exp(tv)^3 &= \sum_{n=0}^{\infty} \frac{t^n}{n!} v^3(n) = \sum_{n=1}^{\infty} \frac{t^n}{n!} v^3(n) \\ &= \sum_{n=0}^{\infty} \frac{t^{n+1}}{(n+1)!} v^3(n+1) = \sum_{n=0}^{\infty} \frac{t^{n+1}}{(n+1)!} (-\bar{B}^T v^1(n)) \\ &= -\bar{B}^T \frac{1}{i\sqrt{x^2+4B\bar{B}^T}} e^{-\frac{ti}{2}(\sqrt{x^2+4B\bar{B}^T}-x)} \left( e^{ti\sqrt{x^2+4B\bar{B}^T}} - 1 \right), \end{aligned}$$

$$\begin{aligned} \gamma_v^3(t) &= \exp(tv)^3 \exp(-tix) \\ &= -\bar{B}^T \frac{1}{i\sqrt{x^2+4B\bar{B}^T}} e^{-\frac{ti}{2}(\sqrt{x^2+4B\bar{B}^T}+x)} \left( e^{ti\sqrt{x^2+4B\bar{B}^T}} - 1 \right). \end{aligned}$$

It follows that  $\gamma_v^3(t) = 0$  for  $t = \frac{2\pi}{\sqrt{x^2+4B\bar{B}^T}}$ . That means that the geodesic reaches the set  $L$  at  $t = \frac{2\pi}{\sqrt{x^2+4B\bar{B}^T}}$ , and since  $L \subset K_{Id}$  it is also reaches the cut locus. This implies that the geodesic loses its optimality at the latest  $t = \frac{2\pi}{\sqrt{x^2+4B\bar{B}^T}}$ .

Having exact formulas for coordinates of geodesics we proceed to the core of the proof. Suppose  $q \in V_{n,1} \setminus L$  but  $q \in K_{Id}$ , and there exist two optimal normal geodesics  $\gamma_{v_1}$  and  $\gamma_{v_2}$  with  $\gamma_{v_1}(0) = \gamma_{v_2}(0) = Id$ ,  $\gamma_{v_1}(T^*) = \gamma_{v_2}(T^*) = q$  and  $v_1 = \begin{pmatrix} x_1 i & B \\ -\bar{B}^T & 0 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} x_2 i & E \\ -\bar{E}^T & 0 \end{pmatrix}$  such that  $v_1 \neq v_2$  and  $x_j \in \mathbb{R}$ ,  $j = 1, 2$  and  $B, E \in \mathbb{C}^{1 \times (n-1)}$ .

Further we argue mostly as in the proof of Lemma 1.

**Case 1.** Assume  $B\bar{B}^T \neq E\bar{E}^T$ . Since the length of both geodesics  $\gamma_{v_1}$  and  $\gamma_{v_2}$  should coincide, it can be shown, as in the proof of Proposition 3, that

$$\|B\|^2 = B\bar{B}^T = \text{tr}(B\bar{B}^T) = \text{tr}(E\bar{E}^T) = E\bar{E}^T = \|E\|^2.$$

This is a contradiction.

**Case 2.** Let  $x_1 = x_2$  and  $\|B\|^2 = \|E\|^2$ . It follows from

$$\begin{aligned} \gamma_{v_1}^3(T^*) &= \gamma_{v_2}^3(T^*) && \iff \\ &- \bar{B}^T \frac{1}{i\sqrt{x_1^2 + 4B\bar{B}^T}} e^{-\frac{iT}{2}(\sqrt{x_1^2 + 4B\bar{B}^T} + x_1)} (e^{iT\sqrt{x_1^2 + 4B\bar{B}^T}} - 1) \\ &= - \bar{E}^T \frac{1}{i\sqrt{x_2^2 + 4E\bar{E}^T}} e^{-\frac{iT}{2}(\sqrt{x_2^2 + 4E\bar{E}^T} + x_2)} (e^{iT\sqrt{x_2^2 + 4E\bar{E}^T}} - 1) \end{aligned}$$

that  $\bar{B}^T = \bar{E}^T$  and so  $B = E$ , that leads to the contradiction with  $v_1 \neq v_2$ .

**Case 3.** Let now  $x_1 \neq x_2$  and  $\|B\|^2 = \|E\|^2$ . We know that  $\gamma_{v_1}^3(T) = \gamma_{v_2}^3(T) \neq 0$ , which implies  $\|\gamma_{v_1}^3(T)\| = \|\gamma_{v_2}^3(T)\|$ . Thus

$$\frac{\|B\|}{\sqrt{x_1^2 + 4B\bar{B}^T}} \|2 \sin\left(\frac{T}{2} \sqrt{x_1^2 + 4B\bar{B}^T}\right)\| = \frac{\|E\|}{\sqrt{x_2^2 + 4E\bar{E}^T}} \|2 \sin\left(\frac{T}{2} \sqrt{x_2^2 + 4E\bar{E}^T}\right)\|,$$

and we get a contradiction as was shown in Case 3 of the proof of Lemma 1.  $\square$

**Remark 2.** We mentioned that the distribution  $\mathcal{H}$  is strongly bracket generating. It may be worth mentioning that  $V_{n,1}$  is also a contact manifold, which is stated in [16] and also in [24]. To show that statement we note that the submersion  $U(1) \rightarrow V_{n,1} \rightarrow Gr_{n,1}$  can be written as  $S^1 \rightarrow S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$ . In [16] it is shown that for submersion  $S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$  the vertical vector space is spanned by

$$V(q) = -y_0 \partial_{x_0} + x_0 \partial_{y_0} - \dots - y_{n-1} \partial_{x_{n-1}} + x_{n-1} \partial_{y_{n-1}}, \quad q \in S^{2n-1}.$$

The horizontal distribution  $D$  is defined as the orthogonal complement to  $\text{span}\{V\}$  in  $TS^{2n-1}$  with respect to the Euclidean metric in  $\mathbb{R}^{2n} \cong \mathbb{C}^n$ . At the point  $(1, 0, \dots, 0) \in S^{2n-1}$  the vertical vector  $V = (i, 0, \dots, 0)$  coincides with the generator  $\xi = \begin{pmatrix} i & 0 \\ 0 & 0 \end{pmatrix}$  of the Lie algebra  $\mathfrak{u}_n(1)$  and the horizontal distribution  $D = V^\perp$  coincides with the horizontal distribution  $\mathcal{H} = \left\{ \begin{pmatrix} 0 & B \\ -\bar{B}^T & 0 \end{pmatrix} \mid B \in \mathbb{C}^{1 \times (n-1)} \right\}$  that is orthogonal to  $\xi$  with respect to the trace metric. Since metrics, vertical and horizontal distributions are invariant

under the action of  $U(n)$  we conclude that sub-Riemannian geometries are essentially the same. It is shown in [16] that the distribution  $D$  coincides with the holomorphic tangent space  $HS^{2n-1}$  of  $S^{2n-1}$  thought of as an embedded CR-manifold and that it also coincides with the contact distribution given by  $\ker(\omega)$  with respect to the contact form

$$\omega = -y_0 dx_0 + x_0 dy_0 - \dots - y_{n-1} dx_{n-1} + x_{n-1} dy_{n-1}.$$

Thus this information can be transferred to the Stiefel manifold.

## 5. THE CUT LOCI OF $V_{n,k}$

In the present section we show that some of the properties of the cut locus of  $V_{n,1}$  is preserved in the case  $V_{n,k}$ . In general we were not able to describe the total cut locus, since the distribution is not always strongly bracket generating, that leads to the existence of abnormal minimizers. Abnormal minimizers are also have to be taken into account since they can be minimizers due to [23]. The interested reader can find a further information about abnormal minimizers in [2, 7, 8, 9, 13, 14, 18, 20].

The fact that the distribution is in general not strongly bracket generating follows from the following proposition in [24].

**Proposition 1.** *Let  $Q$  be an  $m$ -dimensional manifold and  $\mathcal{H}$  an  $l$ -dimensional strongly bracket generating distribution of codimension 2 or greater. Then at least one of the following conditions*

- (1)  $l$  is a multiple of 4
- (2)  $l \geq (m - l) + 1$ .

have to be fulfilled.

It is obvious that it is not always the case for  $V_{n,k}$ , where  $m = 2nk - k^2$  and  $l = 2nk - 2k^2$ . Moreover, it is technically hard to write the exact form of normal sub-Riemannian geodesics for an arbitrary  $V_{n,k}$ .

**Proposition 2.** *The distribution  $\mathcal{H}$  on  $V_{n,k}$  is bracket generating.*

*Proof.* First we note that the commutator  $[\mathcal{H}, \mathcal{H}]$  is given by

$$\left[ \begin{pmatrix} 0 & B \\ -\bar{B}^T & 0 \end{pmatrix}, \begin{pmatrix} 0 & C \\ -\bar{C}^T & 0 \end{pmatrix} \right] = \begin{pmatrix} -B\bar{C}^T + C\bar{B}^T & 0 \\ 0 & -\bar{C}^T B + \bar{B}^T C \end{pmatrix}.$$

It is sufficient if we show that for every upper triangular  $(k \times k)$ -matrix  $D_{lm}$ ,  $m > l$  with an entry  $d_{lm} \neq 0$  on the intersection of  $l$ -row and  $m$ -column and all other entries vanish we can find  $B, C \in \mathbb{C}^{k \times (n-k)}$  such that  $D_{lm} = -B\bar{C}^T$ . For instance, if we choose

$$B = (b_{\alpha\beta}) = \quad \text{by} \quad b_{\alpha\beta} = \begin{cases} d_{lm} & \text{for } \alpha = l, \beta = \min\{m, n - k\} \\ 0 & \text{otherwise,} \end{cases}$$

and

$$-C^T = (c_{\alpha\beta}) \quad \text{by} \quad c_{\alpha\beta} = \begin{cases} 1 & \text{for } \alpha = \min\{m, n - k\}, \beta = m \\ 0 & \text{otherwise,} \end{cases}$$

then we deduce that  $D_{lm} = -B\bar{C}^T$ .

We also need to construct diagonal-form  $(k \times k)$ -matrices  $D_j$  with  $i \in \mathbb{C}$  on the intersection of  $j$ -row and  $j$ -column and all other entries vanish and show that there are  $B, C \in \mathbb{C}^{k \times (n-k)}$  such that  $D_j = -B\bar{C}^T$ . In this case we choose

$$B = (b_{\alpha\beta}) \quad \text{by} \quad b_{\alpha\beta} = \begin{cases} i & \text{for } \alpha = j, \beta = \min\{j, n-k\} \\ 0 & \text{otherwise,} \end{cases}$$

and

$$-\bar{C}^T = (c_{\alpha\beta}) \quad \text{by} \quad c_{\alpha\beta} = \begin{cases} 1 & \text{for } \alpha = \min\{j, n-k\}, \beta = j \\ 0 & \text{otherwise.} \end{cases}$$

Then we obtain that  $D_j = -B\bar{C}^T$ . It implies that  $\mathcal{H}$  is bracket generating.  $\square$

**Proposition 3.** *Suppose  $\gamma_v(t)$  is a normal sub-Riemannian geodesic, which connects the identity  $\text{Id}$  with the point  $q \in V_{n,k}$ ,  $q \neq \text{Id}$ , at the time  $T > 0$ , where  $v = \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \in \mathfrak{u}(n)$ . The length of  $\gamma_v$  is given by  $l(\gamma_v, T) = 2T\sqrt{n \operatorname{tr}(B\bar{B}^T)}$ .*

*Proof.* First of all we calculate the velocity vector of  $\gamma_v(t)$  at the time  $t$ , which is  $\dot{\gamma}_v(t) = \gamma_v(t)v_D$  for  $v_D \in \mathfrak{u}(n)$ . We omit the subscript  $U(n)$  or  $U(k)$  from  $\exp(\cdot)$ , since it is clear which one we use from the context. By the chain rule we get that

$$\begin{aligned} \dot{\gamma}_v(t) &= d_{f(t)}\pi_1 \left[ \left( \exp \left\{ t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right\} \right) \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \left( \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \right) \right. \\ &\quad \left. + \left( \exp \left\{ t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right\} \right) \left( \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \right) \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right], \end{aligned}$$

where  $f(t) := \exp \left( t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right) \exp \left( t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right)$ . We note that

$$\begin{aligned} &\begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} = \begin{pmatrix} A \exp(-tA) & B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix} \\ &= \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \begin{pmatrix} \exp(tA)A \exp(-tA) & \exp(tA)B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix} \\ &= \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \begin{pmatrix} A & \exp(tA)B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix}. \end{aligned}$$

Thus

$$\begin{aligned} \dot{\gamma}_v(t) &= d_{f(t)}\pi_1 \left[ \exp \left\{ t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right\} \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \right. \\ &\quad \left. \times \left( \begin{pmatrix} A & \exp(tA)B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix} + \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right) \right] \\ &= \gamma_v(t) \begin{pmatrix} 0 & \exp(tA)B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix} \end{aligned}$$

and

$$v_D = \begin{pmatrix} 0 & \exp(tA)B \\ -\bar{B}^T \exp(-tA) & 0 \end{pmatrix}.$$

It follows that

$$\begin{aligned}
g(\dot{\gamma}_v(t), \dot{\gamma}_v(t)) &= -2n \operatorname{tr}(v_D^2) \\
&= -2n \operatorname{tr} \left( \begin{pmatrix} -\exp(tA)B\bar{B}^T \exp(-tA) & 0 \\ 0 & -\bar{B}^T B \end{pmatrix} \right) \\
&= -2n \left( -\operatorname{tr}(\exp(tA)B\bar{B}^T \exp(-tA)) - \operatorname{tr}(\bar{B}^T B) \right) \\
&= 4n \operatorname{tr}(B\bar{B}^T).
\end{aligned}$$

In the last equation we used  $\operatorname{tr}(XY) = \operatorname{tr}(YX)$  and  $\operatorname{tr}(-X) = -\operatorname{tr}(X)$ .

It follows that the length of  $\gamma_v$  does not depend on  $A$ , but depend on  $T$  and the trace of the matrix  $B\bar{B}^T$ .  $\square$

**Theorem 4.** *The set*

$$L = \left\{ \left[ \begin{pmatrix} C & 0 \\ 0 & D \end{pmatrix} \right]_{V_{n,k}} \mid C \in U(k), D \in U(n-k) \right\} \setminus [\operatorname{Id}]_{V_{n,k}}$$

is a subset of the cut locus  $K_{\operatorname{Id}}$  on  $V_{n,k}$ .

*Proof.* Suppose the point  $[g]_{V_{n,k}} = \left[ \begin{pmatrix} C & 0 \\ 0 & D \end{pmatrix} \right]_{V_{n,k}} \in L$ . This point is reached optimally

by the geodesic  $\gamma_v = \left[ \begin{pmatrix} \gamma_v^1 & \gamma_v^2 \\ \gamma_v^3 & \gamma_v^4 \end{pmatrix} \right]_{V_{n,k}}$  at some  $T$  from the initial point  $\operatorname{Id} \in V_{n,k}$  and the

initial velocity vector  $v = \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \in \mathfrak{u}(n)$ . Let us see how  $\gamma_v^j$ ,  $j = 1, 2, 3, 4$ , depend on  $A$  and  $B$ . We recall that

$$\begin{aligned}
\gamma_v(t) &= \pi_1 \left( \exp \left\{ t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right\} \exp \left\{ t \begin{pmatrix} -A & 0 \\ 0 & 0 \end{pmatrix} \right\} \right) \\
&= \left[ \begin{pmatrix} \gamma_v^1(t) & \gamma_v^2(t) \\ \gamma_v^3(t) & \gamma_v^4(t) \end{pmatrix} \right]_{V_{n,k}}.
\end{aligned}$$

We start from calculating  $\exp \left( t \begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix} \right) = \begin{pmatrix} v_1(t) & v_2(t) \\ v_3(t) & v_4(t) \end{pmatrix}$ . Using the notation

$\begin{pmatrix} A & B \\ -\bar{B}^T & 0 \end{pmatrix}^n := \begin{pmatrix} v_1^n & v_2^n \\ v_3^n & v_4^n \end{pmatrix}$ , we find that  $v_1^n = v_1^{n-1}A - v_1^{n-2}B\bar{B}^T$ ,  $n \geq 2$ , for initial values  $v_1^0 = \operatorname{Id}$  and  $v_1^1 = A$ . This implies that  $v_1(t)$  depends on  $A$  and  $B\bar{B}^T$ . Furthermore, we get the formulas  $v_2^n = v_1^{n-1}B$ ,  $v_3^n = -\bar{B}^T v_1^{n-1}$  and  $v_4^n = -\bar{B}^T v_1^{n-2}B$ .

Now we claim that the geodesic  $\gamma_{v^*}$  with  $v^* := \begin{pmatrix} A & -B \\ \bar{B}^T & 0 \end{pmatrix}$  is also minimizing from  $\operatorname{Id}$  to  $[g]_{V_{n,k}}$  such that with  $\gamma_{v^*}(T) = [g]_{V_{n,k}}$ . Indeed, since  $(-B)(-\bar{B}^T) = B\bar{B}^T$  and  $(-\bar{B}^T)(-B) = \bar{B}^T B$  the length of both geodesics coincides. It remains to show that  $\gamma_{v^*}(T) = [g]_{V_{n,k}}$ . Observe, that the value  $v_1(t)$  depends on  $A$ ,  $B\bar{B}^T$  and  $t$ , and therefore  $\gamma_{v^*}^1(T) = \gamma_v^1(T)$ . Finally  $\gamma_{v^*}^2(T) = \gamma_v^2(T) = 0$  implies  $\gamma_{v^*}^2(T) = -\gamma_v^2(T) = 0 = \gamma_v^2(T)$  and  $\gamma_{v^*}^3(T) = -\gamma_v^3(T) = 0 = \gamma_v^3(T)$ . We conclude that  $\gamma_{v^*}(T) = \gamma_v(T)$  and that  $L \subset K_{\operatorname{Id}}$ .

The geodesic  $\gamma_{v^*}$  can be replaced by  $\gamma_{\hat{v}}$  with  $\hat{v} = \begin{pmatrix} A & -BU \\ (\bar{B}U)^T & 0 \end{pmatrix}$  for all  $U \in U(n-k)$ . It is also a minimizing geodesic from  $\operatorname{Id}$  to  $[g]_{V_{n,k}}$ , with  $\gamma_{\hat{v}}(T) = [g]_{V_{n,k}}$ .  $\square$

## REFERENCES

- [1] AGRACHEV, A. A.; SACHKOV, Y. L., *Control theory from the geometric viewpoint*. Encyclopaedia of Mathematical Sciences, 87. Control Theory and Optimization, II. Springer-Verlag, Berlin, 2004. pp. 412.
- [2] AGRACHEV, A. A.; SARYCHEV, A. V., *Abnormal sub-Riemannian geodesics: Morse index and rigidity*. Ann. Inst. H. Poincaré Anal. Non Linéaire. **13** (1996), pp. 635-690.
- [3] AGRACHEV, A. A.; SARYCHEV, A. V., *Sub-Riemannian metrics: minimality of abnormal geodesics versus subanalyticity*. ESAIM Control Optim. Calc. Var. **4** (1999), pp. 377-403
- [4] AGRACHEV, A.; BONNARD, B.; CHYBA, M.; KUPKA, I., *Sub-Riemannian sphere in Martinet flat case*. ESAIM Control, Optimisation and Calculus of Variations. **2** (1997), pp. 377-448.
- [5] AGRACHEV, A. A.; GAUTHIER, J. P., *Sub-Riemannian metrics and isoperimetric problems in the contact case*. In honor of L. Pontriaguine, 90th birthday commemoration, Contemporary Mathematics, Tome (in Russian). **64** (1999), pp. 5-48.
- [6] EDELMAN, A.; ARIAS, T. A.; SMITH, S. T., *The geometry of algorithms with orthogonality constraints*. SIAM Journal on Matrix Analysis and Applications. **20** (1999), pp. 303-353.
- [7] BONNARD, B.; CHYBA, M., *Méthodes géométriques et analytiques pour étudier l'application exponentielle, la sphère et le front d'onde en géométrie sous-riemannienne dans le cas Martinet*. ESAIM. Control, Optimisation and Calculus of Variations. **4** (1999), pp. 245-334.
- [8] BONNARD, B.; CHYBA, M.; KUPKA, I., *Nonintegrable geodesics in SR-Martinet geometry*. In "Differential geometry and control (Boulder, CO, 1997)", pp. 119-134. Amer. Math. Soc., Providence, RI.
- [9] BONNARD, B.; TRÉLAT, E., *On the role of abnormal minimizers in sub-Riemannian geometry*. Ann. Fac. Sci. Toulouse Math. (6) **10**(2001), pp. 405-491.
- [10] BOSCAIN, U.; ROSSI, F., *Invariant Carnot-Carathéodory metrics on  $S^3$ ,  $SO(3)$ ,  $SL(2)$ , and lens spaces*. SIAM J. Control Optim. **47** (2008), pp. 1851-1878.
- [11] CAPOGNA, L.; DANIELLI, D.; PAULS, S. D.; TYSON, J.T., *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*. Progress in Mathematics, 259. Birkhäuser Verlag, Basel, 2007. pp. 223.
- [12] CHOW, W.-L., *Über Systeme von linearen partiellen Differentialgleichungen erster Ordnung*. Math. Ann. **117** (1939), pp. 98-105.
- [13] CHYBA, M., *Le front d'onde en géométrie sous-riemannienne: le cas Martinet*. Séminaire de Théorie Spectrale et Géométrie, **16** (1997-1998), pp. 81-105. Univ. Grenoble I, Saint.
- [14] CHYBA, M., *La cas Martinet en Géométrie Sous-Riemannienne*. Ph.D. thesis, Univ. of Geneva.
- [15] GALLIER, J., *Notes on Differential Geometry and Lie Groups*. Unpublished manuscript. Retrieved from <http://www.cis.upenn.edu/~cis610/diffgeom-n.pdf>
- [16] GODOY MOLINA, M.; MARKINA, I., *Sub-Riemannian geodesics and heat operator on odd dimensional spheres*. Analysis and Mathematical Physics **2** (2012), pp. 123-147.
- [17] GROCHOWSKI, M., *Normal forms and reachable sets for analytic Martinet sub-Lorentzian structures of Hamiltonian type*. Journal of Dynamical and Control Systems **17** (2011), pp. 49-75.
- [18] HUANG, T.; YANG, X., *Extremals in some classes of Carnot groups*. Sci. China Math. **55** (2012), pp. 633-646.
- [19] KNAPP, A. W., *Lie groups beyond an introduction*. Second edition. Progress in Mathematics, 140. Birkhäuser Boston, Inc., Boston, MA, 2002. 812 pp.
- [20] LIU, W.; SUSSMANN, H. J., *Shortest paths for sub-Riemannian metrics on rank-two distributions*. Mem. Amer. Math. Soc. **118** (1995), pp. 104.
- [21] MANTON, J. H., *Optimization algorithms exploiting unitary constraints*. IEEE Transactions on Signal Processing. **50** (2002), pp. 635-650.
- [22] MILNOR, J., *Curvatures of left invariant metrics on Lie groups*, Advances in Mathematics **21** (1976), pp. 293-329.
- [23] MONTGOMERY, R., *Abnormal Minimizers*. SIAM Journal on Control and Optimization **32** (1994), pp. 1605-1620.
- [24] MONTGOMERY, R., *A tour of subriemannian geometries, their geodesics and applications*. Mathematical Surveys and Monographs, 91. American Mathematical Society, Providence, RI, 2002.

- [25] NAITOH, H.; SAKANE, Y., *On conjugate points of a nilpotent Lie group*. Tsukuba J. Math. **5** (1981), no. 1, 143-152.
- [26] RASHEVSKIĬ, P. K., *About connecting two points of complete nonholonomic space by admissible curve*, Uch. Zapiski Ped. Inst. K. Liebknecht **2** (1938), pp. 83-94.
- [27] SACHKOV, Y., *Cut locus and optimal synthesis in the sub-Riemannian problem on the group of motions of a plane*. ESAIM. Control, Optimisation and Calculus of Variations **17** (2011), pp. 293-321.
- [28] STRICHARTZ, R. S., *Sub-Riemannian geometry*. J. Differential Geom. **24** (1986), no. 2, pp. 221-263.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, NORWAY.

*E-mail address:* christian.autenried@math.uib.no

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, NORWAY.

*E-mail address:* irina.markina@math.uib.no