

THE ROYAL  
SWEDISH  
ACADEMY OF  
SCIENCES



**INSTITUT  
MITTAG-LEFFLER**

Auravägen 17, SE-182 60 Djursholm, Sweden  
Tel. +46 8 622 05 60 Fax. +46 8 622 05 89  
info@mittag-leffler.se www.mittag-leffler.se

**The choice number or complete multipartite  
graphs with part size four**

H. A. Kierstead, A. Salmon and R. Wang

REPORT No. 47, 2013/2014, spring

ISSN 1103-467X

ISRN IML-R- -47-13/14- -SE+spring

# ON THE CHOICE NUMBER OF COMPLETE MULTIPARTITE GRAPHS WITH PART SIZE FOUR

H. A. KIERSTEAD, ANDREW SALMON, AND RAN WANG

ABSTRACT. Let  $\text{ch}(G)$  denote the choice number of a graph  $G$ , and let  $K_{s*k}$  be the complete  $k$ -partite graph with  $s$  vertices in each part. Erdős, Rubin, and Taylor showed that  $\text{ch}(K_{2*k}) = k$ , and suggested the problem of determining the choice number of  $K_{s*k}$ . The first author established  $\text{ch}(K_{3*k}) = \lceil \frac{4k-1}{3} \rceil$ . Here we prove  $\text{ch}(K_{4*k}) = \lceil \frac{3k-1}{2} \rceil$ .

## 1. INTRODUCTION

Let  $G = (V, E)$  be a graph. A *list assignment*  $L$  for  $G$  is a function  $L : V \rightarrow 2^{\mathbb{N}}$ , where  $\mathbb{N}$  is the set of natural numbers and  $2^{\mathbb{N}}$  is the power set of  $\mathbb{N}$ . If  $|L(v)| = k$  for all vertices  $v \in V$ , then  $L$  is a  *$k$ -list assignment* for  $G$ . An  *$L$ -coloring*  $f$  from a list assignment  $L$  is a function  $f : V \rightarrow \mathbb{N}$  such that  $f(v) \in L(v)$  for all vertices  $v \in V$  and  $f(x) \neq f(y)$  whenever  $xy \in E$ .  $G$  is  *$L$ -colorable* if there exists an  $L$ -coloring of  $G$ ; it is  *$k$ -choosable* if it is  $L$ -colorable for all  $k$ -list assignments  $L$ . The *list chromatic number* or *choice number* of  $G$ , denoted  $\text{ch}(G)$ , is the smallest integer  $k$  such that  $G$  is  $k$ -choosable. The general list coloring problem may consider list assignments with uneven list sizes.

The study of list coloring was initiated by Vizing [16] and by Erdős, Rubin and Taylor [4]. It is a generalization of two well studied areas of combinatorics—graph coloring and transversal theory. Restricting the list assignment to a constant function, yields ordinary graph coloring; restricting the graph to a clique yields the problem of finding a system of distinct representatives (SDR) for the family of lists. Both restrictions play a role in this paper. Given the general nature of this parameter, it is hardly surprising that there are not many graphs whose exact choice number is known. However, there are some amazingly elegant results that add to the subject's charm. For example, Thomassen [15] proved that planar graphs have choice number at most 5, Voigt [18] proved that this is tight, and Galvin [5] proved that line graphs of bipartite graphs have choice number equal to their clique number.

Erdős et al. [4] suggested determining the choice number of uniform complete multipartite graphs. More generally, let  $K_{1*k_1, 2*k_2, \dots}$  denote the complete multipartite graph with  $k_i$  parts of size  $i$ , where zero terms in the subscript are deleted. Since  $K_{1*k}$  is a clique and  $K_{s*1}$  is an independent set, these cases are trivial. Alon [1] proved the general bounds  $c_1 k \log s \leq \text{ch}(K_{s*k}) \leq c_2 k \log s$  for some constants  $c_1, c_2 > 0$ . This was tightened by Gazit and Krivelevich [6].

**Theorem 1** (Gazit and Krivelevich [6]).  $\text{ch}(K_{s*k}) = (1 + o(1)) \frac{\log s}{\log(1+1/k)}$ .

The next well-known example provides the best lower bounds for small values of  $s$ .

---

2000 *Mathematics Subject Classification.* 05C15.

*Key words and phrases.* List coloring, choice number, choosable, on-line choice number.

The first author thanks Institut Mittag-Leffler (Djursholm, Sweden) for its hospitality and creative environment.

Research of the first author was supported in part by NSA grant H98230-12-1-0212.

**Example 2.**  $\text{ch}(K_{s**k}) \geq l = \lceil \frac{2(s-1)k-s+2}{s} \rceil$ : Let  $G = K_{s**k}$  have parts  $\{X_1, \dots, X_k\}$  with  $X_i = \{v_{i,1}, \dots, v_{i,s}\}$ . We will construct an  $(l-1)$ -list assignment  $L$  from which  $G$  cannot be colored. Equitably partition  $C := [2k-1]$  into  $s$  parts  $C_1, \dots, C_s$ . Define a list assignment  $L$  for  $G$  by  $L(v_{i,j}) = C \setminus C_j$ . Then each list has size at least

$$2k-1 - \left\lfloor \frac{2k-1}{s} \right\rfloor = \left\lfloor \frac{2ks-s-2k+1}{s} \right\rfloor = \left\lfloor \frac{2(s-1)k-2s+2}{s} \right\rfloor = l-1.$$

Consider any color  $\alpha \in C$ . Then  $\alpha \in C_i$  for some  $i \in [s]$ . So  $\alpha \notin L(x_{i,j})$  for every  $j \in [k]$ . Thus any  $L$ -coloring of  $G$  uses at least two colors for every part  $X_j$ . Since vertices in distinct parts are adjacent, they require distinct colors. As there are  $k$  parts this would require  $2k > |C|$  colors, which is impossible.

Restricting the question of Erdős et al., we ask for those integers  $s$  such that:

$$(1.1) \quad (\forall k \in \mathbb{Z}^+) \left[ \text{ch}(K_{s**k}) = l(s, k) := \left\lceil \frac{2(s-1)k-s+2}{s} \right\rceil \right].$$

The first two cases  $s=2$  and  $s=3$  have been solved:

**Theorem 3** (Erdős, Rubin and Taylor [4]). *All positive integers  $k$  satisfy  $\text{ch}(K_{2**k}) = k$ .*

**Theorem 4** (Kierstead [7]). *All positive integers  $k$  satisfy  $\text{ch}(K_{3**k}) = \lceil \frac{4k-1}{3} \rceil$ .*

Recently, Kozik, Micek, and Zhu [8] gave a very different proof of Theorem 4. The following more general result appears in [11].

**Theorem 5** (Ohba [11]).  $\text{ch}(K_{1**k_1, 3**k_3}) = \max\{k, \lceil \frac{n+k-1}{3} \rceil\}$ , where  $k = k_1 + k_3$  and  $n = k_1 + 3k_3$ .

The next example shows that the largest  $s$  satisfying (1.1) is at most 14.

**Example 6.** If  $k$  is even then  $\text{ch}(K_{15**k}) \geq l := 2k$ : Let  $G = K_{15**k}$  have parts  $\{X_1, \dots, X_k\}$  with  $X_i = \{v_{i,1}, \dots, v_{i,15}\}$ . We will construct an  $(l-1)$ -list assignment  $L$  from which  $G$  cannot be colored. Equitably partition  $C := [3k-1]$  into 6 parts  $C_1, \dots, C_6$ , and fix a bijection  $f : [15] \rightarrow \binom{[6]}{2}$ . Define a list assignment  $L$  for  $G$  by

$$L(v_{i,j}) = C \setminus \bigcup \{C_h : h \in f(i)\}.$$

Then each list has size at least

$$3k-1 - 2 \left\lfloor \frac{3k-1}{6} \right\rfloor = 2k-1 = l-1.$$

Consider any two colors  $\alpha, \beta \in C$ . Then  $\alpha, \beta \in \bigcup \{C_h : h \in f(i)\}$  for some  $i \in [15]$ . So  $\alpha, \beta \notin L(x_{i,j})$  for every  $j \in [k]$ . Thus any  $L$ -coloring of  $G$  uses at least three colors for every part  $X_j$ . Since  $3k > |C|$ , this is impossible.

Yang [19] proved  $\lceil \frac{3k}{2} \rceil \leq \text{ch}(K_{4**k}) \leq \lceil \frac{7k}{4} \rceil$ , and Noel, West, Wu and Zhu [10] improved the upper bound to  $\lceil \frac{5k-1}{3} \rceil$ . The main result of this paper is that (1.1) holds for  $s=4$ . To prove this theorem we first extract a simple proof of Theorem 4 from [10], and then elaborate on it.

**Theorem 7.**  $\text{ch}(K_{4**k}) = l(4, k) := \lceil \frac{3k-1}{2} \rceil$ .

Some of the recent development of list coloring of complete multipartite graphs has been motivated by paintability, or on-line choosability. Introduced independently by Schauz [14] and Zhu [20] (as on-line list colouring), *paintability* is a coloring game played between two players Alice and Bob on a graph  $G = (V, E)$  and a function  $f : V \rightarrow \mathbb{N}$ .

Let  $V_i$  denote the vertex set at the start of round  $i$ ; so  $V_1 = V$ . At round  $i$ , Alice selects a nonempty set of vertices  $A_i \subseteq V_i$ , and Bob selects an independent set  $B_i \subseteq A_i$ . Then  $B_i$  is deleted from the graph so that  $V_{i+1} = V_i \setminus B_i$ , and the rounds are continued until  $V_n = \emptyset$ . Alice's goal is to present some vertex  $v$  more than  $f(v)$  times, while Bob's goal is to choose every vertex before it has been presented  $f(v) + 1$  times. We say that  $G$  is *on-line  $f$ -choosable* if Bob has a strategy such that any vertex  $v \in V$  is in at most  $f(v)$  sets  $A_i$ , and *on-line  $k$ -choosable* if  $G$  is on-line  $f$ -choosable when  $f(v) = k$  for all  $v \in V$ . The *on-line choice number*, denoted  $\text{ch}^{\text{ol}}(G)$ , is the least  $k$  such that  $G$  is on-line  $k$ -choosable.

This game formulation hides the on-line nature of the problem. Another way of thinking about it is that Alice has secretly assigned lists of colors to all the vertices. At round  $i$  she reveals all vertices whose list contains color  $i$ , and Bob colors an independent set of them with color  $i$ . In this formulation it is clear that  $\text{ch}(G) \leq \text{ch}^{\text{ol}}(G)$ .

Schauer [14] proved that surprisingly many results on choice number, including Brooks' theorem [3], Thomassen's theorem [15], and the Bondy–Boppana–Siegel kernel lemma carry over to on-line choice number. It is unknown whether  $\text{ch}^{\text{ol}}(G) - \text{ch}(G)$  is bounded by a constant. Indeed, no graphs are known for which  $\text{ch}^{\text{ol}}(G) - \text{ch}(G) \geq 2$ . It is known that

$$\text{ch}(K_{2,2,3}) = 3 < 4 = \text{ch}^{\text{ol}}(K_{2,2,3}).$$

The explicit value of  $\text{ch}(K_{4*k})$  provided by Theorem 7 may be useful for establishing larger gaps. In Section 4 we show that  $\text{ch}(K_{4*3}) < \text{ch}^{\text{ol}}(K_{4*3})$ .

## 2. SET-UP

Fix  $s, k \in \mathbb{Z}^+$ . Let  $G = (V, E) = K_{s*k}$ , and let  $\mathcal{P}$  be the partition of  $V$  into  $k$  independent  $s$ -sets. Let  $l = l(k, s) = \lceil \frac{(s-1)2k-s+2}{s} \rceil$ , and consider any  $l$ -list assignment  $L$  for  $G$ . Put  $C^* = \bigcup_{x \in V} L(x)$ . Let  $L-\alpha$  be the result of deleting  $\alpha$  from every list of  $L$ .

We may write  $x_1 \dots x_t$  for the subpart  $S = \{x_1, \dots, x_t\} \subseteq X \in \mathcal{P}$ ; when we use this notation we implicitly assume the  $x_i$  are distinct. Also set  $\bar{S} = X \setminus S$ . For a set of vertices  $S \subseteq V$  let  $\mathcal{L}(S) = \{L(x) : x \in S\}$ ,  $L(S) = \bigcap \mathcal{L}(S)$ ,  $W(S) = \bigcup \mathcal{L}(S)$ , and  $l(S) = |L(S)|$ . The operation of replacing the vertices in  $S$  by a new vertex  $v_S$  with the same neighborhood as  $S$  is called *merging*. The new vertex  $v_S$  is said to be *merged*; vertices that are not merged are called *original*. When merging a set  $S$  we also create a list  $L(v_S) = L(S)$ .

For a color  $\alpha \in C^*$ , let  $|X, \alpha| = |\{x \in X : \alpha \in L(x)\}|$  be the number of times  $\alpha$  appears in the lists of vertices of  $X$ ,  $N_i(X) = \{\alpha \in C^* : |X, \alpha| = i\}$  be the set of colors that appear exactly  $i$  times in the lists of vertices in  $X$ ,  $n_i(X) = |N_i(X)|$ ,  $N(X) = N_2(X) \cup N_3(X)$ , and  $n(X) = |N(X)|$ . Let  $\sigma_i(X) = \sum \{l(I) : I \subseteq X \wedge |I| = i\}$  and  $\mu_i(X) = \max \{l(I) : I \subseteq X \wedge |I| = i\}$ .

For a set  $S$  and element  $x$  we use the notation  $S + x = S \cup \{x\}$  and  $S - x = S \setminus \{x\}$ . The range of a function  $f$  is denoted by  $\text{ran}(f)$ .

The following lemma was proved independently by Kierstead [7], and by Reed and Sudakov [12], [13], and named by Rabern.

**Lemma 8** (Small Pot Lemma). *If  $\text{ch}(G) > r$  then there exists a  $r$ -list assignment  $L$  using fewer than  $|V(G)|$  colors such that  $G$  is not  $L$ -colorable.*

If  $s$  does not satisfy (1.1) then there is a minimum counterexample  $k$  with  $\text{ch}(K_{s*k}) > l(s, k)$ . By the Small Pot Lemma, this is witnessed by a list assignment  $L$  with  $|\bigcup \{L(x) : x \in V(G)\}| < |V|$ . We always assume  $L$  has this property.

**Lemma 9.** *If  $G$  is a minimum counterexample then every part  $X$  of  $G$  satisfies  $L(X) = \emptyset$ .*

*Proof.* Otherwise there exists a list assignment  $L$ , a color  $\alpha$ , and a part  $X$  such that  $\alpha \in L(X)$ . Color each vertex in  $X$  with  $\alpha$ , set  $G' = G - X$ , and put  $L' = L - \alpha$ . Then  $L'$  witnesses that  $k - 1$  is a smaller counterexample, a contradiction.  $\square$

By Lemma 9,  $n_s(X) = 0$  for each part  $X \in \mathcal{P}$ . So by the Small Pot Lemma,  $|W(X)| = \sum_{i=1}^{s-1} n_i(X) < sk$ . Also  $\sum_{i=1}^{s-1} i n_i(X) = sl$  is the number of occurrences of colors in the lists of vertices of  $X$ . Thus

$$(2.1) \quad \sum_{i=1}^{s-1} (i-1)n_i(X) \geq sl - |W(X)| \geq s(l-k) + 1.$$

Now we warm-up by giving a short proof extracted from [10] of Theorem 4.

*Proof of Theorem 4.* Let  $s = 3$ ,  $l = l(3, k) = \text{ch}(K_{3*k}) = \lceil \frac{4k-1}{3} \rceil$ , and assume  $G$  is a counterexample with  $k$  minimal. Then  $k > 1$ . By Lemma 9,  $n_3(X) = 0$  for all  $X \in \mathcal{P}$ . We obtain a contradiction by  $L$ -coloring  $G$ . First we use the following steps to partition  $V$  into sets of vertices that will receive the same color. Then we *merge* each set  $I$  into a single vertex  $v_I$ , and assign  $v_I$  the set of colors in  $L(I)$ . Finally we apply Hall's Theorem to choose a system of distinct representatives (SDR) for these new lists; this induces an  $L$ -coloring of  $G$ .

**Step 1.** Partition  $\mathcal{P}$  into a set  $\mathcal{R}$  of  $l - k$  reserved parts and a set  $\mathcal{U} = \mathcal{P} \setminus \mathcal{R}$  of  $2k - l$  unreserved parts.

**Step 2.** Choose  $\mathcal{U}_1 \subseteq \mathcal{U}$  to maximize  $\nu = \sum_{X \in \mathcal{U}_1} \mu_2(X)$  subject to the constraint  $|\mathcal{U}_1| \leq \mu_2(X)$  for all  $X \in \mathcal{U}_1$ . Set  $u_1 = |\mathcal{U}_1|$ . For each  $X \in \mathcal{U}_1$  choose a pair  $I_X \subseteq X$  with  $l(I_X) \geq u_1$  maximum. Put  $\mathcal{U}_2 = \mathcal{U} \setminus \mathcal{U}_1$  and  $u_2 = |\mathcal{U}_2|$ .

The maximality of  $\nu$  implies

$$(2.2) \quad \mu_2(X) \leq u_1 \text{ for all } X \in \mathcal{U}_2,$$

as otherwise adding  $X$  to  $\mathcal{U}_1$ , and deleting one part  $Y \in \mathcal{U}_1$  with  $\mu_2(Y) = u_1$ , if such a part  $Y$  exists, would increase  $\nu$ .

**Step 3.** Using (2.1), each part  $X \in \mathcal{P}$  satisfies

$$n_2(X) \geq 3(l-k) + 1 = 3 \left\lceil \frac{k-1}{3} \right\rceil + 1 \geq k - 1 + 1 = k.$$

Form an SDR  $f$  for  $\{L(v_{I_X}) : X \in \mathcal{U}_1\} \cup \{N(X) : X \in \mathcal{R}\}$  by greedily choosing representatives for the first family and then for the second family. For each  $X \in \mathcal{R}$  choose a pair  $I_X \subseteq X$  so that  $f(x) \in L(I_X)$ .

**Step 4.** For each  $X \in \mathcal{U}_1 \cup \mathcal{R}$ , merge  $I_X$  to a new vertex  $v_{I_X}$ , let  $z_X \in X \setminus I_X$ , and set  $X' = \{v_{I_X}, z_X\}$ . If  $X \in \mathcal{U}_2$ , set  $X' = X$ . This yields a graph  $G'$  with parts  $\mathcal{P}' = \{X' : X \in \mathcal{P}\}$ , and list assignment  $L$ .

Next we use Hall's Theorem to prove that  $\{L(x) : x \in V(G')\}$  has an SDR. For this it suffices to prove:

$$(2.3) \quad |S| \leq \left| \bigcup \{L(x) : x \in S\} \right| \text{ for every } S \subseteq V(G').$$

To prove (2.3), let  $S \subseteq V(G')$  be arbitrary, and set  $W = W(S) := \bigcup \{L(x) : x \in S\}$ . We consider several cases in order, always assuming all previous cases fail.

**Case 1:** There exists  $X \in \mathcal{P}$  with  $|S \cap X'| = 3$ . Then  $X' = X \in \mathcal{U}_2$ ,  $u_2 \geq 1$ , and  $|S| \leq |G'| \leq 2k + u_2$ . By (2.2),  $u_1 \geq \mu_2(X) \geq \sigma_2(X)/3$ . Using inclusion-exclusion, and Lemma 9,

$$\begin{aligned} |W| &\geq |W(X)| = \sigma_1(X) - \sigma_2(X) + \sigma_3(X) \geq 3l - 3u_1 = 3l - 3(2k - l - u_2) \\ &\geq 6(l - k) + 3u_2 \geq (2k - 2) + (2 + u_2) \geq 2k + u_2 \geq |S|. \end{aligned}$$

**Case 2:** There is  $X \in \mathcal{U}_2$  with  $|S \cap X'| = 2$ . Since  $u_1 = 2k - l - u_2 < 2k - l$ , By (2.2),  $|W| \geq |W(S \cap X)| \geq 2l - l(S \cap X) \geq 2l - u_1 \geq 2l - (2k - l - u_2) \geq 3l + 1 - 2k \geq 2k \geq |S|$ .

**Case 3:** There is  $X \in \mathcal{U}_1$  with  $|S \cap X'| = 2$ . As  $|S| \leq 2k - u_2 = l + u_1$  and  $L(v_{I_X} z_X) = L(X) = \emptyset$ ,

$$|W| \geq |W(S \cap X')| \geq l(v_{I_X}) + l(z_X) - l(v_{I_X} z_X) \geq u_1 + l \geq |S|.$$

**Case 4:**  $S$  has an original vertex. Then  $|S| \leq 2k - u_1 - u_2 = l \leq |W|$ .

**Case 5:** All vertices of  $S$  have been merged. Then  $|S| \leq |f(S)| \leq |W|$ .

□

### 3. THE MAIN THEOREM

In this section we prove our main result, Theorem 7. The case when  $k$  is odd is considerably more technical. Casual or first time readers may wish to avoid these additional details; the proof is organized so that this is possible. In particular, in the even case Step 7(b), Step 11, Lemma 10(b), and Lemma 14 are not involved. Furthermore, only the first conclusion of Lemma 13 that  $k$  is odd (in the bad case covered by its hypothesis) is used. Let  $b \in \{0, 1\}$  with  $b \equiv k \pmod{2}$  and  $l = l(4, k) = \lceil \frac{3k-1}{2} \rceil$ . We often use the partition  $k = (2k - l) + (l - k)$  of the integer  $k$ , and note that  $2k - l = l - k + b$ .

*Proof of Theorem 7.* Our set-up is the same as in the proof of Theorem 4. Let  $s = 4$ ,  $l = l(4, k)$ , and  $G = K_{4+k}$ . The theorem is trivial if  $k = 1$ . Let  $k > 1$  be a minimal counterexample, and let  $L$  be an  $l$ -list assignment for  $G$  with  $|W(V)| \leq 4k - 1$  and  $L(X) = \emptyset$  for all parts  $X \in \mathcal{P}$ . Again we partition  $V$  into sets of vertices that will receive the same color, and then find an SDR for the induced list assignment that in turn induces an  $L$ -coloring of  $G$ . See Figure 3.1.

**Step 1.** Reserve notation for a partition  $\mathcal{P} = \mathcal{R} \cup \mathcal{U}$  of  $V$  with  $|\mathcal{R}| = l - k$ ,  $|\mathcal{U}| = 2k - l$ ,  $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2 \cup \mathcal{R}_3$ , and  $\mathcal{U} = \mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3 \cup \mathcal{U}_4$ , where  $\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3, \mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3, \mathcal{U}_4$  are to be defined in the following steps.

**Step 2.** Choose  $\mathcal{U}_1 \subseteq \mathcal{P}$  so that  $|\mathcal{U}_1| \leq 2k - l$ , for every  $X \in \mathcal{U}_1$  there is a pair  $I_X \subseteq X$  with (\*)  $l(I_X), l(\bar{I}_X) \geq k$ , and subject to these constraints  $|\mathcal{U}_1|$  is maximum. For each  $X \in \mathcal{U}_1$  fix  $I_X$  witnessing (\*). Let  $u_1 := |\mathcal{U}_1|$ . Then:

$$(3.1) \quad \text{If } u_1 < 2k - l \text{ then } (\forall X \in \mathcal{P} \setminus \mathcal{U}_1)(\forall I \subseteq X)[|I| = 2 \rightarrow \min\{l(I), l(\bar{I})\} \leq k - 1].$$

**Step 3.** Choose  $\mathcal{U}_2 \subseteq \mathcal{P} \setminus \mathcal{U}_1$  so that  $|\mathcal{U}_2| \leq 2k - l - u_1$  and  $|\mathcal{U}_2| \leq \mu_3(X)$  for all  $X \in \mathcal{U}_2$ ; subject to this let  $\nu_3 = \sum_{X \in \mathcal{U}_2} \mu_3(X)$  be maximum. Let  $u_2 = |\mathcal{U}_2|$ . If  $\mathcal{U}_2 \neq \emptyset$  we select a part  $\dot{Z} \in \mathcal{U}_2$ ; else  $\dot{Z} = \emptyset$ . For each  $X \in \mathcal{U}_2$  choose a triple  $I_X \subseteq X$  with  $l(I_X) \geq u_2$  maximum. Since  $\nu_3$  cannot be increased:

$$(3.2) \quad \text{If } u_1 + u_2 < 2k - l \text{ then } (\forall X \in \mathcal{U}_3 \cup \mathcal{U}_4 \cup \mathcal{R})[\mu_3(X) \leq u_2].$$

**Step 4.** Choose  $\mathcal{R}_1 \subseteq \mathcal{P} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2)$  so that  $|\mathcal{R}_1| \leq l - k$  and there is a family of sets  $\{I_X : X \in \mathcal{R}_1\}$  such that (\*)  $I_X \subseteq X$ ,  $|I_X| = 3$ , and there is an SDR  $f_1$  of  $\mathcal{L}(M_1)$ , where

$M_1 := \{v_{I_X} : X \in \mathcal{U}_2 \cup \mathcal{R}_1\}$ ; subject to this constraint, choose  $\mathcal{R}_1$  with  $|\mathcal{R}_1|$  maximum. Fix  $\{I_X : X \in \mathcal{R}_1\}$ ,  $f_1$  and  $M_1$  satisfying (\*). Let  $C_1 = \text{ran}(f_1)$  and  $r_1 := |\mathcal{R}_1|$ . Then:

$$(3.3) \quad \text{If } r_1 < l - k \text{ then } (\forall X \in \mathcal{U}_3 \cup \mathcal{U}_4 \cup \mathcal{R}_2 \cup \mathcal{R}_3)[N_3(X) \subseteq C_1].$$

Moreover, by Lemma 9,  $L(T) \cap L(T') = \emptyset$  for any two triples  $T, T' \subseteq X$ , and so

$$(3.4) \quad \text{If } r_1 < l - k \text{ then } (\forall X \in \mathcal{U}_3 \cup \mathcal{U}_4 \cup \mathcal{R}_2 \cup \mathcal{R}_3)[\sigma_3(X) \leq u_2 + r_1].$$

**Step 5.** Choose  $\mathcal{U}_3 \subseteq \mathcal{P} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{R}_1)$  so that  $|\mathcal{U}_3| \leq 2k - l - u_1 - u_2$  and  $l - k + u_2 + |\mathcal{U}_3| \leq \mu_2(X)$  for all  $X \in \mathcal{U}_3$ ; subject to this constraint let  $\nu_5 = \sum_{X \in \mathcal{U}_3} \mu_2(X)$  be maximum. Let  $u_3 = |\mathcal{U}_3|$ . Since  $\nu_5$  cannot be increased:

$$(3.5) \quad \text{If } u_1 + u_2 + u_3 < 2k - l \text{ then } (\forall X \in \mathcal{U}_4 \cup \mathcal{R}_2 \cup \mathcal{R}_3)[\mu_2(X) \leq l - k + u_2 + u_3].$$

For all  $X \in \mathcal{U}_3$  choose a pair  $I_X = xy \subseteq X$  with  $l(I_X) \geq l - k + u_2 + u_3$  maximum; subject to this choose  $I_X$  so that  $\Delta_1(I_X) := l(I_X) - l(\bar{I}_X)$  is maximum. Set  $\Delta_2(I_X) := 2u_2 - l(xyz) - l(xyw)$ , where  $zw = \bar{I}_X$ .

**Step 6.** Choose  $\mathcal{R}_2 \subseteq \mathcal{P} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3 \cup \mathcal{R}_1)$  so that  $|\mathcal{R}_2| \leq l - k - r_1$  and  $\sigma_2(X) - \sigma_3(X) \geq 5(l - k) + 2u_1 + 2u_2 + u_3 + r_1 + |\mathcal{R}_2|$  for all  $X \in \mathcal{R}_2$ ; subject to this constraint let  $\nu_6 = \sum_{X \in \mathcal{R}_2} (\sigma_2(X) - \sigma_3(X))$  be maximum. Set  $r_2 = |\mathcal{R}_2|$ . Then:

$$(3.6) \quad \begin{aligned} &\text{If } r_1 + r_2 < l - k \text{ then } (\forall X \in \mathcal{U}_4 \cup \mathcal{R}_3) \\ &[\sigma_2(X) - \sigma_3(X) \leq 5(l - k) + 2u_1 + 2u_2 + u_3 + r_1 + r_2]. \end{aligned}$$

**Step 7.** Choose  $\mathcal{R}_3 \subseteq \mathcal{P} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3 \cup \mathcal{R}_1 \cup \mathcal{R}_2)$  with  $|\mathcal{R}_3| = l - k - r_1 - r_2$ . Let  $r_3 = |\mathcal{R}_3|$ . If  $r_3 = 0$  then go to Step 8. Otherwise, for  $I \subseteq X \in \mathcal{R}_3$ , put  $L^1(I) = L(I) \setminus C_1$ ,  $l^1(I) = |L^1(I)|$ , and if  $|I| = 2$ , let  $\Delta_1^1(I) = l^1(I) - l^1(\bar{I})$ . By Lemma 10, all 2-sets  $I \subset X$  satisfy

$$(3.7) \quad W(\bar{I} + v_I) \geq 3k - b + \Delta_1^1(I_X) - u_2 - r_1.$$

Set  $\dot{r} = 0$  and do one of (a) or (b) below.

(a) If  $b = 0$  or there are both  $X \in \mathcal{R}_3$  and a 2-set  $I \subset X$  with  $\Delta_1^1(I) \geq 1$  then choose  $\dot{Y}$  and a 2-set  $I_{\dot{Y}} \subset \dot{Y}$ , so that  $\Delta_1^1(I)$  is maximum. By Lemma 10(a) there is a family  $\mathcal{J} = \{I_X : X \in \mathcal{R}_3 - \dot{Y}\}$  such that  $I_X \subseteq X$ ,  $|I_X| = 2$ , and  $\Delta_1^1(I_X) \geq 0$  for all  $X \in \mathcal{R}_3$ ; and  $\mathcal{L}(M_2 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\})$  has an SDR  $f_2$  extending  $f_1$ . Fix such  $\mathcal{I} = \mathcal{J} + I_{\dot{Y}} = \{I_X : X \in \mathcal{R}_3\}$  and  $f_2$ , and set  $M_2 = M_1 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\}$  and  $C_2 = \text{ran}(f_2)$ .

(b) Otherwise  $b = 1$  and every  $X \in \mathcal{R}_3$  and every 2-set  $I \subset X$  satisfies  $\Delta_1^1(I) = 0$ . Pick a class  $\dot{Y} \in \mathcal{R}_3$ . By Lemma 10(b), there is a 2-set  $I_{\dot{Y}} \subset \dot{Y}$  such that  $\mathcal{I}$  can be chosen so that  $\mathcal{L}(M_2 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\} + v_{\bar{I}_{\dot{Y}}})$  has an SDR  $h$  extending  $f_1$ . Let  $M_2 = M_1 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\}$ ,  $f_2$  equal  $h$  restricted to  $M_2$ , and  $C_2 = \text{ran}(f_2)$ . (b\*) If  $u_1 = 0 = r_2$  then Step 9 is degenerate; set  $f_3 = h$ ,  $M_3 = M_2 + v_{\bar{I}_{\dot{Y}}}$ ,  $C_3 = \text{ran}(f_3)$ , and  $\dot{r} = 1$ .

**Step 8.** Put  $\mathcal{U}_4 := \mathcal{U} \setminus (\mathcal{R} \cup \mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3)$ , and  $u_4 := |\mathcal{U}_4|$ .

**Step 9.** Using Lemma 12, choose a family  $\mathcal{I} = \{I_X : X \in \mathcal{R}_2\}$  such that  $I_X \subseteq X$  and  $|I_X| = 2$  for all  $X \in \mathcal{R}_2$ , and  $\mathcal{L}(M_2 \cup \{v_{I_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\} \cup \{v_{\bar{I}_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\})$  has an SDR  $f_3$  that extends  $f_2$ . Set  $M_3 = M_2 \cup \{v_{I_X}, v_{\bar{I}_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\}$  and  $C_3 = \text{ran}(f_3)$ .

**Step 10.** Let  $G' := (V', E')$  be the graph obtained from  $G$  by merging each  $I_X$  with  $X \in \mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3 \cup \mathcal{R}_1 \cup \mathcal{R}_2 \cup \mathcal{R}_3$  and each  $\bar{I}_X$  with  $X \in \mathcal{U}_1 \cup \mathcal{R}_2$ . Note that this does not include  $\bar{I}_{\dot{Y}}$  even if Step 7(b\*) is executed. For a part  $X$ , let  $X'$  be the corresponding part in  $G'$ , and set  $\mathcal{P}' = \{X' : X \in \mathcal{P}\}$ .




FIGURE 3.1. The partition  $\mathcal{P}$  of  $K_{4+k}$ : The ovals indicate sets of vertices that are merged to form  $\mathcal{P}'$ .

**Step 11.** Set  $\dot{u} = \ddot{u} = 0$ . If  $k$  is odd ( $b = 1$ ) then we merge one more pair of vertices under the following special circumstances:

- (a) there exists  $X \in \mathcal{U}_4$  with  $|W(X)| < |G'|$ . Fix such an  $X = \dot{X}$ . By Lemma 13,  $u_1 = 0 = r_3$  and there is a pair  $I_{\dot{X}} \subseteq \dot{X}$  such that (i)  $f_3$  can be extended to an SDR  $f$  of  $\mathcal{L}(M)$ , where  $M := M_3 + v_{I_{\dot{X}}}$ ; (ii)  $|W(\{v_{I_{\dot{X}}}, v\})| \geq 2k - 1$ , and if equality holds then  $|W(\{v_{I_{\dot{X}}}, v\} \cup \dot{Z}') \cup C_4| \geq 2k$  for both  $v \in \bar{I}_{\dot{X}}$ ; and (iii)  $W(\bar{I}_{\dot{X}} + v_{I_{\dot{X}}}) \geq |G'| - 1$ . Merge  $I_{\dot{X}}$  and set  $\dot{u} = 1$ .
- (b) condition (a) fails and there exist  $X \in \mathcal{U}_4$  and  $xyz \subseteq X$  with

$$|W(xyz \cup \dot{Z}')| \leq 2k + u_4 - 1 < |W(X)|.$$

Fix such an  $X = xyzw = \ddot{X}$ . By Lemma 14 there is a pair  $I_{\ddot{X}} \subseteq xyz$  such that (i)  $f_3$  can be extended to an SDR  $f$  of  $\mathcal{L}(M)$ , where  $M := M_3 + v_{I_{\ddot{X}}}$ ; (ii)  $|W(\{v_{I_{\ddot{X}}}, v\})| \geq 2k$  for  $v \in xyz \setminus I_{\ddot{X}}$  and  $|W(\{v_{I_{\ddot{X}}}, w\})| \geq 2k - 1 + u_3$ ; and (iii)  $|W(\bar{I}_{\ddot{X}} + v_{I_{\ddot{X}}})| \geq 2k + u_4$ . Merge  $I_{\ddot{X}}$  and set  $\ddot{u} = 1$ .

**Step 12.** Recall that  $G'$  is the graph obtained after the first ten steps. Let  $H$  be the final graph obtained by this merging procedure, including  $\bar{I}_{\dot{Y}}$  if Step 7(b\*) is executed. (If  $b = 0$ , and possibly otherwise,  $H = G'$ ). Also let  $M$  be the final set of merged vertices,  $f$  be the final SDR of  $\mathcal{L}(V(H))$ , and  $C = \text{ran}(f)$ .

Recall that  $f_i$  is an SDR for  $\mathcal{L}(M_i)$  with  $\text{ran}(f_i) = C_i$ ,  $M$  is the final set of merged vertices,  $f$  is the final SDR of  $\mathcal{L}(M)$ , and  $C = \text{ran}(f)$ . Also, depending on whether Step 7(b\*) is executed,  $M_3 = M_2 + v_{I_{\dot{Y}}}$  or  $M_3 = M_2 \cup \{v_{I_X}, v_{\bar{I}_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\}$ . The following table summarizes some of the notation from the algorithm before Step 11.

$i$	$X \in \mathcal{U}_i$	$X \in \mathcal{U}_i$	$X \in \mathcal{R}_i$	$M_i$
1	$ I_X  = 2$	$l(I_X), l(\bar{I}_X) \geq k$	$ I_X  = 3$	$\{v_{I_X} : X \in \mathcal{U}_2 \cup \mathcal{R}_1\}$
2	$ I_X  = 3$	$l(I_X) \geq u_2$	$ I_X  = 2$	$M_1 \cup \{v_{I_X} : X \in \mathcal{U}_3 \cup \mathcal{R}_3\}$
3	$ I_X  = 2$	$l(I_X) \geq l - k + u_2 + u_3$	$ I_X  = 2$	$M_2 \cup \{v_{I_X}, v_{\bar{I}_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\}$ or $M_2 + v_{\bar{I}_{\dot{Y}}}$

TABLE 1. Notation and facts from the algorithm



Our next task is to state and prove the four lemmas on which the algorithm is based. The first lemma is used for Step 7. The statement of the lemma uses the notation from that step.

**Lemma 10.** *Every 2-set  $I \subset X \in \mathcal{R}_3$  satisfies*

$$(3.8) \quad W(\bar{I}_X + v_{I_X}) \geq 3k - b + \Delta_1^1(I_X) - u_2 - r_1.$$

Furthermore, for all  $Y \in \mathcal{R}_3$ :

(a) For every 2-set  $I_Y \subset Y$  with  $L^1(I_Y) \neq \emptyset$  and  $\Delta_1^1(I_Y) \geq 0$  there is a family  $\mathcal{I} = \{I_X : X \in \mathcal{R}_3\}$  with  $I_Y \in \mathcal{I}$  such that  $I_X \subseteq X$ ,  $|I_X| = 2$ , and  $\Delta_1^1(I_X) \geq 0$  for all  $X \in \mathcal{R}_3$ ; and  $\mathcal{L}(M_1 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\})$  has an SDR  $f_2$  extending  $f_1$ .

(b) Additionally, if (H)  $W(\bar{I} \cup L(I)) \leq 3k - 1 - u_2 - r_1$  for all  $X \in \mathcal{R}_3$  and all 2-sets  $I \subset X$ , then for some 2-set  $I_Y \subset Y$ , the family  $\mathcal{I}$  and SDR  $f_2$  from part (a) can be chosen so that there is an SDR  $f_3$  of  $\mathcal{L}(M_1 \cup \{v_{I_X} : X \in \mathcal{R}_3 \cup \mathcal{U}_3\} + v_{\bar{I}_Y})$  extending  $f_2$ .

*Proof.* Consider any  $X = xyzw \in \mathcal{R}_3$ . For  $I \subseteq X$ , let  $l^2(I) = |L(I) \cap C_1|$ . Then  $l(I) = l^1(I) + l^2(I)$ . First we prove (3.8) with  $I = xy$ . By (3.3),  $l^1(wxy) = 0 = l^1(zxy)$ , and by Lemma 9,  $l(X) = 0$ , and so  $l^2(xy) - l^2(wxy) - l^2(zxy) \geq 0$ . Thus

$$\begin{aligned} |W(wz + v_{xy})| &= l^1(w) + l^1(z) + l^1(xy) - l^1(wz) - l^1(wxy) - l^1(zxy) + \\ &\quad l^2(w) + l^2(z) + l^2(xy) - l^2(wz) - l^2(wxy) - l^2(zxy) \\ &\geq 2l + \Delta_1^1(xy) - l^2(wz) \geq 3k - b + \Delta_1^1(xy) - u_2 - r_1. \end{aligned}$$

Let  $A(X) = N_1(X) \setminus C_1$ . By (3.3),  $N_3(X) \subseteq C_1$ . So  $A(X)$  is the set of colors available for coloring a merged pair of vertices from  $X$ , and  $L^1(I) = L(I) \setminus C_1 = L(I) \cap A(X)$  for all pairs  $I \subseteq X$ . Moreover  $\{L^1(I) : I \subseteq X \wedge |I| = 2\}$  is a partition of  $A(X)$ . For each color  $\alpha \in A(x)$ , set  $I(\alpha) = \{x \in X : \alpha \in L(x)\}$ . As  $A(X) \subseteq N_2(X)$ ,  $|I(\alpha)| = 2$ . Let  $B(X) = \{\alpha \in A(X) : \Delta_1^1(I(\alpha)) \geq 0\}$ . Colors  $\beta \in B(X)$ , do not conflict with  $f_1$  and can be representatives for  $L(I(\beta))$ , while  $\Delta_1^1(I(\beta)) \geq 0$ .

Setting  $X = wx_1x_2x_3$ , and using  $N_3(X) \subseteq C_1$ ,

$$(3.9) \quad |A(X)| = \sum_{i=1}^3 (l^1(wx_i) + l^1(\overline{wx}_i)) \leq 2 \sum_{i=1}^3 \max(l^1(wx_i), l^1(\overline{wx}_i)) = 2|B(X)|.$$

By (2.1)

$$(3.10) \quad n_2(X) + 2n_3(X) \geq 4l - |W(X)| \geq 4(l - k) + 1 \geq 2k - 1.$$

As  $N_3(X) \subseteq C_1$ ,  $n_3(X) \leq |C_1| = u_2 + r_1$ , and

$$(3.11) \quad \begin{aligned} |A(X)| &\geq n_2(X) + n_3(X) - |C_1| \geq n_2(X) + 2n_3(X) - n_3(X) - |C_1| \\ &\geq 2k - 1 - (2u_2 + 2r_1) \geq 2r_3 + 2u_3 - 1. \end{aligned}$$

By (3.9),  $|B(X)| \geq \lceil |A(X)|/2 \rceil \geq r_3 + u_3$ .

For (a), we construct the family  $\mathcal{I}$  and SDR  $f_2$  by a greedy algorithm. Start with the special class  $Y \in \mathcal{R}_3$ , and its preassigned subset  $I_Y$ . As  $L^1(I_Y) \neq \emptyset$ , and  $\Delta_1^1(I_Y) \geq 0$  there is a color  $\alpha \in L^1(I_Y) \cap B(Y)$ . Let  $f_2(L(v_{I_Y})) = \alpha$ . Next process the  $X \in \mathcal{R}_3 - Y$  one at a time. When  $X$  is considered, at most  $r_3 - 1$  of the  $r_3 + u_3$  colors of  $B(X)$  have been used. Let  $\beta$  be an unused color, set  $I_X = I(\beta)$ , and put  $f_2(L(v_{I_X})) = \beta$ . Finally consider the  $Z \in \mathcal{U}_3$ . Recall that  $I_Z$  has been assigned in Step 3 so that  $l(I_Z) \geq l - k + u_2 + u_3 \geq u_2 + r_1 + u_3 + r_3$ . So there is a color  $\gamma \in L(I_Z) \setminus C_1$  that has not been used for any previous choices. Set  $f_2(L(v_{I_Z})) = \gamma$ .

For (b), suppose (H) holds. Then  $\Delta_1^1(I) = 0$  for all  $X \in \mathcal{R}_3$  and all 2-sets  $I \subset X$ . Thus  $|A(X)|$  is even and  $A(X) = B(X)$  for all  $X \in \mathcal{R}_3$ . Again we use a greedy procedure.

First choose representatives for each  $L(v_{I_Z})$  with  $Z \in \mathcal{U}_3$ . Also, for each representative  $\alpha$  of  $L(v_{I_Z})$ ,  $Z \in \mathcal{U}_3$ , remove  $\alpha$  from all lists  $L^1(I)$ ,  $I \subset X \in \mathcal{R}_3$ ; and for bookkeeping also remove some additional colors so that for the new lists  $L^-(I)$ ,  $I \subset xyzw := Y$  satisfy

$$\begin{aligned} |L^-(xy)| &= |L^-(wz)|, |L^-(xz)| = |L^-(wy)|, |L^-(xw)| = |L^-(yz)|, \text{ and} \\ r_3 &= |L^-(wx)| + |L^-(wy)| + |L^-(wz)|. \end{aligned}$$

Finish the construction by first choosing a 2-set  $I_Y \subset Y$  with  $L^-(I_Y) \neq \emptyset$ , and setting  $f_2(L(v_{I_Y})) = \alpha \in L^-(I_Y)$  and  $f_3(L(v_{\bar{I}_Y})) = \beta \in L^-(\bar{I}_Y)$ . Then for each  $X \in \mathcal{R}_3 - Y$ , greedily choose a 2-set  $I_X \subset X$  so that  $L^-(I_X)$  has an unused color  $\gamma$  and set  $f_2(L(v_{I_X})) = \gamma$ . This is possible since  $\sum_{I \subset X, |I|=2} |L^-(I)| = 2r_3$ .  $\square$

The next lemma is used in Step 9. The statement of the lemma uses the notation from that step. We will need the following easy claim.

**Claim 11.** Let  $\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3$  be the three partitions of a 4-set  $X$  into pairs. For all  $I_1 \in \mathcal{P}_1, I_2 \in \mathcal{P}_2, I_3 \in \mathcal{P}_3$  there exists  $v \in X$  such that either (i)  $v \in I_1 \cap I_2 \cap I_3$  or (ii)  $v \notin I_1 \cup I_2 \cup I_3$ .

**Lemma 12.** *There is a family  $\mathcal{I} = \{I_X : X \in \mathcal{R}_2\}$  such that  $I_X \subseteq X$  and  $|I_X| = 2$  for all  $X \in \mathcal{R}_2$ , and  $\mathcal{L}(M_2 \cup \{v_{I_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\} \cup \{v_{\bar{I}_X} : X \in \mathcal{U}_1 \cup \mathcal{R}_2\})$  has an SDR  $f_3$  that extends  $f_2$ .*

*Proof.* If  $r_2 = 0 = u_1$  then there is nothing to prove; else Step 7(b\*) is not executed. Each  $X \in \mathcal{U}_1$  satisfies  $L(I_X), L(\bar{I}_X) \geq k$  by Step 2 and  $L(I_X) \cap L(\bar{I}_X) = \emptyset$  by Lemma 9. Thus  $|L(I_X) \setminus C_2|, |L(\bar{I}_X) \setminus C_2| \geq k - u_2 - u_3 - r_1 - r_3 \geq u_1$ . By Theorem 3,  $\{L(I_X) \setminus C_2, L(\bar{I}_X) \setminus C_2 : X \in \mathcal{U}_1\}$  has an SDR, and so  $f_2$  can be extended to an SDR  $g$  for  $\mathcal{L}(M'_2)$ , where  $M'_2 := M_2 \cup \{I_X, \bar{I}_X : X \in \mathcal{U}_1\}$ . Let

$$(3.12) \quad C^g = \text{ran}(g).$$

Then  $|C^g| = 2u_1 + u_2 + u_3 + r_1 + r_3$ .

Next consider any  $X \in \mathcal{R}_2$ . Let  $A(X) = N_2(X) \setminus C^g$ . Again by Theorem 3 it suffices to show:

$$(3.13) \quad (\exists I_X \subseteq X)[|I_X| = 2 \wedge |L(I_X) \cap A(X)| \geq r_2 \wedge |L(\bar{I}_X) \cap A(X)| \geq r_2].$$

Observe  $\sigma_2(X) = n_2(X) + 3n_3(X)$  and  $\sigma_3(X) = n_3(X)$ . So  $n(X) = n_2(X) + n_3(X) = \sigma_2(X) - 2\sigma_3(X)$ . By (3.3),  $N_3(X) \subseteq C^g$  and by (3.4)  $\sigma_3(X) \leq u_2 + r_1$ . So by (3.6),

$$(3.14) \quad \begin{aligned} n(X) &= \sigma_2(X) - 2\sigma_3(X) \geq 5(l - k) + 2u_1 + 2u_2 + u_3 + r_1 + r_2 - (u_2 + r_1) \\ &\geq 5(l - k) + 2u_1 + u_2 + u_3 + r_2 \end{aligned}$$

and

$$(3.15) \quad \begin{aligned} |A(X)| &= |N_2(X) \setminus C^g| = |(N_2(X) \cup N_3(X)) \setminus C^g| \geq n(X) - |C^g| \\ &\geq 5(l - k) + 2u_1 + u_2 + u_3 + r_2 - (2u_1 + u_2 + u_3 + r_1 + r_3) \\ &\geq 5(l - k) - r_1 + r_2 - r_3 \geq 4(l - k) + 2r_2. \end{aligned}$$

Suppose (3.13) fails. Then for each of the three partitions of  $X$  into pairs, there is a pair  $uv$  with  $|L(uv) \cap A(X)| \leq r_2 - 1$ . Using Claim 11, there exists  $w \in X$  such that either (i)  $|L(vw) \cap A(X)| \leq r_2 - 1$  for all  $v \in X - w$  or (ii)  $|L(uw) \cap A(X)| \leq r_2 - 1$  for all  $u, v \in X - w$ . For all  $v \in X - w$ ,

$$(3.16) \quad \text{(a) } |L(v) \cap N(X)| \leq l \text{ and (b) } |L(v) \cap N(X)| \leq \sum_{u \in X - v} l(uw).$$

If (i) holds then

$$(3.17) \quad |L(w) \cap N(X)| \leq |C^g| + \sum_{v \in X-w} |L(vw) \cap A(X)| \leq |C^g| + 3r_2 - 3.$$

Using (3.17) and (3.16a),

$$(3.18) \quad 2n(X) \leq |L(w) \cap N(X)| + \sum_{v \in X-w} |L(v) \cap N(X)| \leq (|C^g| + 3r_2 - 3) + 3l.$$

By (3.14), (3.12) and (3.18) we have

$$(3.19) \quad \begin{aligned} 10(l-k) + 4u_1 + 2u_2 + 2u_3 + 2r_2 &\leq 3l + 2u_1 + u_2 + u_3 + r_1 + r_3 + 3r_2 - 3 \\ (6l - 9k + 3) + 2u_1 + u_2 + u_3 &\leq k - l + r_1 + r_2 + r_3 = 0. \end{aligned}$$

Since  $6l - 9k = -3b$ , both  $b = 1$  and  $0 = u_1 = u_2 = u_3$ . By (3.5),

$$\mu_2(X) \leq l - k + u_2 + u_3 = l - k.$$

Using this with (3.16b) in (3.18) to strengthen the estimate in (3.19) yields the contradiction:

$$\begin{aligned} 10(l-k) + 2r_2 &\leq 9(l-k) + (|C^g| + 3r_2 - 3) \\ l - k &\leq r_1 + r_2 + r_3 - 3 < l - k. \end{aligned}$$

Thus (ii) holds. So

$$(3.20) \quad |A(X)| \leq l(w) + \sum_{uv \subseteq X-w} |L(uv) \cap A(X)| \leq l + 3(r_2 - 1).$$

Using (3.15) and (3.20),

$$\begin{aligned} 4(l-k) + 2r_2 &\leq |A(X)| \leq l + 3(r_2 - 1) \\ 3l - 4k + 3 &\leq r_2. \end{aligned}$$

As  $2l - 3k = -b$ , this yields the contradiction  $l - k + 2 \leq r_2 \leq l - k$ .  $\square$

The next Lemma is used in Step 11(a). Recall that  $\dot{Z}$  is defined in Step 3, and  $\Delta_1$  and  $\Delta_2$  are defined in Step 5.

**Lemma 13.** *Suppose  $X = xyzw \in \mathcal{U}_4$  and  $|W(X)| < |G'|$ . Then  $b = 1$ ,  $u_1 = 0 = r_3$ ,  $u_2 + u_3 \geq 1$ , and there exists a pair  $J \subseteq X$  such that:*

- (1)  $L(J) \not\subseteq C_3$ , and so  $f_3$  can be extended to an SDR of  $\mathcal{L}(M_3 + v_J)$ ;
- (2) for both  $v \in \bar{J}$ , both  $|W(\{v_J, v\})| \geq 2k - 1$  and if  $|W(\{v_J, v\})| = 2k - 1$  then  $|W(\{v_J, v\} \cup \dot{Z}') \cup C_3| \geq 2k$ ;
- (3)  $|W(\bar{J} + v_J)| \geq |G'| - 1$ ; in particular  $|W(X)| \geq |G'| - 1$ .

*Proof.* Now  $|G'| = 3k - u_1 - u_2 + u_4 - r_1 - r_2$ . Observe that

$$(3.21) \quad \sigma_2(X) - \sigma_3(X) \geq 5(l-k) + 2u_1 + 2u_2 + u_3 + r_1 + r_2 + 1,$$

since otherwise inclusion-exclusion yields the contradiction:

$$\begin{aligned} |W(X)| &= \sigma_1(X) - \sigma_2(X) + \sigma_3(X) \\ &\geq 4l - 5(l-k) - 2u_1 - 2u_2 - u_3 - r_1 - r_2 \\ &\geq 3k + (2k - l - u_1 - u_2 - u_3) - u_1 - u_2 - r_1 - r_2 \\ &\geq 3k + u_4 - u_1 - u_2 - r_1 - r_2 = |G'| > |W(X)|. \end{aligned}$$

By (3.21) and (3.6),  $r_1 + r_2 = l - k$  and  $r_3 = 0$ . Consider any pair  $I = xy \subseteq X$ . Then

$$\begin{aligned} |W(\bar{I} + v_I)| &\geq l(z) + l(w) + l(xy) - l(xyz) - l(xyw) - l(zw) \\ &= 2l + \Delta_1(I) - 2u_2 + \Delta_2(I) \\ 1 \leq |G'| - |W(\bar{I} + v_I)| &\leq b - 2u_1 + (u_1 + u_2 + u_4 - l + k) - \Delta_1(I) - \Delta_2(I) \\ (3.22) \qquad \qquad \qquad &= 2b - 2u_1 - u_3 - \Delta_1(I) - \Delta_2(I); \end{aligned}$$

$$(3.23) \qquad \Delta_1(I) + \Delta_2(I) \leq 2b - 1 - 2u_1 - u_3, \text{ and so}$$

$$(3.24) \qquad \Delta_1(\bar{I}) + \Delta_2(\bar{I}) \leq 2b - 1 - 2u_1 - u_3.$$

As  $\Delta_1(I) = -\Delta_1(\bar{I})$ , and  $\Delta_2(I), \Delta_2(\bar{I}) \geq 0$  by (3.2), either  $\Delta_1(I) + \Delta_2(I) \geq 0$  or  $\Delta_1(\bar{I}) + \Delta_2(\bar{I}) \geq 0$ . So  $b = 1$ ,  $u_1 = 0$ ,  $u_3 \leq 1$ , and

$$(3.25) \qquad 0 \leq 4u_2 - \sigma_3(X) = \Delta_2(I) + \Delta_2(\bar{I}) = \Delta_1(I) + \Delta_2(I) + \Delta_1(\bar{I}) + \Delta_2(\bar{I}) \leq 2.$$

By (3.21),  $r_1 + r_2 = l - k$ ,  $\sigma_2(X) \leq 6\mu_2(X)$ , (3.5), and  $\sigma_3 = 4u_2 - \Delta_2(I) - \Delta_2(\bar{I})$ ,

$$(3.26) \qquad 1 + 6(l - k) + 2u_2 + u_3 + \sigma_3(X) \leq \sigma_2(X) \leq 6(l - k + u_2 + u_3)$$

$$(3.27) \qquad 1 + u_3 + 6(l - k + u_2) - \Delta_2(I) - \Delta_2(\bar{I}) \leq \sigma_2(X) \leq 6(l - k + u_2 + u_3).$$

By (3.26)  $u_2 + u_3 \geq 1$ . So the first three assertions of the lemma have been proved. It remains to find a pair  $J \subseteq X$  satisfying (1–3).

First suppose  $u_3 = 1$ . By (3.23),  $\Delta_1(I) + \Delta_2(I) = 0$  for all pairs  $I \subseteq X$ . So  $\Delta_1(I) \leq 0$  and  $\Delta_1(\bar{I}) \leq 0$ . As  $\Delta_1(I) = -\Delta_1(\bar{I})$ , this implies  $\Delta_1(I) = 0 = \Delta_1(\bar{I})$ . Thus  $\Delta_2(I) = 0 = \Delta_2(\bar{I})$ . By (3.27) and (3.5), there exists a pair  $I \subseteq X$  with  $l(I) \geq l - k + u_2 + u_3$ . As  $\Delta_1(I) = 0$ ,  $l(\bar{I}) \geq l - k + u_2 + u_3$ . Thus, using Lemma 9 and (3.2),

$$|W(\{v_I, v_{\bar{I}}\})| = l(I) + l(\bar{I}) \geq 2(l - k + u_2 + u_3) > 2(l - k) + u_2 + u_3 \geq |C_3|.$$

Pick  $J \in \{I, \bar{I}\}$  such that  $L(J) \not\subseteq C_3$ . Then (1) holds. For (2), let  $v' \in \bar{J}$ . Using (3.2),

$$|W(\{v_J, v'\})| = l(J) + l(v') - l(J + v') \geq 2l - k + u_2 + u_3 - u_2 \geq 2k.$$

Thus (2) holds. As  $u_3 = 1$ , (3.22) implies (3).

Otherwise  $u_3 = 0$ . Then  $u_2 \geq 1$ , and so  $\dot{Z} \neq \emptyset$ . Put  $C_0 := C_3 \cup W(\dot{Z}')$ . By Step 3 and Lemma 9,  $|C_0| \geq |W(\dot{Z}')| \geq l + u_2$ . Call a vertex  $x \in X$  *bad* if  $|L(x) \cup C_0| \leq 2k - 1$ ; otherwise  $x$  is *good*. If  $x$  is bad then  $|C_0 \setminus L(x)| \leq 2k - 1 - l = l - k$ . If another vertex  $y$  is also bad, then using (3.5) and  $b = 1$ ,

$$\begin{aligned} l - k + u_2 \geq l(xy) &\geq |L(xy) \cap C_0| \geq |C_0| - |C_0 \setminus L(x)| - |C_0 \setminus L(y)| \\ &\geq l + u_2 - 2(l - k) \geq l - k + u_2 + 1, \end{aligned}$$

a contradiction. So at most one vertex of  $X$  is bad.

Call a pair  $I \subseteq X$  *bad* if  $L(I) \subseteq C_3$ ; otherwise  $I$  is *good*. Note that a good pair satisfies (1). Using (3.25), (3.27) and  $u_3 = 0$  yields

$$(3.28) \qquad 6(l - k + u_2) - 1 \leq \sigma_2 \leq 6(l - k + u_2)$$

so by (3.5), every pair  $I \subseteq X$  satisfies

$$l - k + u_2 - 1 \leq l(I) \leq l - k + u_2.$$

If the upper bound is sharp then call  $I$  *normal*; otherwise call  $I$  *abnormal*. By (3.28) there is at most one abnormal pair. If  $I$  is normal then  $l(\bar{I}) \leq l(I)$ ; so  $\Delta_1(I) \geq 0$ .

By (3.2), every triple  $T \subseteq X$  satisfies  $l(T) \leq u_2$ . If equality holds then call  $T$  *normal*; otherwise call  $T$  *abnormal*; if  $l(T) \leq u_2 - 2$  then call  $T$  *very abnormal*. Suppose two pairs  $I, J \subseteq T$  are both bad. At least one, say  $I$ , is normal. As  $u_1 = 0 = u_3$ ,

$$(3.29) \quad 2(l - k) + u_2 \geq |C_3| \geq |L(I) \cup L(J)| \geq l - k + u_2 + l(J) - l(I \cup J)$$

$$l(T) = l(I \cup J) \geq l(J) - l + k = \begin{cases} u_2 & \text{if } J \text{ is normal} \\ u_2 - 1 & \text{if } J \text{ is abnormal} \end{cases}.$$

So an abnormal triple contains at most one bad, normal pair, and a very abnormal triple contains at most one bad pair.

A pair  $I$  contained in an abnormal triple satisfies  $\Delta_2(I) \geq 1$ . Suppose  $J$  is a good, normal pair contained in an abnormal triple  $T$  with  $w \in X \setminus T$ . Then  $\Delta_1(J) + \Delta_2(J) \geq 1$ . So  $J$  satisfies (3) by (3.22). Also, for both  $v \in X \setminus J$ ,

$$|W(v_J, v)| = l(J) + l(v) - l(J + v) \geq \begin{cases} 2l - k + u_2 - (u_2 - 1) = 2k & \text{if } v \in T \setminus J \\ 2l - k + u_2 - u_2 = 2k - 1 & \text{if } v = w \end{cases}.$$

We have proved the following observation.

*Observation 1.* If  $J \subseteq T \subseteq X$  and  $w \in X \setminus T$ , where  $J$  is a good, normal pair and  $T$  is an abnormal triple, then  $J$  satisfies (1–3) provided  $w$  is good or  $C_0 \not\subseteq W(v_J, w)$ .

By  $u_3 = 0$ , (3.27) and (3.25),  $1 \leq \Delta_2(I) + \Delta_2(\bar{I}) \leq 2$ . As  $\sigma_3 = 4u_2 - \Delta_2(I) - \Delta_2(\bar{I})$ , we have  $\sigma_3(X) = 4u_2 - 2$  or  $\sigma_3(X) = 4u_2 - 1$ . In the first case either there are two abnormal triples or there is a very abnormal triple. In the second case, there is one abnormal triple, and there are no abnormal pairs since equality holds in (3.27).

First suppose there are two abnormal triples. Choose an abnormal triple  $T$  so that if there is a bad vertex then it is in  $T$ . As  $T$  contains three pairs, of which at most one is abnormal, and at most one is bad and normal,  $T$  contains a good, normal pair  $J$ . Say  $J = yz$ ,  $T = xyz$ , and  $w \in X \setminus T$ . Then  $w$  is good, and thus  $J$  satisfies (1–3) by Observation 1.

Otherwise, let  $T = xyz$  be the only abnormal triple and  $w \in X \setminus T$ . There is at most one abnormal pair, and only if  $T$  is very abnormal. So  $T$  contains at most one bad pair.

Now suppose  $T$  has two good, normal pairs  $xy$  and  $yz$ . By Observation 1, some  $J \in P := \{xy, yz\}$  satisfies (1–3), unless  $C_0 \subseteq L(J) \cup L(w)$  for both  $J \in P$ . Then

$$C_0 \subseteq (L(xy) \cup L(w)) \cap (L(yz) \cup L(w)) = L(T) \cup L(w).$$

As  $T$  is abnormal, and using  $u_1 = u_3 = r_3 = 0$ , this yields the contradiction

$$l + u_2 \leq |C_0| \leq |L(T) \cup L(w)| = u_2 - 1 + l.$$

Otherwise,  $T$  does not contain two good, normal pairs. So  $T$  is very abnormal. As  $T$  has at least two normal pairs and at most one bad pair, it has a bad, normal pair (say)  $xz$  and a good normal pair (say)  $J = yz$  is normal. Since  $xz$  is bad,  $L(xz) \subseteq C_0$ . Now

$$\begin{aligned} |C_0 \setminus W(\{v_J, w\})| &\geq |L(xz) \setminus (L(w) \cup L(J))| \geq l(xz) - l(xzw) - l(xzy) \\ &\geq l - k + u_2 - u_2 - (u_2 - 2) \geq 2, \end{aligned}$$

and so  $C_0 \not\subseteq W(v_J, w)$ . Thus  $J$  satisfies (1–3) by Observation 1.  $\square$

The next Lemma is needed for Step 11(b).

**Lemma 14.** *Suppose  $b = 1$  and  $X = xyzw \in \mathcal{U}_4$ . If*

$$(3.30) \quad |W(xyz)| \leq 2k + u_4 - 1 < |W(X)|$$

*then  $u_1 = 0$  and there exists a pair  $J \subseteq X$  such that:*

- (1)  $L(J) \not\subseteq C_3$ , and so  $f_3$  can be extended to an SDR of  $\mathcal{L}(M_3 + v_J)$ ;
- (2)  $|W(\{v_J, v\})| \geq 2k$  for  $v \in xyz \setminus J$  and  $|W(\{v_J, w\})| \geq 2k - 1 + u_3$ ; and
- (3)  $|W(\bar{J} + v_J)| \geq 2k + u_4$ .

*Proof.* Consider a pair  $vv' \subseteq xyz$ . By (3.5) and (3.30),

$$\begin{aligned} 2k + u_4 - 1 &\geq |W(xyz)| \geq |W(vv')| \geq l(v) + l(v') - l(vv') \\ &\geq 2l - (l - k + u_2 + u_3) \geq 3k - 1 - k + u_1 + u_4 \\ &\geq 2k + u_1 + u_4 - 1. \end{aligned}$$

So  $u_1 = 0$ ,  $l(vv') = l - k + u_2 + u_3$ , and  $W(xyz) = W(vv')$ . Since  $vv'$  is arbitrary, every color in  $W(xyz)$  appears in at least two of the lists  $L(x)$ ,  $L(y)$ ,  $L(z)$ . So  $W(\{v_J, v\}) = W(xyz)$  and  $|W(\{v_J, v\})| \geq 2k$  for every pair  $J \subseteq xyz$  and vertex  $v \in xyz \setminus J$ . As  $|C_3| < 2k \leq |W(xyz)|$ , there is a pair  $J \subseteq xyz$  with  $L(J) \not\subseteq C_3$ . Furthermore, by (3.2),

$$|W(\{v_J, w\})| \geq l(J) + l(w) - l(J + w) \geq l - k + u_2 + u_3 + l - u_2 = 2k - 1 + u_3.$$

Finally, as  $W(\{v_J, v\}) = W(xyz)$  for  $v \in xyz \setminus J$ , and using (3.30),

$$|W(\bar{J} + v_J)| = |W(\{v_J, v\}) \cup W(w)| = |W(xyzw)| \geq 2k + u_4. \quad \square$$

The next lemma completes the proof of our main theorem. The reader should keep Figure 3.1 and Table 1 in mind.

**Lemma 15.**  $G'$  is  $L$ -choosable.

*Proof.* First observe that if  $k$  is even then  $b = \dot{u} = \ddot{u} = \dot{r} = 0$  and  $H = G'$ . In this case the following argument is much simpler.

We will prove  $\mathcal{L}(V(H))$  has an SDR. Using Hall's Theorem it suffices to show  $|S| \leq |W| := |\bigcup_{x \in S} L(x)|$  for every  $S \subseteq V(H)$ . Suppose for a contradiction that  $|S| > |W|$  for some  $S \subseteq V(H)$ . We consider several cases. Each case assumes the previous cases fail.

**Case 1:** There is  $X \in \mathcal{U}_4$  with  $|S \cap X| = 4$ . Since  $|W| < |S| \leq |H| \leq |G'|$ , Lemma 13 yields  $b = 1$  and  $|G'| - 1 = |W(X)| < |S| \leq |H|$ . Furthermore, Step 11(a) is executed, and so  $|H| \leq |G'| - 1$ , a contradiction.

**Case 2:** There exists  $X = xyzw \in \mathcal{U}_3$  with  $|S \cap X| = 3$ . Since Case 1 fails,

$$(3.31) \quad |S| \leq |H| \leq 3k - u_1 - u_2 - r_1 - r_2 - \dot{r}.$$

Say  $I_X = xy$ . By Step 5,  $\Delta_1(xy) \geq 0$  and  $l(xyz) + l(xyw) = 2u_2 - \Delta_2(xy)$ . By (3.4),  $l(xyz) + l(xyw) \leq u_2 + r_1$  if  $r_1 < l - k$ ; and it also holds if  $r_1 = l - k$  since  $u_2 \leq 2k - l - u_3 \leq l - k$  and  $l(xyz) + l(xyw) \leq 2u_2$ . Thus

$$\begin{aligned} (3.32) \quad |W| &\geq |W(X')| \geq l(xy) + l(z) + l(w) - l(xyz) - l(xyw) - l(zw) \\ &= 2l + \Delta_1(xy) - 2u_2 + \Delta_2(xy) = 3k - b + \Delta_1(xy) - 2u_2 + \Delta_2(xy) \\ &\geq 3k - b + \Delta_1(xy) - u_2 - r_1 \geq |S| - b \geq |W|. \end{aligned}$$

So equality holds throughout. Thus (3.31) is sharp,  $b = 1$ ,  $u_1 = r_2 = \dot{r} = \Delta_1(xy) = 0$ ,

$$|W| = 3k - 1 - u_2 - r_1,$$

and  $r_1 \leq u_2$  since  $\Delta_2 \geq 0$ . So Step 7(a) is not executed. Since  $\dot{r} = 0$ , Step 7(b\*) is not executed. As  $u_2 = 0 = r_1$ ,  $r_3 = 0$  and Step 7 is degenerate. As  $X \in \mathcal{U}_3$ ,

$$l - k = r_1 \leq u_2 \leq 2k - l - u_3 - u_4 \leq l - k - u_4.$$

So  $u_4 = 0$ , and by Step 5,

$$k = l - k + u_2 + u_3 \leq l(xy) = l(\bar{xy}) + \Delta_1(xy) = l(\bar{xy}).$$

By (3.1) this contradicts  $u_1 = 0$ .

**Case 3:** There exists  $X = wxyz \in \mathcal{R}_3$  with  $|S \cap X'| = 3$ . Say  $I_X = xy$ . As the previous cases fail,  $|S| \leq 3k - u_1 - u_2 - u_3 - r_1 - r_2 - \dot{r}$ . By Step 7,  $\Delta_1^1(xy) \geq 0$ , and by Lemma 10,

$$|W| \geq |W(X')| \geq 3k - b + \Delta_1^1(xy) - u_2 - r_1 \geq |S| - b \geq |W|.$$

Thus  $b = 1$ ,  $0 = r_2 = u_1 = \dot{r}$ ,  $|S| = 3k - 1 - u_2 - r_1$ , and  $\dot{Y}' \subseteq S$ . As  $\dot{r} = 0$ , Step 7(b\*) is not executed. Thus, as  $0 = r_2 = u_1$  and  $r_3 \neq 0$ , Step 7(a) is executed. So  $\Delta_1^1(I_{\dot{Y}}) \geq 1$ . This yields the contradiction

$$|W| \geq |W(\dot{Y}')| \geq 3k - 1 + \Delta_1^1(I_{\dot{Y}}) - u_2 - r_1 \geq 3k - u_2 - r_1.$$

**Case 4:** There exists  $X \in \mathcal{U}_4$  with  $|S \cap X'| = 3$ . As the previous cases fail,

$$|S| \leq 2k + u_4 - \dot{u} - \ddot{u}$$

Let  $xy \subseteq (S \cap X') \setminus M$ . By (3.5),

$$\begin{aligned} |W| &\geq l(x) + l(y) - l(xy) \geq 3k - b - (l - k + u_2 + u_3) \\ &= 2k + (2k - l) - (u_2 + u_3) - b \geq 2k + u_1 + u_4 - b \geq |S| - b \geq |W|. \end{aligned}$$

So  $b = 1$ ,  $0 = u_1 = \dot{u} = \ddot{u}$ ,  $|W| = 2k + u_4 - 1$ , and  $|S| = 2k + u_4$ . Thus  $S$  has exactly two vertices in every class  $X'$  with  $X \in \mathcal{P} \setminus \mathcal{U}_4$  and exactly three vertices in every class  $X'$  with  $X \in \mathcal{U}_4$ . In particular,  $\dot{Z}' \subseteq S$ . Since  $\dot{u} = \ddot{u} = 0$ , we have  $X = X'$ . As Step 11(a) is not executed,  $|W(X)| \geq |G'| \geq |S| = 2k + u_4$ . Thus, as Step 11(b) is not executed, we have the contradiction

$$|W| \geq |W((S \cap X) \cup \dot{Z}')| \geq 2k + u_4 = |S|.$$

**Case 5:** There exists  $X \in \mathcal{U}_1$  with  $|S \cap X'| = 2$ . Then  $v_{I_X}, v_{\bar{I}_X} \in S$ . As the previous cases fail,  $|S| \leq 2k$ . Now

$$|W| \geq L(v_{I_X}) + L(v_{\bar{I}_X}) \geq 2k \geq |S|.$$

**Case 6:** There exists  $X \in \mathcal{U}_3$  with  $|S \cap X'| = 2$ . Say  $S \cap X' = vv'$ . As the previous cases fail,  $|S| \leq 2k - u_1$ . If (say)  $v, v' \notin M$  then  $\bar{I}_X = vv'$ . By (3.1),  $l(\bar{I}_X) \leq k - 1$ . So

$$|W(vv')| \geq l(v) + l(v') - l(vv') \geq 2l - (k - 1) \geq 2k \geq |S|.$$

Otherwise (say)  $v = v_{xy}$ ; so  $I_X = xy$  and  $v' = z \notin M$ . By Step 5,  $l(v_{xy}) \geq l - k + u_2 + u_3$ , and  $l(xyz) \leq u_2$  by (3.2). So

$$\begin{aligned} |W(vv')| &\geq l(v_{xy}) + l(z) - l(xyz) \\ &\geq l - k + u_2 + u_3 + l - u_2 \geq 2k - b + u_3 \geq 2k \geq |S|. \end{aligned}$$

**Case 7:** There exists  $X \in \mathcal{U}_4$  with  $|S \cap X'| = 2$ . Say  $S \cap X' = vv'$ . If possible, choose  $X$  so that  $S \cap X' \cap M = \emptyset$ . As the previous cases fail,  $|S| \leq 2k - u_1 - u_3$ . If  $v, v' \notin M$  then, using (3.2),

$$\begin{aligned} |W(vv')| &= l(v) + l(v') - l(vv') \geq 2l - (l - k + u_2 + u_3) \\ (3.33) \quad &= 2k - b + u_1 + u_4 \geq 2k \geq |S|. \end{aligned}$$

Else  $b = 1$ , and (say)  $v \in M$ . By Step 11, either  $v = v_{I_X}$  or  $v = v_{\bar{I}_X}$ . By Lemmas 13 and 14,  $u_1 = 0$ .

If  $v = v_{I_X}$  then Step 11(a) was executed. So (i)  $u_1 = 0 = r_3$ , (ii)  $|W(vv')| \geq 2k - 1$ , and (iii) if  $|W(vv')| = 2k - 1$  then  $|W(vv' \cup \dot{Z}') \cup C_3| \geq 2k$ . Since

$$2k \geq |S| > |W| \geq |W(vv')| \geq 2k - 1,$$

we have  $|S| = 2k$  and  $u_3 = 0$ . Thus  $S$  contains exactly two vertices of each part  $Y' \in \mathcal{P}'$ . As at most one class of  $\mathcal{U}_4$  has merged vertices, the choice of  $X$  implies  $u_4 = 1$ ; thus

$u_2 = l - k$ . Also,  $\dot{Z}' \subseteq S$ . Since  $u_3 = 0 = r_3$ ,  $M_3 \subseteq S$ . So  $|W| \geq |W(vv' \cup \dot{Z}') \cup C_4| \geq 2k$ , a contradiction.

Otherwise  $v = v_{I_{\ddot{X}}}$ . Then Step 11(b) was executed. As only one part in  $\mathcal{U}_4$  can have contracted vertices,  $X = \ddot{X} = xyzw \in \mathcal{U}_4$  with (say)  $I_{\ddot{X}} = xy$ ,

$$|W(xyz \cup \dot{Z}')| \leq 2k + u_4 - 1 < |W(\ddot{X})|,$$

$|W(\{v_{xy}, z\})| \geq 2k$  and  $|W(\{v_{xy}, w\})| \geq 2k - 1 + u_3$ . So we have the contradiction  $|W(vv')| \geq 2k$  unless  $v' = w$  and

$$2k \geq |S| > |W(\{v_{xy}, w\})| \geq 2k - 1 + u_3.$$

Thus  $u_1 = 0 = u_3$  and  $|S| = 2k$ . Again  $S$  contains exactly two vertices of each class  $Y' \in \mathcal{P}'$ , and the choice of  $X$  implies  $u_4 = 1$ . So  $u_2 = l - k$ . Also,  $\dot{Z}' \subseteq S$ . As  $|W(\ddot{X})| > |W(xyz \cup \dot{Z}')|$ , we have  $|L(w) \setminus W(xyz \cup \dot{Z}')| \geq 1$ . So we have the contradiction

$$|W| \geq |W(\{v_{xy}, w\} \cup \dot{Z}')| \geq |W(\dot{Z}')| + 1 = l + u_2 + 1 = 2l - k + 1 = 2k.$$

**Case 8:** There exists  $X = xyzw \in \mathcal{U}_2$  with  $|S \cap X'| = 2$ . Say  $S \cap X' = \{v_I, w\}$ . As the previous cases fail,  $|S| \leq 2k - u_1 - u_3 - u_4 = l + u_2$ . Since  $L(xyz) \cap L(w) = \emptyset$ , we have

$$|W| \geq |W(X')| \geq l(xyz) + l(w) \geq u_2 + l \geq |S|.$$

**Case 9:** Otherwise. As the previous cases fail,

$$|S| \leq u_1 + u_2 + u_3 + u_4 + 2|\mathcal{R}| = l.$$

As  $\mathcal{L}(M)$  has an SDR, there is a vertex  $x \in S \setminus M$ . Thus  $|W| \geq l(x) = l \geq |S|$ . □

This completes the proof of Theorem 7. □

#### 4. ON-LINE CHOOSABILITY

By Theorem 7,  $\text{ch}(K_{4*3}) = 4$ . Using a computer we have checked that  $\text{ch}^{\text{ol}}(K_{4*3}) = 5$ , but do not have a readable argument to verify the upper bound. Here we prove the lower bound.

**Theorem 16.**  $\text{ch}^{\text{ol}}(K_{4*3}) \geq 5$ .

*Proof.* Figure 4.1 describes a strategy for Alice. The top left matrix depicts the initial game position, and Alice's first move. The positions in the matrix correspond to the vertices of  $K_{4*3}$  arranged so that vertices in the same part correspond to positions in the same column. The order of vertices within a column is irrelevant, as is the order of the columns. The numbers represent the size of the list of each corresponding vertex. The sequence of numbers represents a function  $f$ . The shaded positions represent the vertices that Alice presents on her first move.

As play progresses Bob chooses certain vertices presented by Alice and passes over others. When a vertex is chosen its position is removed from the next matrix (and the positions in its column of the remaining vertices and the order of the columns may be rearranged). When he passes over a vertex its list size is decreased by one (and its position in its column and the order of the columns may change). The arrows between the matrices point to the possible new game positions that arise from Bob's choice, not counting equivalent positions and omitting clearly inferior positions for Bob. In particular we assume Bob always chooses a maximal independent set.

For example, after Bob's first move there is only one possible game position, provided Bob chooses a maximal independent set. It is shown in the second column of the first row,






FIGURE 4.1. Strategy for Alice demonstrating  $\text{ch}^{\text{ol}}(K_{4*3}) \geq 5$ .

along with Alice's second move. Now Bob has two possible responses that are pointed to by two arrows. Also consider the matrix in the third row and third column. There are three nonequivalent responses for Bob, but choosing the offered vertex in the second column of the matrix results in a position that is inferior to choosing the offered vertex in the first column. So this option is not shown.

Eventually, Alice forces one of five positions  $(G, f)$  such that  $G$  is not  $f$ -choosable, and Bob, being a gentleman, resigns.  $\square$

**Acknowledgment.** The authors would like to thank two exceptionally thorough referees whose suggestions greatly improved the presentation.

#### REFERENCES

1. Noga Alon, *Choice numbers of graphs: a probabilistic approach.*, *Combinatorics, Probability and Computing* **1** (1992), 107–114.
2. ———, *Restricted colorings of graphs*, *Surveys in combinatorics, 1993* (Keele), *London Math. Soc. Lecture Note Ser.*, vol. 187, Cambridge Univ. Press, Cambridge, 1993, pp. 1–33.
3. R. L. Brooks, *On colouring the nodes of a network*, *Proc. Cambridge Philos. Soc.* **37** (1941), 194–197.
4. Paul Erdős, Arthur L. Rubin, and Herbert Taylor, *Choosability in graphs*, *Congress. Numer.*, XXVI (1980), 125–157.
5. Fred Galvin, *The list chromatic index of a bipartite multigraph*, *J. Combin. Theory Ser. B* **63** (1995), no. 1, 153–158.
6. Nurit Gazit and Michael Krivelevich, *On the asymptotic value of the choice number of complete multi-partite graphs*, *J. Graph Theory* **52** (2006), no. 2, 123–134.
7. H.A. Kierstead, *On the choosability of complete multipartite graphs with part size three*, *Discrete Mathematics* **211** (2000), 255–259.
8. Jakub Kozik, Piotr Micek, and Xuding Zhu, *Towards an on-line version of Ohba's conjecture*, *European Journal of Combinatorics* **36** (2014), no. 0, 110–121.
9. Jonathan A. Noel, Bruce A. Reed, and Hehui Wu, *A proof of a conjecture of Ohba*, *J. Graph Theory* (2014), in press.

10. Jonathan A. Noel, Douglas B. West, Hehui Wu, and Xuding Zhu, *Beyond Ohba's conjecture: a bound on the choice number of  $k$ -chromatic graphs with  $n$  vertices*, European J. Combin. **43** (2015), 295–305. MR 3266299
11. Kyoji Ohba, *Choice number of complete multipartite graphs with part size at most three.*, Ars Comb. **72** (2004), 133–139.
12. Bruce A. Reed and Benny Sudakov, *List colouring of graphs with at most  $(2 - o(1))\chi$  vertices*, Proceedings of the International Congress of Mathematicians **Vol. III** (2002), 587–603.
13. ———, *List colouring when the chromatic number is close to the order of the graph.*, Combinatorica **25** (2004), no. 1, 117–123.
14. Uwe Schauz, *Mr. Paint and Mrs. Correct*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 77, 18.
15. Carsten Thomassen, *Every planar graph is 5-choosable*, J. Combin. Theory Ser. B **62** (1994), no. 1, 180–181.
16. V. G. Vizing, *Coloring the vertices of a graph in prescribed colors*, Diskret. Analiz (1976), no. 29 Metody Diskret. Anal. v Teorii Kodov i Shem, 3–10, 101.
17. ———, *Coloring the vertices of a graph in prescribed colors*, Diskret. Analiz (1976), no. 29 Metody Diskret. Anal. v Teorii Kodov i Shem, 3–10, 101.
18. Margit Voigt, *List colourings of planar graphs*, Discrete Math. **120** (1993), no. 1–3, 215–219.
19. D. Yang, *Extension of the game coloring number and some results on the choosability of complete multipartite graphs*, Ph.D. thesis, Arizona State University, 2003.
20. Xuding Zhu, *On-line list colouring of graphs*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 127, 16.

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287, USA

*E-mail address:* `kierstead@asu.edu`

6328 E. CACTUS WREN RD., PARADISE VALLEY, AZ 85253, USA

*E-mail address:* `andrew.salmon@yale.edu`

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287, USA

*E-mail address:* `rwang31@asu.edu`