

A General Mass Transference Principle

Demi Allen* Simon Baker†

March 7, 2018

Abstract

In this paper we prove a general form of the Mass Transference Principle for lim sup sets defined via neighbourhoods of sets satisfying a certain local scaling property. Such sets include self-similar sets satisfying the open set condition and smooth compact manifolds embedded in \mathbb{R}^n . Our main result is applicable in locally compact metric spaces and allows one to transfer Hausdorff g -measure statements to Hausdorff f -measure statements. This work extends previous results of this type in several distinct directions.

2000 *Mathematics Subject Classification*: Primary 11J83, 28A78; Secondary 11K60.

Keywords and phrases: Mass Transference Principle, Hausdorff measures, lim sup sets, Diophantine approximation.

1 Introduction

1.1 Background

Within Diophantine approximation and metric number theory, many sets of interest can be characterised as lim sup sets. Recall that, given a countable collection of sets $(E_j)_{j \in \mathbb{N}}$, we define the corresponding lim sup set to be

$$\begin{aligned} \limsup E_j &:= \bigcap_{j=1}^{\infty} \bigcup_{n=j}^{\infty} E_n \\ &= \{x : x \in E_j \text{ for infinitely many } j \in \mathbb{N}\}. \end{aligned}$$

Often one is interested in determining the metric properties of $\limsup E_j$. When the sequence $(E_j)_{j \in \mathbb{N}}$ is a collection of balls, a useful tool in determining the metric properties of $\limsup E_j$ is the Mass Transference Principle [5]. This result allows one to determine Hausdorff measure statements from seemingly less general Lebesgue measure statements.

*EPSRC Doctoral Prize Fellow supported by grant EP/N509565/1

†Supported by EPSRC grant EP/M001903/1

Given a ball $B := B(x, r)$ in \mathbb{R}^n and a dimension function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ (see Section 2 for definitions), define another corresponding ball $B^f := B(x, f(r)^{\frac{1}{n}})$. When $f(r) = r^s$ for some $s > 0$, we write B^s in place of B^f . In particular, $B^n = B$. The following was established by Beresnevich and Velani in [5].

Mass Transference Principle. *Let $(B_j)_{j \in \mathbb{N}}$ be a sequence of balls in \mathbb{R}^n with $r(B_j) \rightarrow 0$ as $j \rightarrow \infty$. Let f be a dimension function such that $x^{-n}f(x)$ is monotonic and suppose that, for any ball B in \mathbb{R}^n ,*

$$\mathcal{H}^n(B \cap \limsup_{j \rightarrow \infty} B_j^f) = \mathcal{H}^n(B) .$$

Then, for any ball B in \mathbb{R}^n ,

$$\mathcal{H}^f(B \cap \limsup_{j \rightarrow \infty} B_j^k) = \mathcal{H}^f(B) .$$

We denote by $\mathcal{H}^f(X)$ the Hausdorff f -measure of a set $X \subset \mathbb{R}^n$. For $s \geq 0$, $\mathcal{H}^s(X)$ denotes the standard Hausdorff s -measure of X . These notions will be formally introduced in Section 2. Although, it is worth noting at this point that if X is a Borel subset of \mathbb{R}^n , then $\mathcal{H}^n(X)$ is a constant multiple times the n -dimensional Lebesgue measure of X (see [9] for further details). Thus, as discussed previously, the Mass Transference Principle genuinely does enable us to transfer Lebesgue measure statements to Hausdorff measure ones.

The Mass Transference Principle was originally motivated by a desire to establish a Hausdorff measure analogue of the famous Duffin–Schaeffer Conjecture in metric number theory and has some other surprising consequences in Diophantine approximation, see for example [2, 4, 5] and references therein.

A generalisation of the Mass Transference Principle, which is applicable in locally compact metric spaces and not restricted to lim sup sets in \mathbb{R}^n , was also given by Beresnevich and Velani in [5]. Before stating this result formally, we require some preliminaries.

We say that a function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is doubling if there exists a constant $\lambda > 1$ such that

$$f(2x) < \lambda f(x) \tag{1}$$

for all $x > 0$. Here $\mathbb{R}^+ := [0, \infty)$. Let (X, d) be a locally compact metric space and let g be a doubling dimension function. Moreover, suppose that there exist constants $0 < c_1 < 1 < c_2 < \infty$ and $r_0 > 0$ such that

$$c_1 g(r) \leq \mathcal{H}^g(B(x, r)) \leq c_2 g(r) \tag{2}$$

for all $x \in X$ and $0 < r < r_0$. Here, \mathcal{H}^g denotes the Hausdorff g -measure defined in Section 2. Given another dimension function f and a ball $B := B(x, r)$ we define

$$B^{f,g} := B(x, g^{-1}(f(r))) .$$

The following theorem was established in [5].

Theorem MTP*. *Let (X, d) and g be as above and let $(B_j)_{j \in \mathbb{N}}$ be a sequence of balls in X with $r(B_j) \rightarrow 0$ as $j \rightarrow \infty$. Let f be a dimension function such that f/g is monotonic and suppose that for any ball B in X we have*

$$\mathcal{H}^g(B \cap \limsup_{j \rightarrow \infty} B_j^{f,g}) = \mathcal{H}^g(B).$$

Then, for any ball B in X ,

$$\mathcal{H}^f(B \cap \limsup_{j \rightarrow \infty} B_j) = \mathcal{H}^f(B).$$

Compared with the Mass Transference Principle, this general theorem applies to lim sup sets of balls in more general metric spaces. As one example, Theorem MTP* is applicable when X is, say, the middle-third Cantor set. In this case, Theorem MTP* has been used in [15] as a tool for addressing a problem posed by Mahler regarding the existence of very well approximable points in the middle-third Cantor set.

In the Euclidean setting, Beresnevich and Velani also extended the Mass Transference Principle in another direction to deal with lim sup sets determined by sequences of neighbourhoods of “approximating” planes. The result they obtained in this case, [6, Theorem 3], carried an extra technical conditions as a result of the “slicing” technique used for the proof. Recently, the first author and Beresnevich have removed these additional constraints in [1]. This improvement gives rise to a number of previously inaccessible applications in Diophantine approximation which are also explored in [1]. For completeness, we remark here that, in the Euclidean setting, progress towards proving a mass transference principle for rectangles has been made by Wang, Wu and Xu in [19].

In this paper, we extend the results of [1, 5, 6] by proving a general version of the Mass Transference Principle that applies to lim sup sets in a locally compact metric space which are defined in terms of sets satisfying a certain local scaling property. Our main result, Theorem 1 below, extends the known mass transference principles of [1, 5, 6] in several manners. First of all, while it incorporates mass transference principles for balls and approximating planes, Theorem 1 is also applicable to more exotic sets. For example, we are able to consider lim sup sets generated by sequences of neighbourhoods of smooth compact manifolds or self-similar fractals satisfying the open set condition. Furthermore, as long as the local scaling property is satisfied, the sets generating a lim sup set need not all be of the same type (e.g. all balls or all planes). Secondly, we deal with lim sup sets in a locally compact metric space (X, d) and are not constrained to the Euclidean setting. Finally, the result we derive allows us to transfer Hausdorff g -measure statements to Hausdorff f -measure statements, where f and g are dimension functions subject to some mild conditions. Aside from Theorem MTP*, the previous results mentioned above all deal with transferring Lebesgue measure statements (or, more precisely, Hausdorff n -measure statements) to Hausdorff f -measure statements for lim sup sets in Euclidean space.

1.2 The Main Result

Let (X, d) be a locally compact metric space. Given $F \subset X$ and $\delta \geq 0$, we define the δ -neighbourhood of F to be

$$\Delta(F, \delta) := \{x \in X : d(x, F) < \delta\}.$$

Let g be a dimension function satisfying (2). Given a sequence of sets $\mathcal{F} := (F_j)_{j \in \mathbb{N}}$ in X and $0 \leq \kappa < 1$, we say that \mathcal{F} satisfies the *local scaling property (LSP) with respect to κ* if there exist constants $c_3, c_4, r_1 > 0$ such that, for any $0 < r < r_1$, $\delta < r$, $j \in \mathbb{N}$ and $x \in F_j$, we have

$$c_3 g(\delta)^{1-\kappa} \cdot g(r)^\kappa \leq \mathcal{H}^g(B(x, r) \cap \Delta(F_j, \delta)) \leq c_4 g(\delta)^{1-\kappa} \cdot g(r)^\kappa. \quad (3)$$

If (3) is satisfied for one specific set F , say, we will also say that F satisfies the local scaling property with respect to κ . It should be clear from the context when we are referring to an individual set and when we are referring to a sequence of sets.

Restricting to the case where $X = \mathbb{R}^n$ and $\mathcal{H}^g = \mathcal{H}^n$ (that is, essentially, Lebesgue measure) then (3) takes the form

$$c_3 \delta^{n-\kappa n} \cdot r^{\kappa n} \leq \mathcal{H}^n(B(x, r) \cap \Delta(F_j, \delta)) \leq c_4 \delta^{n-\kappa n} \cdot r^{\kappa n}. \quad (4)$$

Many familiar subsets of Euclidean space satisfy (4) for an appropriate choice of κ . For example, in Section 5 we show that any smooth compact manifold embedded in \mathbb{R}^n satisfies (4). We also show that any self-similar set satisfying the open set condition satisfies the LSP. In these examples we will see that κ is related to the box counting dimension of the sets we are considering. In Section 1.3, we provide some discussion as to why this should be the case. In the meantime, we present here two trivial examples of sets satisfying the LSP.

Example 1. If \mathcal{F} were a sequence of points, then it follows from (2) that \mathcal{F} satisfies the LSP with respect to $\kappa = 0$.

Example 2. Suppose $F \subset \mathbb{R}^n$ is a specific set satisfying (4). One can then define a sequence $\mathcal{F} := (F_j)_{j \in \mathbb{N}}$ by defining each F_j to be the image of F under some isometry. Clearly \mathcal{F} provides us with a sequence of sets satisfying the LSP.

Next, suppose $(\Upsilon_j)_{j \in \mathbb{N}}$ is a sequence of non-negative reals such that $\Upsilon_j \rightarrow 0$ as $j \rightarrow \infty$. Consider the lim sup set

$$\Lambda(\Upsilon) := \{x \in X : x \in \Delta(F_j, \Upsilon_j) \text{ for infinitely many } j \in \mathbb{N}\}.$$

Our main result is the following theorem.

Theorem 1. *Let (X, d) and g be as above. Let $\mathcal{F} := (F_j)_{j \in \mathbb{N}}$ be a sequence of sets satisfying the LSP with respect to some $0 \leq \kappa < 1$, and let $\Upsilon := (\Upsilon_j)_{j \in \mathbb{N}}$ be a sequence of non-negative reals such that $\Upsilon_j \rightarrow 0$ as $j \rightarrow \infty$. Let f be*

a dimension function such that f/g is monotonic and f/g^κ is a dimension function. Suppose that for any ball B in X we have

$$\mathcal{H}^g \left(B \cap \Lambda \left(g^{-1} \left(\left(\frac{f(\Upsilon)}{g(\Upsilon)^\kappa} \right)^{\frac{1}{1-\kappa}} \right) \right) \right) = \mathcal{H}^g(B). \quad (5)$$

Then, for any ball B in X we have

$$\mathcal{H}^f(B \cap \Lambda(\Upsilon)) = \mathcal{H}^f(B).$$

Taking $(F_j)_{j \in \mathbb{N}}$ to be a sequence of points in X and $\kappa = 0$, as in Example 1, Theorem 1 coincides with Theorem MTP* given above. If, further, we insist that $X = \mathbb{R}^n$ and $g(r) = r^n$, we recover the usual Mass Transference Principle. One can recover [1, Theorem 1] by taking $X = \mathbb{R}^n$, $(F_j)_{j \in \mathbb{N}}$ to be a sequence of l -dimensional planes in \mathbb{R}^n , $\kappa = \frac{l}{n}$, and $g(r) = r^n$ in Theorem 1.

We believe that the greater generality of Theorem 1 should open up a number of new interesting applications compared with its predecessors. Within Diophantine approximation, as one example, we suspect that Theorem 1 may enable the establishment of further Hausdorff measure statements relating to approximation on manifolds. For an overview of some existing results of this type we refer the reader to [3, 4] and references within. In addition to its potential uses in Diophantine approximation, Theorem 1 lends itself naturally to wider applications. Indeed, we will provide an example in Section 6 which shows how we may use Theorem 1 to deduce Hausdorff measure and dimension results for certain random lim sup sets.

Restricting our attention to Euclidean space and s -dimensional Hausdorff measures, Theorem 1 takes the following simpler form.

Corollary 1. *Let $\mathcal{F} := (F_j)_{j \in \mathbb{N}}$ be a sequence of subsets of \mathbb{R}^n satisfying the LSP with respect to some $0 \leq \kappa < 1$ and let $\Upsilon := (\Upsilon_j)_{j \in \mathbb{N}}$ be a sequence of non-negative reals such that $\Upsilon_j \rightarrow 0$ as $j \rightarrow \infty$. Let $s > \kappa n$ and suppose that for any ball B in \mathbb{R}^n we have*

$$\mathcal{H}^n(B \cap \Lambda(\Upsilon^{\frac{s-\kappa n}{(1-\kappa)n}})) = \mathcal{H}^n(B).$$

Then, for any ball B in X we have

$$\mathcal{H}^s(B \cap \Lambda(\Upsilon)) = \mathcal{H}^s(B).$$

1.3 Box counting dimension, Minkowski content and the LSP

In this section, we highlight the connection between (4) and the Minkowski/box counting dimension of a set and why these quantities are likely to be related to κ for sets satisfying the LSP.

Instead of the usual box counting definition of lower and upper box counting dimension (see, for example, [9]), there is the following equivalent notion that is

defined using the volume of a δ neighbourhood. Given a bounded set $F \subset \mathbb{R}^n$, the lower and upper box counting dimension of F are defined to be, respectively,

$$\underline{\dim}_B(F) := n - \limsup_{\delta \rightarrow 0} \frac{\log \mathcal{H}^n(\Delta(F, \delta))}{\log \delta}$$

and

$$\overline{\dim}_B(F) := n - \liminf_{\delta \rightarrow 0} \frac{\log \mathcal{H}^n(\Delta(F, \delta))}{\log \delta}.$$

When these limits coincide we call the common value the box counting dimension of F and denote it by $\dim_B(F)$. It follows from this definition that if F were a set whose box counting dimension exists, and we were interested in whether (4) held for some κ , the natural candidate for κ would be $\dim_B(F) \cdot n^{-1}$.

Two other useful quantities that describe how the volume of a δ neighbourhood of a set F scales are the lower and upper Minkowski content. Given a bounded set F contained in \mathbb{R}^n whose box counting dimension exists, we define the lower and upper Minkowski content to be, respectively,

$$\underline{M}(F) := \liminf_{\delta \rightarrow 0} \delta^{n - \dim_B(F)} \cdot \mathcal{L}^n(\Delta(F, \delta))$$

and

$$\overline{M}(F) := \limsup_{\delta \rightarrow 0} \delta^{n - \dim_B(F)} \cdot \mathcal{L}^n(\Delta(F, \delta)).$$

Here \mathcal{L}^n is the n -dimensional Lebesgue measure. When $\underline{M}(F) = \overline{M}(F)$ we call the common value the Minkowski content and denote it by $M(F)$. Determining conditions under which a set F has both $\underline{M}(F)$ and $\overline{M}(F)$ positive and finite is a well studied problem, see [14]. Equation (4) doesn't follow directly from the positivity and finiteness of both $\underline{M}(F)$ and $\overline{M}(F)$. However, if F were reasonably homogeneous in small neighbourhoods, one would expect this property to be sufficient to deduce (4).

1.4 Structure of the paper

The rest of the paper is arranged as follows. In Section 2 we recall some useful preliminaries from geometric measure theory. In Section 3 we prove some key technical results which will be required throughout the proof of Theorem 1. The proof of Theorem 1 will be presented in full in Section 4. In Section 5 we give some examples of sets satisfying the LSP, and in Section 6 we use Theorem 1 to deduce the Hausdorff dimension and measures of some random lim sup sets.

2 Preliminaries

In this section we state some definitions and recall some well known facts from geometric measure theory. Throughout this paper we will say that $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a *dimension function* if f is a left continuous, non-decreasing function such

that $f(r) \rightarrow 0$ as $r \rightarrow 0$. Given a ball $B := B(x, r)$ in X and a dimension function f , we define

$$V^f(B) := f(r).$$

The Hausdorff f -measure with respect to the dimension function f is defined as follows. Suppose $F \subset X$, let f be a dimension function and let $\rho > 0$. A ρ -cover for F is any countable collection of balls $\{B_i\}_{i \in \mathbb{N}}$ with $r(B_i) < \rho$ for every $i \in \mathbb{N}$ and $F \subset \bigcup_{i \in \mathbb{N}} B_i$. We define

$$\mathcal{H}_\rho^f(F) := \inf \left\{ \sum_i V^f(B_i) : \{B_i\} \text{ is a } \rho\text{-cover for } F \right\}.$$

The Hausdorff f -measure of F with respect to the dimension function f is then defined as

$$\mathcal{H}^f(F) := \lim_{\rho \rightarrow 0} \mathcal{H}_\rho^f(F).$$

A simple consequence of the definition of \mathcal{H}^f is the following useful fact (see, for example, [9]).

Lemma 1. *Let (X, d) be as above. Suppose f and g are dimension functions such that the ratio $f(r)/g(r) \rightarrow 0$ as $r \rightarrow 0$, then $\mathcal{H}^f(F) = 0$ whenever $\mathcal{H}^g(F) < \infty$.*

When $f(r) = r^s$ ($s \geq 0$), the measure \mathcal{H}^f is the familiar s -dimensional Hausdorff measure, which we denote by \mathcal{H}^s . The Hausdorff dimension, $\dim_{\mathbb{H}} F$, of a set F is defined as

$$\dim_{\mathbb{H}}(F) := \inf \{s \geq 0 : \mathcal{H}^s(F) = 0\}.$$

When calculating the Hausdorff dimension of a set, a usual strategy is to obtain upper and lower bounds separately. It is often the case that calculating an upper bound is relatively straightforward while determining a lower bound is much more difficult. Nevertheless, a standard tool which can frequently be employed in obtaining lower bounds for Hausdorff dimension is the following Mass Distribution Principle.

Lemma 2 (Mass Distribution Principle). *Let μ be a probability measure supported on a subset F of X . Suppose there are positive constants c and r_0 such that*

$$\mu(B) \leq c V^f(B)$$

for any ball B with radius $r \leq r_0$. If E is a subset of F with $\mu(E) = \lambda > 0$ then $\mathcal{H}^f(E) \geq \lambda/c$.

For this precise statement of the Mass Distribution Principle, see [5, Section 2]. For further general information regarding Hausdorff measures and dimension we refer the reader to [9, 16].

Let $B := B(x, r)$ be a ball in (X, d) . For any $\alpha > 0$, we denote by αB the ball B scaled by a factor α ; i.e. $\alpha B(x, r) := B(x, \alpha r)$. A useful covering lemma which we will use throughout is the following (see [16]).

Lemma 3 (The $5r$ -covering lemma). *Let (X, d) be a metric space. Every family \mathcal{F} of balls of uniformly bounded diameter in X contains a disjoint subfamily \mathcal{G} such that*

$$\bigcup_{B \in \mathcal{F}} B \subset \bigcup_{B \in \mathcal{G}} 5B.$$

We will also make use of the following adaptation of [7, Lemma 4].

Lemma 4. *Let B be a ball in the locally compact metric space (X, d) and let g be a doubling dimension function. Let $(S_i)_{i \in \mathbb{N}}$ be a sequence of subsets in B and let $(\delta_i)_{i \in \mathbb{N}}$ be a sequence of positive numbers such that $\delta_i \rightarrow 0$ as $i \rightarrow \infty$. Let*

$$\Delta(S_i, \delta_i) := \{x \in X : d(S_i, x) < \delta_i\}.$$

Then, for any real number $C > 1$,

$$\mathcal{H}^g(\limsup_{i \rightarrow \infty} \Delta(S_i, \delta_i)) = \mathcal{H}^g(\limsup_{i \rightarrow \infty} \Delta(S_i, C\delta_i)).$$

Note that [7, Lemma 4] is stated in the setting of Euclidean space. Going through the steps in the proof of this lemma one can verify the above analogue holds in our setting. To prove this analogue we require a notion of the Lebesgue Density Theorem that holds for our metric space (X, d) equipped with the measure \mathcal{H}^g . Such an analogue is known to exist when g satisfies our doubling bound (1), see for example [18].

In what follows we use the Vinogradov notation, writing $A \ll B$ if $A \leq cB$ for some positive constant c and $A \gg B$ if $A \geq c'B$ for some positive constant c' . If $A \ll B$ and $A \gg B$ we write $A \asymp B$ and say that A and B are *comparable*.

3 The $K_{G,B}$ -Lemma

Before proving Theorem 1, we formulate suitable analogues of [1, Lemma 4] and [1, Lemma 5] which will be required in the present setting. Let

$$\tilde{\Upsilon}_j := g^{-1} \left(\left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)^\kappa} \right)^{\frac{1}{1-\kappa}} \right).$$

Given a ball B in X and $j \in \mathbb{N}$, we define

$$\Phi_j(B) := \{B(x, \tilde{\Upsilon}_j) \subset B : x \in F_j\}.$$

The following is the analogue of [1, Lemma 4] or [5, Lemma 5], the so-called $K_{G,B}$ -Lemma, we obtain in the setting currently under consideration.

Lemma 5. *Let (X, d) , \mathcal{F} , Υ , g and f be as given in Theorem 1 and assume that the hypotheses of Theorem 1 hold. Then, for any ball B in X and any $G \in \mathbb{N}$, there exists a finite collection*

$$K_{G,B} \subset \{(A; j) : j \geq G, A \in \Phi_j(B)\}$$

satisfying the following properties:

- (i) if $(A; j) \in K_{G,B}$ then $3A \subset B$;
- (ii) if $(A; j), (A', j') \in K_{G,B}$ are distinct then $3A \cap 3A' \neq \emptyset$; and
- (iii) there exists a constant $c_5 \geq 0$ independent of our choice of ball B such that

$$\mathcal{H}^g \left(\bigcup_{(A;j) \in K_{G,B}} A \right) \geq c_5 \mathcal{H}^g(B).$$

Similarly to [1, Lemma 4], the collection $K_{G,B}$ here is a collection of balls drawn from the families $\Phi_j(B)$. These balls correspond to the lim sup set $\Lambda(\tilde{\Upsilon})$. From each of these balls what we are actually interested in is extracting a suitable collection of balls corresponding to the lim sup set $\Lambda(\Upsilon)$. We adopt the notation from [1] and write $(A; j)$ for a generic ball from $K_{G,B}$ to “remember” the index j of the family $\Phi_j(B)$ that the ball A comes from but just write A if we are referring only to the ball A (as opposed to the pair $(A; j)$). Making such a distinction is necessary for us to be able to choose the “right” collection of balls within A that at the same time lie in an Υ_j -neighbourhood of the relevant F_j . Indeed, for $j \neq j'$ we could have $A = A'$ for some $A \in \Phi_j(B)$ and $A' \in \Phi_{j'}(B)$.

Proof of Lemma 5. For $j \in \mathbb{N}$ and a fixed ball B in X , consider the set of balls

$$\Phi_j^3(B) := \{B(x, 3\tilde{\Upsilon}_j) \subset B : x \in F_j\}.$$

It follows from our assumption (5) that for any $G \geq 1$ we have

$$\mathcal{H}^g \left(\bigcup_{j \geq G} (\Delta(F_j, 3\tilde{\Upsilon}_j) \cap B) \right) = \mathcal{H}^g(B).$$

Observe that $\tilde{\Upsilon}_j \rightarrow 0$ as $j \rightarrow \infty$ because g and f/g^k are dimension functions. Therefore for $j \in \mathbb{N}$ sufficiently large,

$$\bigcup_{L \in \Phi_j^3(B)} L \supset \Delta(F_j, 3\tilde{\Upsilon}_j) \cap \frac{1}{2}B.$$

Therefore for any sufficiently large $G \in \mathbb{N}$, we have

$$\mathcal{H}^g \left(\bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \right) \geq \mathcal{H}^g \left(\bigcup_{j \geq G} \left(\Delta(F_j, 3\tilde{\Upsilon}_j) \cap \frac{1}{2}B \right) \right) = \mathcal{H}^g \left(\frac{1}{2}B \right).$$

Suppose $G' \in \mathbb{N}$ is large enough that the above inequality holds for any $G \geq G'$. Clearly for any $G < G'$ we also have

$$\bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \supset \bigcup_{j \geq G'} \bigcup_{L \in \Phi_j^3(B)} L.$$

Therefore for any $G \in \mathbb{N}$ it follows that

$$\mathcal{H}^g \left(\bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \right) \geq \mathcal{H}^g \left(\frac{1}{2}B \right). \quad (6)$$

Next, by the $5r$ -covering Lemma (Lemma 3), there exists a disjoint subcollection $\mathcal{G} \subset \{(L; j) : j \geq G, L \in \Phi_j^3(B)\}$ satisfying

$$\bigcup_{(L; j) \in \mathcal{G}} L \subset \bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \subset \bigcup_{(L; j) \in \mathcal{G}} 5L.$$

Let $\mathcal{G}' := \{(\frac{1}{3}L; j) : (L; j) \in \mathcal{G}\}$ be the balls from the collection \mathcal{G} all scaled by a factor of $1/3$. Note that the balls in \mathcal{G}' are still disjoint when scaled by 3. By the above, we have that

$$\bigcup_{(A; j) \in \mathcal{G}'} A \subset \bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \subset \bigcup_{(A; j) \in \mathcal{G}'} 15A. \quad (7)$$

It follows from (1), (2), and the disjointness of the balls in \mathcal{G}' that

$$\begin{aligned} \mathcal{H}^g \left(\bigcup_{(A; j) \in \mathcal{G}'} A \right) &= \sum_{(A; j) \in \mathcal{G}'} \mathcal{H}^g(A) \stackrel{(2)}{\asymp} \sum_{(A; j) \in \mathcal{G}'} g(r(A)) \\ &\stackrel{(1)}{\gg} \sum_{(A; j) \in \mathcal{G}'} g(r(15A)) \\ &\stackrel{(2)}{\asymp} \sum_{(A; j) \in \mathcal{G}'} \mathcal{H}^g(15A) \\ &\geq \mathcal{H}^g \left(\bigcup_{(A; j) \in \mathcal{G}'} 15A \right). \end{aligned}$$

Now also using (6) and (7), we see that

$$\begin{aligned} \mathcal{H}^g \left(\bigcup_{(A; j) \in \mathcal{G}'} A \right) &\stackrel{(7)}{\gg} \mathcal{H}^g \left(\bigcup_{j \geq G} \bigcup_{L \in \Phi_j^3(B)} L \right) \\ &\stackrel{(6)}{\geq} \mathcal{H}^g \left(\frac{1}{2}B \right) \\ &\stackrel{(2)}{\asymp} g(r(\frac{1}{2}B)) \\ &\stackrel{(1)}{\gg} g(r(B)) \\ &\stackrel{(2)}{\asymp} \mathcal{H}^g(B). \end{aligned}$$

Thus, there exists a constant $c' > 0$ such that

$$c' \cdot \mathcal{H}^g(B) \leq \mathcal{H}^g \left(\bigcup_{(A;j) \in \mathcal{G}'} A \right).$$

Since the balls in \mathcal{G}' are disjoint and contained in B , it follows that

$$\mathcal{H}^g \left(\bigcup_{\substack{(A;j) \in \mathcal{G}' \\ j \geq N}} A \right) \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Consequently, there must exist $N_0 \in \mathbb{N}$ such that

$$\mathcal{H}^g \left(\bigcup_{\substack{(A;j) \in \mathcal{G}' \\ j \geq N_0}} A \right) < \frac{c'}{2} \mathcal{H}^g(B).$$

We define $K_{G,B}$ to be the subcollection of $(A; j) \in \mathcal{G}'$ with $G \leq j < N_0$. By the above we see that $K_{G,B}$ is a finite collection of balls while still satisfying the required properties (i)–(iii) with $c_5 = \frac{c'}{2}$. \square

As mentioned previously, from each of the balls in $K_{G,B}$ we wish to extract a collection of balls corresponding to $\Lambda(\Upsilon)$. The desired properties and existence of such collections are summarised in the following lemma, which constitutes the required analogue of [1, Lemma 5] in this setting.

Lemma 6. *Let (X, d) , \mathcal{F} , Υ , f , g , and B be as in Lemma 5 and assume that the hypotheses of Theorem 1 hold. Furthermore, assume that $f(r)/g(r) \rightarrow \infty$ as $r \rightarrow 0$. Let $K_{G,B}$ be as in Lemma 5. Then, provided that G is sufficiently large, for any $(A; j) \in K_{G,B}$ there exists a collection $\mathcal{C}(A; j)$ of balls satisfying the following properties:*

- (i) *each ball in $\mathcal{C}(A; j)$ is of radius Υ_j and is centred on F_j ;*
- (ii) *if $L \in \mathcal{C}(A; j)$ then $3L \subset A$;*
- (iii) *if $L, M \in \mathcal{C}(A; j)$ are distinct then $3L \cap 3M = \emptyset$;*
- (iv) $\mathcal{H}^g(\Delta(F_j, \Upsilon_j) \cap \frac{1}{2}A) \ll \mathcal{H}^g \left(\bigcup_{L \in \mathcal{C}(A; j)} L \right) \leq \mathcal{H}^g(\Delta(F_j, \Upsilon_j) \cap A)$; *and*
- (v) *there exist some constants $d_1, d_2 > 0$ such that*

$$d_1 \times \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)} \right)^{\frac{\kappa}{1-\kappa}} \leq \#\mathcal{C}(A; j) \leq d_2 \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)} \right)^{\frac{\kappa}{1-\kappa}}. \quad (8)$$

Proof. We begin by showing that

$$\frac{\Upsilon_j}{\tilde{\Upsilon}_j} \rightarrow 0 \quad \text{as } j \rightarrow \infty. \quad (9)$$

To this end, suppose that $N \in \mathbb{N}$. We aim to show that for all sufficiently large $j \in \mathbb{N}$ we have

$$\Upsilon_j < \frac{\tilde{\Upsilon}_j}{2^N}. \quad (10)$$

Observe that (10) holds if

$$g(\Upsilon_j) < g\left(\frac{\tilde{\Upsilon}_j}{2^N}\right). \quad (11)$$

Furthermore, by repeated application of (1), we see that

$$g\left(\frac{\tilde{\Upsilon}_j}{2^N}\right) > \frac{g(\tilde{\Upsilon}_j)}{\lambda^N} = \frac{1}{\lambda^N} \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)^\kappa}\right)^{\frac{1}{1-\kappa}},$$

where λ is the doubling constant. Consequently, (11) holds if

$$g(\Upsilon_j) < \frac{1}{\lambda^N} \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)^\kappa}\right)^{\frac{1}{1-\kappa}}.$$

Rearranging the above we get

$$\lambda^{N(1-\kappa)} < \frac{f(\Upsilon_j)}{g(\Upsilon_j)}.$$

By the assumptions that $\frac{f(r)}{g(r)} \rightarrow \infty$ as $r \rightarrow 0$ and $\Upsilon_j \rightarrow 0$ as $j \rightarrow \infty$, we see that this inequality holds for sufficiently large $j \in \mathbb{N}$, thus verifying (9).

In light of (9) we can assume that G is sufficiently large so that

$$6\Upsilon_j < \tilde{\Upsilon}_j \quad \text{for any } j \geq G. \quad (12)$$

Let $x_1, \dots, x_t \in F_j \cap \frac{1}{2}A$ be a maximal collection of points such that

$$d(x_i, x_{i'}) > 6\Upsilon_j \quad \text{if } i \neq i'. \quad (13)$$

Define $\mathcal{C}(A; j)$ to be the collection of balls

$$\mathcal{C}(A; j) := \{B(x_1, \Upsilon_j), \dots, B(x_t, \Upsilon_j)\}.$$

By construction, property (i) is satisfied by the collection $\mathcal{C}(A; j)$. Next, recall that $A \in \Phi_j(B)$ and so $\frac{1}{2}A$ has radius $\frac{1}{2}\tilde{\Upsilon}_j$. If $L := B(x_i, \Upsilon_j) \in \mathcal{C}(A; j)$ then any $y \in 3L$ satisfies $d(y, x_i) < 3\Upsilon_j$. Supposing x_0 is the centre of A , we also have $d(x_i, x_0) \leq \frac{1}{2}\tilde{\Upsilon}_j$. Combining (12) and the triangle inequality we obtain

$$d(y, x_0) \leq d(y, x_i) + d(x_i, x_0) \leq 3\Upsilon_j + \frac{1}{2}\tilde{\Upsilon}_j < \tilde{\Upsilon}_j.$$

Therefore property (ii) follows. In addition property (iii) follows from (13).

It is a consequence of the maximality of x_1, \dots, x_t that for any $x \in F_j \cap \frac{1}{2}A$, there exists an x_i from this collection such that $d(x, x_i) \leq 6\Upsilon_j$. Consequently

$$\Delta(F_j, \Upsilon_j) \cap \frac{1}{2}A \subset \bigcup_{L \in \mathcal{C}(A; j)} 7L.$$

Therefore, by (1) and (2),

$$\begin{aligned} \mathcal{H}^g(\Delta(F_j, \Upsilon_j) \cap \frac{1}{2}A) &\leq \mathcal{H}^g\left(\bigcup_{L \in \mathcal{C}(A; j)} 7L\right) \\ &\leq \sum_{L \in \mathcal{C}(A; j)} \mathcal{H}^g(7L) \\ &\ll \sum_{L \in \mathcal{C}(A; j)} \mathcal{H}^g(L) \\ &\ll \mathcal{H}^g\left(\overset{\circ}{\bigcup}_{L \in \mathcal{C}(A; j)} L\right). \end{aligned}$$

However, by property (ii), we have

$$\overset{\circ}{\bigcup}_{L \in \mathcal{C}(A; j)} L \subset \Delta(F_j, \Upsilon_j) \cap A.$$

This together with the previous inequality proves property (iv).

As a byproduct of (12) we have that $\Upsilon_j < \frac{\tilde{\Upsilon}_j}{2}$ for all $j \geq G$. Therefore, by the LSP we have

$$\mathcal{H}^g\left(\Delta(F_j, \Upsilon_j) \cap \frac{1}{2}A\right) \asymp g\left(\frac{\tilde{\Upsilon}_j}{2}\right)^\kappa \cdot g(\Upsilon_j)^{1-\kappa}.$$

Combining this with the doubling property (1) and the LSP we obtain

$$\mathcal{H}^g\left(\Delta(F_j, \Upsilon_j) \cap \frac{1}{2}A\right) \asymp g(\tilde{\Upsilon}_j)^\kappa \cdot g(\Upsilon_j)^{1-\kappa} \asymp \mathcal{H}^g(\Delta(F_j, \Upsilon_j) \cap A). \quad (14)$$

By (2) and the disjointness of the balls in $\mathcal{C}(A; j)$ we have

$$\mathcal{H}^g\left(\overset{\circ}{\bigcup}_{L \in \mathcal{C}(A; j)} L\right) = \sum_{L \in \mathcal{C}(A; j)} \mathcal{H}^g(L) \asymp \sum_{L \in \mathcal{C}(A; j)} g(\Upsilon_j) = \#\mathcal{C}(A; j) g(\Upsilon_j).$$

Combining the above with (14) and property (iv) we get

$$\#\mathcal{C}(A; j) \asymp \frac{g(\tilde{\Upsilon}_j)^\kappa \cdot g(\Upsilon_j)^{1-\kappa}}{g(\Upsilon_j)} = \left(\frac{g(\tilde{\Upsilon}_j)}{g(\Upsilon_j)}\right)^\kappa = \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)}\right)^{\frac{\kappa}{1-\kappa}}$$

So property (v) holds. \square

4 Proof of Theorem 1

4.1 Strategy

Fix an arbitrary ball B_0 in X and suppose the assumptions of Theorem 1 hold. Our goal is to show that

$$\mathcal{H}^f(B_0 \cap \Lambda(\Upsilon)) = \mathcal{H}^f(B_0). \quad (15)$$

Since f/g is monotonic, there are three situations to consider:

- (a) $\frac{f(r)}{g(r)} \rightarrow \infty$ as $r \rightarrow 0$;
- (b) $\frac{f(r)}{g(r)} \rightarrow 0$ as $r \rightarrow 0$; and
- (c) $\frac{f(r)}{g(r)} \rightarrow \ell$ as $r \rightarrow 0$, where $0 < \ell < \infty$.

If we are in situation (b) it follows from Lemma 1 that $\mathcal{H}^f(B_0) = 0$. Since $B_0 \cap \Lambda(\Upsilon) \subset B_0$ the result follows.

If we are in case (c) it can be shown that $\tilde{\Upsilon}_j \asymp \Upsilon_j$. It then follows from Lemma 4 that $\mathcal{H}^g(B_0 \cap \Lambda(\tilde{\Upsilon})) = \mathcal{H}^g(B_0 \cap \Lambda(\Upsilon))$. In turn, it follows from (5) that $\mathcal{H}^g(B_0 \cap \Lambda(\Upsilon)) = \mathcal{H}^g(B_0)$. Finally, the proof is completed in this case by noting that $\mathcal{H}^f = \ell \cdot \mathcal{H}^g$ and therefore

$$\mathcal{H}^f(B_0 \cap \Lambda(\Upsilon)) = \ell \cdot \mathcal{H}^g(B_0 \cap \Lambda(\Upsilon)) = \ell \cdot \mathcal{H}^g(B_0) = \mathcal{H}^f(B_0).$$

It remains to address case (a). Thus, from now on we will assume that $\frac{f(r)}{g(r)} \rightarrow \infty$ as $r \rightarrow 0$. In this case, it is a consequence of Lemma 1 that $\mathcal{H}^f(B_0) = \infty$. So, to prove Theorem 1 it suffices to show that

$$\mathcal{H}^f(B_0 \cap \Lambda(\Upsilon)) = \infty.$$

To achieve this goal we will show that for any $\eta > 1$, we can construct a Cantor set \mathbb{K}_η contained in $B_0 \cap \Lambda(\Upsilon)$ which supports a probability measure μ satisfying

$$\mu(D) \ll \frac{V^f(D)}{\eta}, \quad (16)$$

for all balls D with sufficiently small radii, where the implicit constants are independent of D and η . The result then follows from the Mass Distribution Principle (Lemma 2) upon taking η to be arbitrarily large since the Mass Distribution Principle yields $\mathcal{H}^f(\mathbb{K}_\eta) \geq \eta$ and $\mathbb{K}_\eta \subset B_0 \cap \Lambda(\Upsilon)$.

4.2 Desired properties of \mathbb{K}_η

The construction of the Cantor set we present here is an adaptation of that given in [1] and [5]. For ease of comparison we will generally adopt the notation used in [1].

Fix $\eta > 1$. Our Cantor set \mathbb{K}_η will take the form

$$\mathbb{K}_\eta = \bigcap_{n=1}^{\infty} \mathbb{K}(n)$$

where $\mathbb{K}(n) \supset \mathbb{K}(n+1)$. The fact that (X, d) is a locally compact metric space guarantees that \mathbb{K}_η is non empty.

Each level $\mathbb{K}(n)$ of the Cantor set will be a union of balls and we will denote the corresponding set of level n balls by $K(n)$. For each ball $B \in K(n-1)$ we will construct an (n, B) -local level, henceforth denoted by $K(n, B)$, which will consist of balls contained in B . The set of level n balls, $K(n)$, will then be defined by

$$K(n) := \bigcup_{B \in K(n-1)} K(n, B).$$

Each (n, B) -local level will be constructed of *local sub-levels* and will take the form

$$K(n, B) := \bigcup_{i=1}^{l_B} K(n, B, i), \quad (17)$$

where $K(n, B, i)$ denotes the i th local sub-level and l_B is the number of local sub-levels forming $K(n, B)$. What is more, each local sub-level will take the form

$$K(n, B, i) := \bigcup_{B' \in \mathcal{G}(n, B, i)} \bigcup_{(A; j) \in K_{G', B'}} \mathcal{C}(A; j). \quad (18)$$

Here, $\mathcal{G}(n, B, i)$ will be a suitable collection of balls contained in B and, for each ball $B' \in \mathcal{G}(n, B, i)$, $K_{G', B'}$ will be the corresponding finite collection whose existence is asserted by Lemma 5. The collections $\mathcal{C}(A; j)$ will be those arising from Lemma 6. The set of pairs $(A; j)$ included in (18) will be denoted by $\tilde{K}(n, B, i)$. As such

$$\tilde{K}(n, B, i) := \bigcup_{B' \in \mathcal{G}(n, B, i)} K_{G', B'} \quad \text{and} \quad K(n, B, i) = \bigcup_{(A; j) \in \tilde{K}(n, B, i)} \mathcal{C}(A; j). \quad (19)$$

We will also require that \mathbb{K}_η satisfies the following properties.

The properties of levels and sub-levels of \mathbb{K}_η

(P0) $K(1) = \{B_0\}$.

(P1) For any $n \geq 2$ and $B \in K(n-1)$ the balls

$$\{3L : L \in K(n, B)\}$$

are disjoint and contained in B .

(P2) For any $n \geq 2$, $B \in K(n-1)$, and $i \in \{1, \dots, l_B\}$, the local sub-level $K(n, B, i)$ is a finite union of some collections $C(A; j)$ of balls satisfying properties (i)–(v) of Lemma 6. Moreover, the balls $3A$ are disjoint and contained in B .

(P3) For any $n \geq 2$, $B \in K(n-1)$, and $i \in \{1, \dots, l_B\}$, we have

$$\sum_{(A;j) \in \tilde{K}(n,B,i)} V^g(A) \geq c_6 V^g(B)$$

where

$$c_6 := \frac{1}{2\lambda} \left(\frac{c_1}{c_2} \right)^2 \frac{c_5}{c_7}.$$

The constants c_1 and c_2 are those appearing in (2), c_5 comes from Lemma 5 (iii), λ is the doubling constant associated with g , and c_7 is a fixed constant such that

$$g(r(5B)) \leq c_7 g(r(B))$$

for any ball B in X . Note that the existence of c_7 is guaranteed by the doubling property (1).

(P4) For any $n \geq 2$, $B \in K(n-1)$, $i \in \{1, \dots, l_B - 1\}$, $L \in K(n, B, i)$, and $M \in K(n, B, i+1)$, we have

$$f(r(M)) \leq \frac{f(r(L))}{2} \quad \text{and} \quad \frac{f(r(M))}{g(r(M))^\kappa} \leq \frac{f(r(L))}{2g(r(L))^\kappa}.$$

(P5) The number of sub-levels is defined by

$$l_B := \begin{cases} \left\lceil \frac{c_2 \eta}{c_6 \mathcal{H}^g(B)} \right\rceil + 1 & \text{if } B = B_0 := \mathbb{K}(1); \\ \left\lceil \frac{V^f(B)}{c_6 V^g(B)} \right\rceil + 1 & \text{if } B \in K(n) \text{ with } n \geq 2. \end{cases}$$

and $l_B \geq 2$ for $B \in K(n)$ with $n \geq 2$.

4.3 Existence of \mathbb{K}_η

We now prove that it is possible to construct a set $\mathbb{K}_\eta \subset B_0 \cap \Lambda(\Upsilon)$ satisfying properties **(P0)**–**(P5)**. To this end, let

$$K_l(n, B) := \bigcup_{i=1}^l K(n, B, i) \quad \text{and} \quad \tilde{K}_l(n, B) := \bigcup_{i=1}^l \tilde{K}(n, B, i). \quad (20)$$

Level 1. Let $\mathbb{K}(1) := B_0$ so **(P0)** holds.

All other levels of \mathbb{K}_η are defined inductively. Therefore, assume levels $\mathbb{K}(1), \dots, \mathbb{K}(n-1)$ have been constructed. To construct the n th level, we need to construct (n, B) -local levels for all balls $B \in K(n-1)$.

Level n. Fix $B \in K(n-1)$ and let $\varepsilon := \varepsilon(B)$ be a small constant which will be explicitly determined later. Let G be sufficiently large so Lemma 5 and Lemma 6 can be invoked. We may also assume that G is large enough that

$$3g(\Upsilon_j)^{1-\kappa} < \frac{f(\Upsilon_j)}{g(\Upsilon_j)^\kappa} \quad \text{for all } j \geq G, \quad (21)$$

$$\frac{g(\Upsilon_j)}{f(\Upsilon_j)} < \varepsilon \frac{g(r(B))}{f(r(B))} \quad \text{for all } j \geq G, \quad (22)$$

and

$$\left[\frac{f(\Upsilon_j)}{c_6 g(\Upsilon_j)} \right] \geq 1 \quad \text{for all } j \geq G. \quad (23)$$

Here, c_6 is the constant appearing in **(P3)**. Inequalities (21)–(23) are achievable since $f(r)/g(r) \rightarrow \infty$ as $r \rightarrow 0$.

Recall that the (n, B) -local level, $K(n, B)$, consists of local sub-levels. These are defined as follows.

Sub-level 1. For B and G as above let $K_{G,B}$ be the collection of balls arising from Lemma 5. We define the first sub-level of $K(n, B)$ to be

$$K(n, B, 1) := \bigcup_{(A;j) \in K_{G,B}} C(A; j).$$

Hence,

$$\tilde{K}(n, B, 1) = K_{G,B} \quad \text{and} \quad \mathcal{G}(n, B, 1) = \{B\}.$$

Higher sub-levels. The higher sub-levels are defined inductively. Suppose the first l sub-levels $K(n, B, 1), \dots, K(n, B, l)$ have been constructed and properties **(P1)**–**(P4)** hold with l in place of l_B . Since we require fairly stringent separation conditions between balls in \mathbb{K}_η , we first verify that there is “space” left over in B for the sub-level $K(n, B, l+1)$ after the first l sub-levels, $K(n, B, 1), \dots, K(n, B, l)$, have been constructed. Let

$$A^{(l)} := \frac{1}{2}B \setminus \bigcup_{L \in K_l(n, B)} 4L.$$

We will show that

$$\mathcal{H}^g(A^{(l)}) \geq \frac{1}{2} \mathcal{H}^g\left(\frac{1}{2}B\right). \quad (24)$$

Using (1), (2), and the upper bound for $\#\mathcal{C}(A; j)$ given in (8), we obtain

$$\begin{aligned}
\mathcal{H}^g \left(\bigcup_{L \in K_l(n, B)} 4L \right) &\leq \sum_{L \in K_l(n, B)} \mathcal{H}^g(4L) \\
&\stackrel{(2)}{\leq} \sum_{L \in K_l(n, B)} c_2 g(r(4L)) \\
&\stackrel{(1)}{\leq} \sum_{L \in K_l(n, B)} c_2 \lambda^2 g(r(L)) \\
&= \sum_{i=1}^l \sum_{L \in K(n, B, i)} c_2 \lambda^2 g(r(L)) \\
&= \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} c_2 \lambda^2 \#\mathcal{C}(A; j) g(\Upsilon_j) \\
&\stackrel{(8)}{\leq} \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} c_2 \lambda^2 d_2 \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)} \right)^{\frac{\kappa}{1-\kappa}} g(\Upsilon_j) \\
&= c_2 \lambda^2 d_2 \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} \frac{f(\Upsilon_j)^{\frac{1}{1-\kappa}}}{g(\Upsilon_j)^{\frac{\kappa}{1-\kappa}}} \cdot \frac{g(\Upsilon_j)}{f(\Upsilon_j)}.
\end{aligned}$$

Recalling condition (22) and property **(P2)** we further see that

$$\begin{aligned}
\mathcal{H}^g \left(\bigcup_{L \in K_l(n, B)} 4L \right) &\stackrel{(22)}{<} \varepsilon c_2 \lambda^2 d_2 \cdot \frac{g(r(B))}{f(r(B))} \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} \frac{f(\Upsilon_j)^{\frac{1}{1-\kappa}}}{g(\Upsilon_j)^{\frac{\kappa}{1-\kappa}}} \\
&= \varepsilon c_2 \lambda^2 d_2 \cdot \frac{g(r(B))}{f(r(B))} \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} g(\tilde{\Upsilon}_j) \\
&\stackrel{(2)}{\leq} \varepsilon \frac{c_2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))}{f(r(B))} \sum_{i=1}^l \sum_{(A; j) \in \tilde{K}(n, B, i)} \mathcal{H}^g(A) \\
&\stackrel{\mathbf{(P2)}}{\leq} \varepsilon \frac{c_2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))}{f(r(B))} l \mathcal{H}^g(B) \\
&\leq \varepsilon \frac{c_2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))}{f(r(B))} (l_B - 1) \mathcal{H}^g(B) \\
&\stackrel{(2)}{\leq} \varepsilon \frac{c_2^2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))^2}{f(r(B))} (l_B - 1). \tag{25}
\end{aligned}$$

To establish (24) we will show that

$$\mathcal{H}^g \left(\bigcup_{L \in K_l(n, B)} 4L \right) < \frac{1}{2} \mathcal{H}^g \left(\frac{1}{2} B \right).$$

By (2) we have

$$\frac{1}{2}\mathcal{H}^g\left(\frac{1}{2}B\right) \geq \frac{c_1}{2}g\left(\frac{1}{2}B\right)$$

and by (1) we have

$$g\left(\frac{1}{2}B\right) > \frac{1}{\lambda}g(r(B)).$$

Therefore it suffices to show that

$$\mathcal{H}^g\left(\bigcup_{L \in K_l(n, B)} 4L\right) < \frac{c_1}{2\lambda}g(r(B)). \quad (26)$$

It follows from (25) that (26) is implied by

$$\varepsilon \frac{c_2^2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))^2}{f(r(B))} (l_B - 1) < \frac{c_1}{2\lambda}g(r(B)). \quad (27)$$

Taking

$$\varepsilon(B) := \frac{c_1}{4\lambda} \left(\frac{c_2^2 \lambda^2 d_2}{c_1} \cdot \frac{g(r(B))}{f(r(B))} (l_B - 1) \right)^{-1}$$

we see that (27) is satisfied and, thus, (26) and (24) both hold.

Next, observe that the quantity

$$d_{\min} := \min\{r(L) : L \in K_l(n, B)\}$$

is well-defined and positive since the collection $K_l(n, B)$ is finite. Let

$$\mathcal{A}(n, B, l) := \{B(x, d_{\min}) : x \in A^{(l)}\}.$$

By Lemma 3 there exists a disjoint subcollection $\mathcal{G}(n, B, l+1)$ of $\mathcal{A}(n, B, l)$ such that

$$A^{(l)} \subset \bigcup_{B' \in \mathcal{A}(n, B, l)} B' \subset \bigcup_{B' \in \mathcal{G}(n, B, l+1)} 5B'. \quad (28)$$

Note that each element of the collection $\mathcal{G}(n, B, l+1)$ is a subset of B . Since the balls in this collection are disjoint and all have the same radius, $\mathcal{G}(n, B, l+1)$ must be finite. Furthermore, by our construction,

$$B' \cap \bigcap_{L \in K_l(n, B)} 3L = \emptyset \quad \text{for any } B' \in \mathcal{G}(n, B, l+1). \quad (29)$$

By the above (28) and (24) we have

$$\mathcal{H}^g\left(\bigcup_{B' \in \mathcal{G}(n, B, l+1)} 5B'\right) \geq \mathcal{H}^g(A^{(l)}) \geq \frac{1}{2}\mathcal{H}^g\left(\frac{1}{2}B\right). \quad (30)$$

Since $\mathcal{G}(n, B, l + 1)$ is a disjoint collection of balls we have the following

$$\begin{aligned}
\mathcal{H}^g \left(\bigcup_{B' \in \mathcal{G}(n, B, l+1)} 5B' \right) &\leq \sum_{B' \in \mathcal{G}(n, B, l+1)} \mathcal{H}^g(5B') \\
&\stackrel{(2)}{\leq} c_2 \sum_{B' \in \mathcal{G}(n, B, l+1)} g(r(5B')) \\
&\leq c_2 c_7 \sum_{B' \in \mathcal{G}(n, B, l+1)} g(r(B')) \\
&\stackrel{(2)}{\leq} \frac{c_2 c_7}{c_1} \sum_{B' \in \mathcal{G}(n, B, l+1)} \mathcal{H}^g(B') \\
&= \frac{c_2 c_7}{c_1} \mathcal{H}^g \left(\bigcup_{B' \in \mathcal{G}(n, B, l+1)} \overset{\circ}{B}' \right).
\end{aligned}$$

Combining this with (30) from above implies

$$\mathcal{H}^g \left(\bigcup_{B' \in \mathcal{G}(n, B, l+1)} \overset{\circ}{B}' \right) \geq \frac{c_1}{2c_2 c_7} \mathcal{H}^g \left(\frac{1}{2} B \right). \quad (31)$$

Now, to construct the $(l + 1)$ th sub-level $K(n, B, l + 1)$, let $G' \geq G$ be sufficiently large so that we can apply Lemmas 5 and 6 to each ball $B' \in \mathcal{G}(n, B, l + 1)$. Moreover, we assume that G' is sufficiently large so that for every $j \geq G'$,

$$f(\Upsilon_j) \leq \frac{1}{2} \min_{L \in K_l(n, B)} f(r(L)) \quad \text{and} \quad \frac{f(\Upsilon_j)}{g(\Upsilon_j)^\kappa} \leq \frac{1}{2} \min_{L \in K_l(n, B)} \frac{f(r(L))}{g(r(L))^\kappa}. \quad (32)$$

Such a G' exists since there are only finitely many balls in $K_l(n, B)$, $\Upsilon_j \rightarrow 0$ as $j \rightarrow \infty$, and because f and f/g^κ are dimension functions.

To each ball $B' \in \mathcal{G}(n, B, l + 1)$ we apply Lemma 5 to obtain a collection of balls $K_{G', B'}$. We then define

$$K(n, B, l + 1) := \bigcup_{B' \in \mathcal{G}(n, B, l+1)} \bigcup_{(A; j) \in K_{G', B'}} \mathcal{C}(A; j).$$

Consequently,

$$\tilde{K}(n, B, l + 1) = \bigcup_{B' \in \mathcal{G}(n, B, l+1)} K_{G', B'}.$$

As $G' \geq G$, properties (21)–(23) remain valid. We now verify that properties **(P1)**–**(P5)** hold for this local sub-level.

To prove **(P1)** holds we first observe that it is satisfied for balls in $\bigcup_{(A; j) \in K_{G', B'}} \mathcal{C}(A; j)$ by the properties of $\mathcal{C}(A; j)$ and the fact that the balls

in $K_{G',B'}$ are disjoint. The balls in $K_{G',B'}$ are by definition contained in B' and the balls $B' \in \mathcal{G}(n, B, l+1)$ are disjoint, therefore **(P1)** is satisfied for all balls L in $K(n, B, l+1)$. Last of all, combining this observation with (29) we can conclude that **(P1)** is satisfied for all balls L in $K_{l+1}(n, B)$. Property **(P2)** is satisfied for this sub-level because of Lemma 5 (i) and (ii) and because the balls $B' \in \mathcal{G}(n, B, l+1)$ are disjoint.

We now prove that **(P3)** still holds for $i = l+1$. Recalling (2), we have

$$\begin{aligned} \sum_{(A;j) \in \tilde{K}(n, B, l+1)} V^g(A) &= \sum_{B' \in \mathcal{G}(n, B, l+1)} \sum_{(A;j) \in K_{G', B'}} V^g(A) \\ &\stackrel{(2)}{\geq} \frac{1}{c_2} \sum_{B' \in \mathcal{G}(n, B, l+1)} \sum_{(A;j) \in K_{G', B'}} \mathcal{H}^g(A). \end{aligned}$$

Combining this with Lemma 5 (iii) and the fact that the balls in $\mathcal{G}(n, B, l+1)$ are disjoint, we see that

$$\begin{aligned} \sum_{(A;j) \in \tilde{K}(n, B, l+1)} V^g(A) &\geq \frac{1}{c_2} \sum_{B' \in \mathcal{G}(n, B, l+1)} c_5 \mathcal{H}^g(B') \\ &= \frac{c_5}{c_2} \mathcal{H}^g \left(\bigcup_{B' \in \mathcal{G}(n, B, l+1)} B' \right) \\ &\stackrel{(31)}{\geq} \frac{c_1}{2c_2c_7} \frac{c_5}{c_2} \mathcal{H}^g \left(\frac{1}{2}B \right) \\ &\stackrel{(2)}{\geq} \frac{c_1}{2c_2c_7} \frac{c_5}{c_2} c_1 g \left(r \left(\frac{1}{2}B \right) \right) \\ &\stackrel{(1)}{\geq} \frac{1}{2\lambda} \left(\frac{c_1}{c_2} \right)^2 \frac{c_5}{c_7} g(r(B)) \\ &= c_6 V^g(B). \end{aligned}$$

Property **(P4)** is satisfied because of (32). Finally, property **(P5)**, that $l_L \geq 2$ for any ball L in $K(n, B, l+1)$, follows from (23).

Therefore properties **(P1)**–**(P5)** are satisfied up to the local sub-level $K(n, B, l+1)$. This establishes the existence of the local level $K(n, B) = K_{l_B}(n, B)$ for each $B \in K(n-1)$. This then establishes the existence of the n th level $K(n)$ (and also $\mathbb{K}(n)$).

4.4 The measure μ on \mathbb{K}_η

In what follows we adopt the notation:

$$h := \frac{f}{g^\kappa}.$$

We now define our measure on \mathbb{K}_η which we will eventually see satisfies (16). For each level we distribute mass according to the following rules.

When $n = 1$ we have that $L = B_0 := \mathbb{K}(1)$ and let $\mu(L) := 1$.

For balls in $K(n)$, with $n > 2$, we distribute mass inductively. Therefore, let $n \geq 2$ and suppose $\mu(B)$ is defined for each $B \in K(n-1)$. Let L be a ball in $K(n)$. Since the balls in $K(n-1)$ are disjoint there is a unique ball $B \in K(n-1)$ satisfying $L \subset B$. By (17), (19) and (20), we know that

$$K(n, B) := \bigcup_{(A;j) \in \tilde{K}_{I_B}(n, B)} \mathcal{C}(A; j).$$

Therefore, L is contained in one of the collections $\mathcal{C}(A'; j')$ appearing in the above union. We define the mass on L to be

$$\mu(L) := \frac{1}{\#\mathcal{C}(A'; j')} \times \frac{h(\Upsilon_{j'})^{\frac{1}{1-\kappa}}}{\sum_{(A;j) \in \tilde{K}_{I_B}(n, B)} h(\Upsilon_j)^{\frac{1}{1-\kappa}}} \times \mu(B).$$

This quantity is well-defined in light of the preceding comment.

Proceeding inductively we see that μ is defined for each ball appearing in the construction of \mathbb{K}_η . We can extend μ uniquely in a standard way to all Borel subsets of X to give a probability measure μ supported on \mathbb{K}_η (see, for example, [9, Proposition 1.7] for further details). Given a Borel subset of X , say F , we let

$$\mu(F) := \mu(F \cap \mathbb{K}_\eta) = \inf \sum_{L \in \mathcal{C}(F)} \mu(L),$$

where the infimum is taken over all covers $\mathcal{C}(F)$ of $F \cap \mathbb{K}_\eta$ by balls $L \in \bigcup_{n \in \mathbb{N}} K(n)$.

Let us conclude this section by observing that for any $L \in K(n)$ we have

$$\begin{aligned} \mu(L) &\leq \frac{1}{d_1 \left(\frac{f(\Upsilon_{j'})}{g(\Upsilon_{j'})} \right)^{\frac{\kappa}{1-\kappa}}} \times \frac{h(\Upsilon_{j'})^{\frac{1}{1-\kappa}}}{\sum_{(A;j) \in \tilde{K}_{I_B}(n, B)} h(\Upsilon_j)^{\frac{1}{1-\kappa}}} \times \mu(B) \\ &= \frac{f(\Upsilon_{j'})}{d_1 \sum_{(A;j) \in \tilde{K}_{I_B}(n, B)} h(\Upsilon_j)^{\frac{1}{1-\kappa}}} \times \mu(B). \end{aligned} \tag{33}$$

This follows from (8) and the definition of h .

4.5 The measure of a ball in the Cantor set construction

Our ultimate goal is to prove that (16) is satisfied for any ball D of sufficiently small radius. Moving towards that goal, we first prove that

$$\mu(L) \ll \frac{V^f(L)}{\eta} \tag{34}$$

for any ball $L \in K(n)$ for $n \geq 2$. We start with $n = 2$ and then tackle higher levels of the Cantor set by induction. Let us fix a ball $L \in K(2) = K(2, B_0)$. Now, let $(A'; j') \in \tilde{K}_{l_{B_0}}(2, B_0)$ be such that $L \in \mathcal{C}(A'; j')$. Using the upper bound given by (33), the definition of μ , and the fact that $\mu(B_0) = 1$, we obtain

$$\mu(L) \leq \frac{f(\Upsilon_{j'})}{d_1 \sum_{(A;j) \in \tilde{K}_{l_{B_0}}(2, B_0)} h(\Upsilon_j)^{\frac{1}{1-\kappa}}} . \quad (35)$$

Using properties **(P3)** and **(P5)** of the Cantor set construction we obtain

$$\begin{aligned} \sum_{(A;j) \in \tilde{K}_{l_{B_0}}(2, B_0)} h(\Upsilon_j)^{\frac{1}{1-\kappa}} &= \sum_{(A;j) \in \tilde{K}_{l_{B_0}}(2, B_0)} V^g(A) \\ &= \sum_{i=1}^{l_{B_0}} \sum_{(A;j) \in \tilde{K}(2, B_0, i)} V^g(A) \\ &\stackrel{\text{(P3)}}{\geq} \sum_{i=1}^{l_{B_0}} c_6 V^g(B_0) \\ &= l_{B_0} c_6 V^g(B_0) \\ &\stackrel{(2)}{\geq} l_{B_0} \frac{c_6}{c_2} \mathcal{H}^g(B_0) \\ &\stackrel{\text{(P5)}}{\geq} \frac{c_2 \eta}{c_6 \mathcal{H}^g(B_0)} \frac{c_6}{c_2} \mathcal{H}^g(B_0) = \eta. \end{aligned}$$

Combining this estimate with (35), and observing that $f(\Upsilon_{j'}) = V^f(L)$, we obtain (34) as required.

We now consider $n > 2$. Assume that (34) holds for all balls in $K(n-1)$. Let L be an arbitrary ball in $K(n)$ and let $B \in K(n-1)$ be the unique ball such that $L \in K(n, B)$. Moreover, suppose $(A'; j') \in \tilde{K}_{l_B}(n, B)$ is the unique $(A'; j')$ such that $L \in \mathcal{C}(A'; j')$. By (33) and our induction hypothesis we have

$$\mu(L) \ll \frac{f(\Upsilon_{j'})}{d_1 \sum_{(A;j) \in \tilde{K}_{l_B}(n, B)} h(\Upsilon_j)^{\frac{1}{1-\kappa}}} \times \frac{V^f(B)}{\eta}. \quad (36)$$

Bounding the denominator of (36) we have

$$\begin{aligned}
\sum_{(A;j) \in \tilde{K}_{l_B}(n,B)} h(\Upsilon_j)^{\frac{1}{1-\kappa}} &= \sum_{i=1}^{l_B} \sum_{(A;j) \in \tilde{K}(n,B,i)} V^g(A) \\
&\stackrel{\text{(P3)}}{\geq} \sum_{i=1}^{l_B} c_6 V^g(B) \\
&= l_B c_6 V^g(B) \\
&\stackrel{\text{(P5)}}{\geq} \frac{V^f(B)}{c_6 V^g(B)} c_6 V^g(B) \\
&= V^f(B). \tag{37}
\end{aligned}$$

Combining (36) and (37) we see that (34) holds for L . By induction (34) holds for all $L \in K(n)$ for $n \geq 2$.

4.6 The measure of an arbitrary ball

Let $r_0 := \min\{r(B) : B \in K(2)\}$ and take an arbitrary ball D such that $r(D) < r_0$. To conclude our proof of Theorem 1 it suffices to prove (16) for D , that is we wish to show that

$$\mu(D) \ll \frac{V^f(D)}{\eta},$$

where the implied constant is independent of D and η . To prove this bound we will make use of the following lemma from [5]. This statement was originally (implicitly) proved in the setting of Euclidean space equipped with the usual metric. With virtually no change required to the proof, the same statement holds in an arbitrary metric space.

Lemma 7. *Let $A := B(x_A, r_A)$ and $M := B(x_M, r_M)$ be arbitrary balls in a metric space (X, d) such that $A \cap M \neq \emptyset$ and $A \setminus (cM) \neq \emptyset$ for some $c \geq 3$. Then $r_M \leq r_A$ and $cM \subset 5A$.*

Recall that our measure μ is supported on \mathbb{K}_η and we proved in the previous section that it satisfies the above inequality whenever D is a ball in our Cantor set construction. Consequently, without loss of generality, we may assume that D satisfies the following two properties:

- $D \cap \mathbb{K}_\eta \neq \emptyset$;
- for every n large enough D intersects at least two balls in $K(n)$.

If $D \cap \mathbb{K}_\eta = \emptyset$ then $\mu(D) = 0$ since μ is supported on \mathbb{K}_η . If the second assumption were false then D would intersect exactly one ball, say L_{n_i} , at level n_i

for infinitely many $i \in \mathbb{N}$. Then, by (34), we would have $\mu(D) \leq \mu(L_{n_i}) \rightarrow 0$ as $i \rightarrow \infty$. So, if either of the above two assumptions fail we have $\mu(D) = 0$ and (16) holds trivially.

By these two assumptions there exists a well-defined maximal integer n such that

$$D \text{ intersects at least 2 balls from } K(n) \quad (38)$$

and

$$D \text{ intersects only one ball } B \text{ from } K(n-1).$$

Since $r_0 = \min\{r(B) : B \in K(2)\}$ it follows that $n > 2$. Suppose $B \in K(n-1)$ is the unique ball which has non-empty intersection with D , then we may also assume that $r(D) < r(B)$. To see why, suppose otherwise that $r(B) \leq r(D)$. Since $D \cap \mathbb{K}_\eta \subset B$ and f is increasing, it would follow from (34) that

$$\mu(D) \leq \mu(B) \ll \frac{V^f(B)}{\eta} = \frac{f(r(B))}{\eta} \leq \frac{f(r(D))}{\eta} = \frac{V^f(D)}{\eta},$$

and (16) would be satisfied.

Note that, since $K(n, B)$ forms a cover of $D \cap \mathbb{K}_\eta$, we have

$$\begin{aligned} \mu(D) &\leq \sum_{i=1}^{l_B} \sum_{L \in K(n, B, i) : L \cap D \neq \emptyset} \mu(L) \\ &= \sum_{i=1}^{l_B} \sum_{(A; j) \in \tilde{K}(n, B, i)} \sum_{\substack{L \in \mathcal{C}(A; j) \\ L \cap D \neq \emptyset}} \mu(L) \\ &\stackrel{(34)}{\ll} \sum_{i=1}^{l_B} \sum_{(A; j) \in \tilde{K}(n, B, i)} \sum_{\substack{L \in \mathcal{C}(A; j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta}. \end{aligned} \quad (39)$$

The remainder of the proof of Theorem 1 will be concerned with showing that (39) is suitably bounded. In order to perform this task, it is useful to partition sub-levels into the following cases:

Case 1 : Sub-levels $K(n, B, i)$ for which

$$\#\{L \in K(n, B, i) : L \cap D \neq \emptyset\} = 1.$$

Case 2 : Sub-levels $K(n, B, i)$ for which

$$\#\{L \in K(n, B, i) : L \cap D \neq \emptyset\} \geq 2 \quad \text{and}$$

$$\#\{(A; j) \in \widetilde{K}(n, B, i) \text{ with } D \cap L \neq \emptyset \text{ for some } L \in \mathcal{C}(A; j)\} \geq 2.$$

Case 3 : Sub-levels $K(n, B, i)$ for which

$$\#\{L \in K(n, B, i) : L \cap D \neq \emptyset\} \geq 2 \quad \text{and}$$

$$\#\{(A; j) \in \widetilde{K}(n, B, i) \text{ with } D \cap L \neq \emptyset \text{ for some } L \in \mathcal{C}(A; j)\} = 1.$$

Technically we should also consider those sub-levels $K(n, B, i)$ for which $\#\{L \in K(n, B, i) : L \cap D \neq \emptyset\} = 0$. However, these sub-levels make no contribution to the sum on the right-hand side of (39) and can therefore be omitted.

Dealing with Case 1. Let $K(n, B, i^*)$ be the first sub-level whose intersection with D is described by Case 1. There is a unique ball L^* in $K(n, B, i^*)$ satisfying $L^* \cap D \neq \emptyset$. We know by (38) that there exists another ball $M \in K(n, B)$ such that $M \cap D \neq \emptyset$. Moreover, we also know that $3L^* \cap 3M = \emptyset$ by property **(P1)**. Therefore $D \setminus 3L^* \neq \emptyset$ and so, applying Lemma 7, we have $r(L^*) \leq r(D)$. Consequently, since f is a dimension function and hence increasing,

$$V^f(L^*) \leq V^f(D). \quad (40)$$

Using property **(P4)** we know that for $i \in \{i^* + 1, \dots, l_B\}$ and $L \in K(n, B, i)$, we have

$$V^f(L) = f(r(L)) \leq 2^{-(i-i^*)} f(r(L^*)) = 2^{-(i-i^*)} V^f(L^*).$$

Combining this inequality with (40), we see that the contribution to the right-hand side of (39) from Case 1 is:

$$\sum_{i \in \text{Case 1}} \sum_{\substack{L \in K(n, B, i) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} \leq \sum_{i \geq i^*} 2^{-(i-i^*)} \frac{V^f(L^*)}{\eta} \leq 2 \frac{V^f(L^*)}{\eta} \leq 2 \frac{V^f(D)}{\eta}. \quad (41)$$

Dealing with Case 2. Let $K(n, B, i)$ be a sub-level whose intersection with D is described by Case 2. Thus, there exist distinct balls $(A; j)$ and $(A'; j')$ in $\widetilde{K}(n, B, i)$, and corresponding balls $L \in \mathcal{C}(A; j)$ and $L' \in \mathcal{C}(A'; j')$ satisfying $L \cap D \neq \emptyset$ and $L' \cap D \neq \emptyset$. Since $L \subset A$ and $L' \subset A'$ we have $A \cap D \neq \emptyset$ and $A' \cap D \neq \emptyset$. By property **(P2)** of our construction we know that the balls $3A$ and $3A'$ are disjoint and contained in B . Therefore $D \setminus 3A \neq \emptyset$ and, applying Lemma 7, we see that $r(A) \leq r(D)$ and $A \subset 3A \subset 5D$. By the same reasoning we also have $A' \subset 3A' \subset 5D$. Hence, on using (8) we get that the

contribution to the right-hand side of (39) from Case 2 is estimated as follows

$$\begin{aligned}
\sum_{i \in \text{Case 2}} \sum_{(A;j) \in \tilde{K}(n,B,i)} \sum_{\substack{L \in \mathcal{C}(A;j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} &\leq \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \#\mathcal{C}(A;j) \frac{f(\Upsilon_j)}{\eta} \\
&\stackrel{(8)}{\ll} \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \left(\frac{f(\Upsilon_j)}{g(\Upsilon_j)} \right)^{\frac{\kappa}{1-\kappa}} \frac{f(\Upsilon_j)}{\eta} \\
&= \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \frac{f(\Upsilon_j)^{\frac{\kappa}{1-\kappa}+1}}{g(\Upsilon_j)^{\frac{\kappa}{1-\kappa}}} \times \frac{1}{\eta} \\
&= \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \frac{f(\Upsilon_j)^{\frac{1}{1-\kappa}}}{g(\Upsilon_j)^{\frac{\kappa}{1-\kappa}}} \times \frac{1}{\eta} \\
&= \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \frac{V^g(A)}{\eta}.
\end{aligned}$$

It follows upon combining this estimate with (1), (2), and the disjointness of balls in $\tilde{K}(n, B, i)$ guaranteed by property **(P2)**, that

$$\begin{aligned}
\sum_{i \in \text{Case 2}} \sum_{(A;j) \in \tilde{K}(n,B,i)} \sum_{\substack{L \in \mathcal{C}(A;j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} &\stackrel{(2)}{\ll} \frac{1}{\eta} \sum_{i \in \text{Case 2}} \sum_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} \mathcal{H}^g(A) \\
&\stackrel{(\mathbf{P2})}{=} \frac{1}{\eta} \sum_{i \in \text{Case 2}} \mathcal{H}^g \left(\bigcup_{\substack{(A;j) \in \tilde{K}(n,B,i) \\ AC5D}} A \right) \\
&\leq \frac{1}{\eta} \sum_{i \in \text{Case 2}} \mathcal{H}^g(5D) \\
&\stackrel{(2)}{\ll} \frac{1}{\eta} \sum_{i \in \text{Case 2}} V^g(5D) \\
&\stackrel{(1)}{\ll} \frac{1}{\eta} \sum_{i \in \text{Case 2}} V^g(D) \\
&\leq \frac{1}{\eta} l_B V^g(D).
\end{aligned}$$

Finally, it follows from the above estimate together with property **(P5)** that

$$\begin{aligned}
\sum_{i \in \text{Case 2}} \sum_{(A;j) \in \tilde{K}(n,B,i)} \sum_{\substack{L \in \mathcal{C}(A;j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} &\stackrel{\text{(P5)}}{\ll} \frac{1}{\eta} \left(\frac{V^f(B)}{V^g(B)} \right) V^g(D) \\
&\ll \frac{1}{\eta} \frac{V^f(D)}{V^k(D)} V^k(D) \\
&= \frac{V^f(D)}{\eta}. \tag{42}
\end{aligned}$$

To deduce the penultimate inequality we used the facts that f/g is decreasing and $r(D) < r(B)$.

Dealing with Case 3. For each sub-level i whose intersection with D is described by Case 3 there exists a unique $(A_i; j_i) \in \tilde{K}(n, B, i)$ such that D has non-empty intersection with balls in $\mathcal{C}(A_i; j_i)$. Let $K(n, B, i^{**})$ denote the first sub-level described by Case 3. There exists a ball L^{**} in $K(n, B, i^{**})$ such that $L^{**} \cap D \neq \emptyset$. By the assumption in (38) there must exist another ball $M \in K(n, B)$ such that $M \cap D \neq \emptyset$. It follows from property **(P1)** that $3L^{**}$ and $3M$ are disjoint and so $D \setminus 3L^{**} \neq \emptyset$. Applying Lemma 7, we have that $r(L^{**}) \leq r(D)$. As h is a dimension function it follows that

$$h(r(L^{**})) \leq h(r(D)). \tag{43}$$

By property **(P4)** we know that for any $i \in \{i^{**} + 1, \dots, l_B\}$ and $L \in K(n, B, i)$ we have

$$h(r(L)) \leq 2^{-(i-i^{**})} h(r(L^{**})). \tag{44}$$

Recall that, by Lemma 6, each $L \in \mathcal{C}(A_i; j_i)$ is centred on F_{j_i} . Combining this fact with the LSP, the relations given by (2), and the fact that the elements of $\mathcal{C}(A_i; j_i)$ are disjoint, by straightforward measure theoretic considerations we have the following estimate

$$\#\{L \in \mathcal{C}(A_i; j_i) : L \cap D \neq \emptyset\} \ll \frac{g(\Upsilon_{j_i})^{1-\kappa} g(r(D))^\kappa}{g(\Upsilon_{j_i})}. \tag{45}$$

Therefore, the contribution to the right-hand side of (39) from Case 3 can be

bounded above as follows:

$$\begin{aligned}
\sum_{i \in \text{Case 3}} \sum_{(A;j) \in \tilde{K}(n,B,i)} \sum_{\substack{L \in \mathcal{C}(A;j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} &\leq \sum_{i \in \text{Case 3}} \sum_{\substack{L \in \mathcal{C}(A_i;j_i) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} \\
&= \sum_{i \in \text{Case 3}} \sum_{\substack{L \in \mathcal{C}(A_i;j_i) \\ L \cap D \neq \emptyset}} \frac{f(\Upsilon_{j_i})}{\eta} \\
&\stackrel{(45)}{\ll} \sum_{i \in \text{Case 3}} \frac{g(\Upsilon_{j_i})^{1-\kappa} g(r(D))^\kappa}{g(\Upsilon_{j_i})} \times \frac{f(\Upsilon_{j_i})}{\eta} \\
&= \frac{g(r(D))^\kappa}{\eta} \sum_{i \in \text{Case 3}} \frac{f(\Upsilon_{j_i})}{g(\Upsilon_{j_i})^\kappa} \\
&= \frac{g(r(D))^\kappa}{\eta} \sum_{i \in \text{Case 3}} h(\Upsilon_{j_i}) \\
&\stackrel{(44)}{\ll} \frac{g(r(D))^\kappa}{\eta} \sum_{i \geq i^{**}} \frac{h(\Upsilon_{j_{i^{**}}})}{2^{i-i^{**}}} \\
&\leq 2 \frac{g(r(D))^\kappa}{\eta} h(\Upsilon_{j_{i^{**}}}).
\end{aligned}$$

Recalling (43) and noting that $\Upsilon_{j_i} = r(L^{**})$, we observe that

$$\begin{aligned}
\sum_{i \in \text{Case 3}} \sum_{(A;j) \in \tilde{K}(n,B,i)} \sum_{\substack{L \in \mathcal{C}(A;j) \\ L \cap D \neq \emptyset}} \frac{V^f(L)}{\eta} &\ll 2 \frac{g(r(D))^\kappa}{\eta} h(r(D)) \\
&= 2 \frac{f(r(D))}{\eta} \ll \frac{V^f(D)}{\eta}. \quad (46)
\end{aligned}$$

Combining estimates (41), (42) and (46) with (39) gives $\mu(D) \ll \frac{V^f(D)}{\eta}$, thus proving (16) as desired. This completes the proof of Theorem 1.

5 Examples of sets satisfying the local scaling property

In this section we briefly detail two collections of sets that satisfy the LSP for appropriate choices of parameters.

5.1 Smooth compact manifolds

Let $M \subset \mathbb{R}^n$ be a smooth compact manifold of dimension l . Let us start by remarking that when M is a smooth compact manifold, the tangent space

map sending $x \rightarrow T_x M$ is a continuous map from M into the Grassmanian of l -dimensional subspaces of \mathbb{R}^n . Fixing $x \in M$, and applying a rotation if necessary, we can identify $T_x M$ with $\mathbb{R}^l \times \{0^{n-l}\}$. Since the tangent space map is continuous we know that for any y sufficiently close to x its tangent space $T_y M$ is approximately $\mathbb{R}^l \times \{0^{n-l}\}$. As such, we can assert that there exists $R_x > 0$ such that the following two properties hold:

- For any $y \in B(x, R_x) \cap M$ and $r > 0$ such that $B(y, r) \subset B(x, R_x)$, we have

$$\mathcal{H}^l(\pi_l(B(y, r) \cap M)) \asymp r^l. \quad (47)$$

Here π_l is the projection map from \mathbb{R}^n to \mathbb{R}^l sending (x_1, \dots, x_n) to (x_1, \dots, x_l) .

- For any $y \in B(x, R_x) \cap M$ and $\delta < R_x$ we have

$$\mathcal{H}^{n-l}(\{z \in \Delta(M, \delta) : \pi(z) = \pi(y)\}) \asymp \delta^{n-l}. \quad (48)$$

The implied constants in (47) and (48) depend only upon M and x . Via an application of (47), (48), and Fubini's Theorem, it can be shown that for any $y \in B(x, R_x/2)$, $r \leq R_x/2$, and $\delta \leq r$ we have

$$\mathcal{H}^n(B(y, r) \cap \Delta(M, \delta)) \asymp r^l \delta^{n-l}. \quad (49)$$

Again, the implied constants in (49) depend only upon M and x . Since $\{B(x, R_x/2)\}_{x \in M}$ covers M , it follows by a compactness argument that there exists $R > 0$ such that for any $y \in M$, $r \leq R$ and $\delta \leq r$ we have

$$\mathcal{H}^n(B(y, r) \cap \Delta(M, \delta)) \asymp r^l \delta^{n-l}.$$

Thus, we conclude that M satisfies the LSP with respect to $\kappa = \frac{l}{n}$.

5.2 Self-similar sets satisfying the open set condition

Let $\Phi = \{\phi_1, \dots, \phi_l\}$ be a collection of contracting similarities acting on \mathbb{R}^n ; that is, Φ is a collection of maps such that

$$|\phi_i(x) - \phi_i(y)| = r_i |x - y| \quad \text{for all } x, y \in \mathbb{R}^n$$

and $0 < r_i < 1$ for each $1 \leq i \leq l$. It is well-known (see, for example, [9]) that there exists a unique non-empty compact set $K \subset \mathbb{R}^n$ such that

$$K = \bigcup_{i=1}^l \phi_i(K).$$

We say that Φ satisfies the *open set condition* if there exists an open set $O \subset \mathbb{R}^n$ such that $\phi_i(O) \subset O$ for all i , and $\phi_i(O) \cap \phi_j(O) = \emptyset$ for all $i \neq j$.

In [11] it is shown that if Φ satisfies the open set condition then there exist constants $c_3, c_4 > 0$ such that for all δ sufficiently small

$$c_3\delta^{n-d} \leq \mathcal{H}^n(\Delta(K, \delta)) \leq c_4\delta^{n-d}, \quad (50)$$

where d is the box counting dimension of K . In [11], (50) was proved for the n -dimensional Lebesgue measure. The statement given above follows since \mathcal{H}^n is equal to a scalar multiple of the n -dimensional Lebesgue measure.

We now prove that the local scaling property holds for K with $\kappa = \frac{d}{n}$. To this end, fix $x \in K$ and some small number $r > 0$. Then there exists a sequence $(a_i)_{i \in \mathbb{N}} \in \{1, \dots, l\}^{\mathbb{N}}$ such that

$$\bigcap_{m=1}^{\infty} (\phi_{a_1} \circ \dots \circ \phi_{a_m})(K) = x.$$

Next, note that there exists $N \in \mathbb{N}$ such that

$$(r_{a_1} \cdots r_{a_{N-1}}) \cdot \text{Diam}K \geq r/2$$

and

$$(r_{a_1} \cdots r_{a_N}) \cdot \text{Diam}K < r/2.$$

For this value of N and $\delta < r/2$, we have

$$\begin{aligned} \mathcal{H}^n(B(x, r) \cap \Delta(K, \delta)) &\geq \mathcal{H}^n(B(x, r) \cap \Delta((\phi_{a_1} \circ \dots \circ \phi_{a_N})(K), \delta)) \\ &\geq \mathcal{H}^n(\Delta((\phi_{a_1} \circ \dots \circ \phi_{a_N})(K), \delta)) \\ &= \mathcal{H}^n((\phi_{a_1} \circ \dots \circ \phi_{a_N})(\Delta(K, \delta \cdot (r_{a_1} \cdots r_{a_N})^{-1}))) \\ &= (r_{a_1} \cdots r_{a_N})^n \mathcal{H}^n(\Delta(K, \delta \cdot (r_{a_1} \cdots r_{a_N})^{-1})) \\ &\stackrel{(50)}{\geq} c_3 \cdot (r_{a_1} \cdots r_{a_N})^n \cdot \left(\frac{\delta}{r_{a_1} \cdots r_{a_N}} \right)^{n-d} \\ &= c_3 \cdot (r_{a_1} \cdots r_{a_N})^d \cdot \delta^{n-d} \\ &\asymp c_3 r^d \delta^{n-d}. \end{aligned} \quad (51)$$

The last line follows since $r_{a_1} \cdots r_{a_N} \asymp r$ by our choice of N . Thus, we have proved that the lower bound in the LSP holds in this case. It remains to prove the upper bound.

Given $r > 0$ let

$$I_r := \{(a_1, \dots, a_k) \in \cup_{j=0}^{\infty} \{1, \dots, l\}^j : r_{a_1} \cdots r_{a_k} \leq r < r_{a_1} \cdots r_{a_{k-1}}\}.$$

Importantly $\{(\phi_{a_1} \circ \dots \circ \phi_{a_k})(K)\}_{(a_1, \dots, a_k) \in I_r}$ forms a cover of K . In [9] it is shown that when the open set condition holds, for any $x \in K$ and $r > 0$ we have

$$\#\{(a_1, \dots, a_k) \in I_r : B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset\} \leq C \quad (52)$$

for some $C > 0$ independent of r .

Fixing $x \in K$ and $r > 0$ and using (50) and (52), for $\delta < \frac{r}{2}$ we obtain

$$\begin{aligned}
\mathcal{H}^n(B(x, r) \cap \Delta(K, \delta)) &\leq \sum_{\substack{(a_1, \dots, a_k) \in I_r \\ B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset}} \mathcal{H}^n(\Delta((\phi_{a_1} \circ \dots \circ \phi_{a_k})(K), \delta)) \\
&= \sum_{\substack{(a_1, \dots, a_k) \in I_r \\ B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset}} \mathcal{H}^n((\phi_{a_1} \circ \dots \circ \phi_{a_k})(\Delta(K, \delta \cdot (r_{a_1} \cdots r_{a_k})^{-1}))) \\
&= \sum_{\substack{(a_1, \dots, a_k) \in I_r \\ B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset}} (r_{a_1} \cdots r_{a_k})^n \mathcal{H}^n(\Delta(K, \delta \cdot (r_{a_1} \cdots r_{a_k})^{-1})) \\
&\stackrel{(50)}{\leq} c_4 \sum_{\substack{(a_1, \dots, a_k) \in I_r \\ B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset}} (r_{a_1} \cdots r_{a_k})^n \left(\frac{\delta}{r_{a_1} \cdots r_{a_k}} \right)^{n-d} \\
&\leq c_4 \sum_{\substack{(a_1, \dots, a_k) \in I_r \\ B(x, r) \cap (\phi_{a_1} \circ \dots \circ \phi_{a_k})(K) \neq \emptyset}} r^d \delta^{n-d} \\
&\stackrel{(52)}{\leq} c_4 C r^d \delta^{n-d}. \tag{53}
\end{aligned}$$

Combining (51) and (53), we see that K satisfies (4) for $\delta < r/2$. Equation (4) trivially holds for $\delta \in [r/2, r)$, since in that case $B(x, r) \cap \Delta(K, \delta)$ contains a ball of radius $r/2$ and is contained in a ball of radius r . Therefore (4) holds for all $\delta < r$ and K satisfies the LSP with respect to $\kappa = \frac{d}{n}$ as claimed.

6 An application of Theorem 1: Random *limsup* sets

In this section we will impose the additional assumptions that (X, d) is a compact metric space and $g(r) := r^n$ for some $n > 0$. By rescaling if necessary, we may assume without loss of generality that $\mathcal{H}^n(X) = 1$. Let $(F_j)_{j \in \mathbb{N}}$ be a sequence of sets satisfying the LSP with respect to some $0 \leq \kappa < 1$. Assume that we are given a set of isometries Φ that are chosen randomly according to some law \mathbb{P} . Given a sequence of non-negative real numbers $\Upsilon := (\Upsilon_j)_{j \in \mathbb{N}}$ and a randomly chosen sequence $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$, we define the corresponding random lim sup set as follows:

$$\Lambda((\phi_j), \Upsilon) := \{x \in \Delta(\phi_j(F_j), \Upsilon_j) \text{ for infinitely many } j \in \mathbb{N}\}.$$

We are interested in determining the \mathbb{P} -almost sure Hausdorff dimension and Hausdorff measure of $\Lambda((\phi_j), \Upsilon)$. When $(F_j)_{j \in \mathbb{N}}$ is a sequence of points, then the \mathbb{P} -almost sure Hausdorff dimension and Hausdorff measure of $\Lambda((\phi_j), \Upsilon)$ is well understood. See [12] and references therein for more on this classical problem. When $(F_j)_{j \in \mathbb{N}}$ is a more exotic sequence of sets, the problem of determining the \mathbb{P} -almost sure metric properties of $\Lambda((\phi_j), \Upsilon)$ is more difficult. That being said, in the Euclidean setting a comprehensive solution to the \mathbb{P} -almost sure metric properties of $\Lambda((\phi_j), \Upsilon)$ is given in [10]. See also [13] and [17]. Our application

below holds in the more general metric space setting. In what follows we assume that \mathbb{P} satisfies the following properties:

- For any $j \in \mathbb{N}$ and $\delta > 0$ sufficiently small:

$$\mathbb{P}(x \in \Delta(\phi_j(F_j), \delta)) = \mathcal{H}^n(\Delta(F_j, \delta)). \quad (54)$$

- For any sequence (δ_j) of sufficiently small numbers, we have that the sequence of events $(E_j)_{j=1}^\infty = (\{x \in \Delta(\phi_j(F_j), \delta_j)\})_{j=1}^\infty$ are independent, i.e. for any finite set $S \subset \mathbb{N}$ we have

$$\mathbb{P}\left(\bigcap_{j \in S} E_j\right) = \prod_{j \in S} \mathbb{P}(E_j).$$

The main result of this section is Theorem 2 below. For convenience it is stated for $(\Upsilon_j)_{j \in \mathbb{N}}$ of the form $(j^{-\tau})_{j \in \mathbb{N}}$, although it holds in greater generality. To prove this theorem we will make use of the following well known result from probability known as the second Borel–Cantelli lemma, see [8].

Lemma 8 (Second Borel–Cantelli Lemma). *Let (X, \mathcal{A}, μ) be a probability space. Let (E_j) be an independent sequence of events such that $\sum_{j=1}^\infty \mu(E_j) = \infty$, then $\mu(\limsup E_j) = 1$.*

Theorem 2. *Let (X, d) , Φ , and \mathbb{P} be as above. Suppose $(F_j)_{j \in \mathbb{N}}$ satisfies the LSP with respect to some $0 \leq \kappa < 1$ and let $\tau > \frac{1}{n-\kappa n}$. Then, for \mathbb{P} -almost every $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$, we have*

$$\dim_{\text{H}}(\Lambda((\phi_j), (j^{-\tau}))) = \kappa n + \frac{1}{\tau} \quad \text{and} \quad \mathcal{H}^{\kappa n + 1/\tau}(\Lambda((\phi_j), (j^{-\tau}))) = \infty.$$

Proof. It follows from the LSP that for any $j \in \mathbb{N}$ and $\delta > 0$ sufficiently small we have

$$\mathcal{H}^n(\Delta(F_j, \delta)) \asymp \delta^{n-\kappa n}. \quad (55)$$

Using the notation of Section 3, we see that when $f(r) = r^s$ we have $\tilde{\Upsilon}_j = j^{\frac{\tau(\kappa n - s)}{n - \kappa n}}$. When $s = \kappa n + 1/\tau$ we have

$$\sum_{j=1}^\infty \mathbb{P}(x \in \Delta(\phi_j(F_j), \tilde{\Upsilon}_j)) \stackrel{(54)}{=} \sum_{j=1}^\infty \mathcal{H}^n(\Delta(F_j, \tilde{\Upsilon}_j)) \stackrel{(55)}{\asymp} \sum_{j=1}^\infty j^{-1} = \infty.$$

Applying Lemma 8, we may assert that for a fixed $x \in X$ we have

$$\mathbb{P}(x \in \Lambda((\phi_j), \tilde{\Upsilon})) = 1.$$

Applying Fubini's Theorem, we obtain that for \mathbb{P} -almost every $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$ we have

$$\mathcal{H}^n(\Lambda((\phi_j), \tilde{\Upsilon})) = 1, \quad (56)$$

i.e. our random limsup set has full measure. Equation (56) tells us that for almost every $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$ the assumptions of Theorem 1 are satisfied. Therefore, we may apply Theorem 1 and conclude that for almost every $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$ we have

$$\mathcal{H}^{\kappa n + 1/\tau}(\Lambda((\phi_j), (j^{-\tau}))) = \infty,$$

since we assumed that $\mathcal{H}^n(X) = 1$ and $\kappa n + 1/\tau < n$.

To complete our proof, it suffices to show that $\mathcal{H}^s(\Lambda((\phi_j), (j^{-\tau}))) = 0$ for any $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$ and $s > \kappa n + 1/\tau$. Therefore fix $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$. Applying the $5r$ -covering lemma (Lemma 3) we may construct for each $j \in \mathbb{N}$ a finite disjoint collection of balls $\{B(x_{l,j}, j^{-\tau})\}_{l \in Y_j}$ such that $\{B(x_{l,j}, 5j^{-\tau})\}_{l \in Y_j}$ covers $\Delta(\phi(F_j), j^{-\tau})$. Notice that the collection $\{B(x_{l,j}, 5j^{-\tau})\}_{l \in Y_j, j \geq N}$ covers $\Lambda((\phi_j), (j^{-\tau}))$ for any $N \in \mathbb{N}$. Furthermore, combining (2) and (55) we see that

$$\#Y_j \ll \frac{\mathcal{H}^n(\Delta(\phi_j(F_j), j^{-\tau}))}{j^{-\tau n}} \underset{(55)}{\asymp} \frac{j^{-\tau(n-\kappa n)}}{j^{-\tau n}} = j^{\tau \kappa n}. \quad (57)$$

Taking $\rho > 0$, $\varepsilon > 0$ and $s > \kappa n + 1/\tau$, we may choose N sufficiently large such that

$$\mathcal{H}_\rho^s(\Lambda((\phi_j), (j^{-\tau}))) \leq \sum_{j=N}^{\infty} \sum_{l \in Y_j} (5j^{-\tau})^s \stackrel{(57)}{\ll} 5^s \sum_{j=N}^{\infty} j^{\tau \kappa n - \tau s} < \varepsilon.$$

Since ε and ρ were arbitrary we may deduce that $\mathcal{H}^s(\Lambda((\phi_j), (j^{-\tau}))) = 0$, as required. Therefore, for \mathbb{P} -almost every $(\phi_j)_{j \in \mathbb{N}} \in \Phi^{\mathbb{N}}$, we have

$$\dim_{\mathbb{H}}(\Lambda((\phi_j), (j^{-\tau}))) = \kappa n + \frac{1}{\tau}. \quad \square$$

Acknowledgements. This work began while both authors were in attendance at the program on *Fractal Geometry and Dynamics* at the Mittag-Leffler Institut in November 2017. We are grateful both to the organisers of the program and the staff at the Institut for making it such a wonderful environment for doing mathematics. DA is grateful to Victor Beresnevich and Sanju Velani, her PhD supervisors, for introducing her to the Mass Transference Principle and the topic of metric Diophantine approximation in general.

References

- [1] D. Allen, V. Beresnevich, *A Mass Transference Principle for systems of linear forms and its applications*, preprint, arXiv:1703.10015.
- [2] V. Beresnevich, V. Bernik, M. Dodson, S. Velani, *Classical metric Diophantine approximation revisited*, Analytic number theory, 38–61, Cambridge Univ. Press, Cambridge, 2009.
- [3] V. Beresnevich, D. Dickinson, S. Velani, *Measure theoretic laws for lim sup sets*, Mem. Amer. Math. Soc. 179 (2006), no. 846, x+91 pp.

- [4] V. Beresnevich, F. Ramírez, S. Velani, *Metric Diophantine Approximation: Aspects of Recent Work*, in D. Badziahin, A. Gorodnik, & N. Peyerimhoff (eds.) *Dynamics and Analytic Number Theory*, London Mathematical Society Lecture Note Series, vol. 437, Cambridge University Press, pp. 1–95.
- [5] V. Beresnevich, S. Velani, *A Mass Transference Principle and the Duffin–Schaeffer conjecture for Hausdorff measures*, *Ann. of Math. (2)* 164 (2006), no. 3, 971–992.
- [6] V. Beresnevich, S. Velani, *Schmidt’s theorem, Hausdorff measures, and slicing*, *Int. Math. Res. Not.* 2006, Art. ID 48794, 24 pp.
- [7] V. Beresnevich, S. Velani, *A note on zero-one laws in metrical Diophantine approximation*, *Acta Arith.* 133 (2008), no. 4, 363–374.
- [8] P. Billingsley, *Probability and measure*, Wiley Series in Probability and Statistics. John Wiley & Sons, Inc., Hoboken, NJ, 2012. xviii+624 pp.
- [9] K. Falconer, *Fractal geometry: Mathematical foundations and applications*, Second edition, John Wiley & Sons, 2003.
- [10] D. J. Feng, E. Järvenpää, M. Järvenpää, V. Suomala, *Dimensions of random covering sets in Riemann manifolds* To appear in *Ann. Probab.*
- [11] D. Gatzouras, *Lacunarity of self-similar and stochastically self-similar sets*. *Trans. Amer. Math. Soc.* 352 (2000), no. 5, 1953–1983.
- [12] M. Järvenpää, *Random Covering Sets, Hitting Probabilities and Variants of the Covering Problem*. In: Barral J., Seuret S. (eds) *Recent Developments in Fractals and Related Fields. FARF3 2015*. Trends in Mathematics. Birkhuser. (2017)
- [13] E. Järvenpää, M. Järvenpää, H. Koivusalo, B. Li, V. Suomala, Y. Xiao, *Hitting probabilities of random covering sets in tori and metric spaces* *Electron. J. Probab.* 22 (2017), Paper No. 1, 18 pp.
- [14] S. Kombrink, *A survey on Minkowski measurability of self-similar and self-conformal fractals in \mathbb{R}^d* , *Fractal geometry and dynamical systems in pure and applied mathematics. I. Fractals in pure mathematics*, 135–159, *Contemp. Math.*, 600, Amer. Math. Soc., Providence, RI, 2013.
- [15] J. Levesley, C. Salp, S. Velani, *On a problem of K. Mahler: Diophantine approximation and Cantor sets*, *Math. Ann.* 338 (2007), no. 1, 97–118.
- [16] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge Studies in Advanced Mathematics, 44., Cambridge University Press, Cambridge, 1995.
- [17] T. Persson, *A Note on Random Coverings of Tori*, *Bulletin of the London Mathematical Society*, 2015, 47 (1), 7–2.

- [18] S. Rigot, *Differentiation of measures in metric spaces*, preprint, (2018), arXiv:1802.02069.
- [19] B. Wang, J. Wu and J. Xu, *Mass transference principle for limsup sets generated by rectangles*, Math. Proc. Cambridge Philos. Soc. 158 (2015), no. 3, 419–437.

D. Allen, SCHOOL OF MATHEMATICS, UNIVERSITY OF MANCHESTER, OXFORD ROAD, MANCHESTER, M13 9PL, UK

E-mail address: `demi.allen@manchester.ac.uk`

S. Baker, MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UK

E-mail address: `simonbaker412@gmail.com`