

THE DIMENSION OF AN AMOEBA

JAN DRAISMA, JOHANNES RAU, AND CHI HO YUEN

ABSTRACT. Answering a question by Nisse and Sottile, we derive a formula for the dimension of the amoeba of an irreducible algebraic variety.

1. INTRODUCTION AND MAIN RESULT

Let $X \subseteq (\mathbb{C}^*)^n$ be an irreducible, closed algebraic subvariety. We define

$$\text{Log} : (\mathbb{C}^*)^n \rightarrow \mathbb{R}^n, \quad (z_1, \dots, z_n) \mapsto (\log |z_1|, \dots, \log |z_n|)$$

and $\mathcal{A}(X) := \text{Log}(X)$, the *amoeba* of X . The amoeba is the image of the semi-algebraic set (*algebraic amoeba*)

$$|X| := \{(|z_1|, \dots, |z_n|) \mid (z_1, \dots, z_n) \in X\} \subseteq \mathbb{R}_{>0}^n,$$

under a diffeomorphism and thus has an obvious notion of dimension, denoted $\dim_{\mathbb{R}} \mathcal{A}(X)$. Clearly, $\dim_{\mathbb{R}} \mathcal{A}(X) \leq 2 \dim_{\mathbb{C}} X$. In [NS18], Nisse and Sottile raise the question when this inequality is strict, as happens in the following two examples.

Example 1 (hypersurfaces). Suppose that $n > 2$ and that X is a hypersurface. Then $\dim_{\mathbb{R}} \mathcal{A}(X) \leq n < 2(n-1) = 2 \dim_{\mathbb{C}} X$. ♣

Example 2 (torus-invariant varieties). Suppose that X is stable under a subtorus $S \subseteq (\mathbb{C}^*)^n$ of dimension $k > 0$. Denote by Y the image of X in the algebraic torus $(\mathbb{C}^*)^n/S \cong (\mathbb{C}^*)^{n-k}$. The map $X \rightarrow Y$ has fibers of complex dimension k , and the corresponding map $\mathcal{A}(X) \rightarrow \mathcal{A}(Y)$ has fibers of real dimension k —namely, translates of $\mathcal{A}(S)$, which is a linear subspace of \mathbb{R}^n spanned by its intersection with \mathbb{Q}^n . Thus we have

$$\dim_{\mathbb{R}} \mathcal{A}(X) = k + \dim_{\mathbb{R}} \mathcal{A}(Y) \leq k + 2 \dim_{\mathbb{C}} Y = -k + 2 \dim_{\mathbb{C}} X < \dim_{\mathbb{C}} X. \quad \clubsuit$$

Our theorem says that these are two instances of the same phenomenon, and that this phenomenon is responsible for all drops in dimension.

Theorem 3. *Let $X \subseteq (\mathbb{C}^*)^n$ be an irreducible, closed algebraic subvariety. Then*

$$\dim_{\mathbb{R}} \mathcal{A}(X) = \min\{2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S \mid T \subseteq S \subseteq (\mathbb{C}^*)^n \text{ subtori and } S \cdot \overline{(T \cdot X)} = \overline{T \cdot X}\}.$$

An equivalent but more concise formula can then be given as

$$\dim_{\mathbb{R}} \mathcal{A}(X) = \min\{2 \dim_{\mathbb{C}} \overline{S \cdot X} - \dim_{\mathbb{C}} S \mid S \subseteq (\mathbb{C}^*)^n \text{ subtorus}\}.$$

In this theorem, $\overline{T \cdot X} = \overline{TX}$ (resp. $\overline{S \cdot X} = \overline{SX}$) is the Zariski closure of the set of all tz with $t \in T$ (resp. all rz with $r \in S$) and $z \in X$. Naturally, S and T may be taken zero-dimensional, in which case we recover the upper bound $2 \dim_{\mathbb{C}} X$.

Example 4 (hypersurfaces revisited). If X is a hypersurface, then most one-dimensional tori $T \subseteq (\mathbb{C}^*)^n$ will satisfy $\overline{T \cdot X} = (\mathbb{C}^*)^n$ (see Lemma 10), so we may take $S = (\mathbb{C}^*)^n$. The bound in the theorem is $2(n-1) + 2 - n = n$. ♣

To motivate the structure of this paper, we now prove the easy inequality \leq in our main theorem.

Proof of \leq in Theorem 3. Let $T \subseteq S \subseteq (\mathbb{C}^*)^n$ be subtori such that $Y := \overline{T \cdot X}$ is S -stable. Then

$$\dim_{\mathbb{R}} \mathcal{A}(X) \leq \dim_{\mathbb{R}} \mathcal{A}(Y) \leq 2 \dim_{\mathbb{C}} Y - \dim_{\mathbb{C}} S \leq 2(\dim_{\mathbb{C}} X + \dim_{\mathbb{C}} T) - \dim_{\mathbb{C}} S,$$

where the second equality follows from Example 2. \square

If we want equality to hold in the proof above, then we need that first, $\dim_{\mathbb{C}} Y = \dim_{\mathbb{C}} X + \dim_{\mathbb{C}} T$; second, the bound in Example 2 for the pair (Y, S) is tight; and third, $\dim_{\mathbb{R}} \mathcal{A}(X) = \dim_{\mathbb{R}} \mathcal{A}(Y) = \dim_{\mathbb{R}}(\mathcal{A}(X) + \mathcal{A}(T))$. Our proof of Theorem 3 consists of first finding a torus T with the latter property (see Section 2):

Proposition 5. *Let $X \subseteq (\mathbb{C}^*)^n$ be a closed, irreducible variety. Then the Zariski-closure \overline{X} in $(\mathbb{R}^*)^n$ of the algebraic amoeba is stable under a subtorus of the real algebraic torus $(\mathbb{R}^*)^n$ of dimension at least $2 \dim_{\mathbb{C}} X - \dim_{\mathbb{R}} \mathcal{A}(X)$.*

In particular, if the amoeba has dimension strictly less than $2 \dim_{\mathbb{C}} X$, then a positive-dimensional real torus acts on \overline{X} . Using this positive-dimensional torus, we prove Theorem 3 by induction in Section 3.

Theorem 3 implies [NS18, Conjecture 4.4], which proposes *near torus actions* (Definition 12 below) as the only cause of dimension drops for the amoeba.

Corollary 6. *For an irreducible, closed subvariety $X \subseteq (\mathbb{C}^*)^n$, we have*

$$\dim_{\mathbb{R}} \mathcal{A}(X) < \min\{n, 2 \dim_{\mathbb{C}} X\}$$

if and only if some nontrivial subtorus $S \subseteq (\mathbb{C}^)^n$ has a near action on X .*

We conclude this introduction with a relation to the tropical variety of X , also to be proved in Section 3.

Corollary 7. *For any irreducible, closed subvariety $X \subseteq (\mathbb{C}^*)^n$ the dimension $\dim_{\mathbb{R}} \mathcal{A}(X)$ is determined by the tropical variety $\text{Trop}(X) \subseteq \mathbb{R}^n$ of X via*

$$\dim_{\mathbb{R}} \mathcal{A}(X) = \min\{ 2 \dim_{\mathbb{R}} \text{Trop}(X) + 2 \dim_{\mathbb{R}} T - \dim_{\mathbb{R}} S \mid$$

$$T \subseteq S \subseteq \mathbb{R}^n \text{ rational linear subspaces with } S + (T + \text{Trop}(X)) = T + \text{Trop}(X) \},$$

where a subspace of \mathbb{R}^n is called rational if it is spanned by vectors in \mathbb{Q}^n . Similar to Theorem 3, we have the equivalent formula

$$\dim_{\mathbb{R}} \mathcal{A}(X) = \min\{ 2 \dim_{\mathbb{R}}(S + \text{Trop}(X)) - \dim_{\mathbb{R}} S \mid S \subseteq \mathbb{R}^n \text{ rational linear subspace} \}.$$

Acknowledgments. JD and CHY were partially and fully, respectively, supported by NWO Vici grant entitled *Stabilisation in Algebra and Geometry*. JD thanks Mounir Nisse and Frank Sottile for sharing their work on amoebas at the Mittag-Leffler semester on *Tropical Geometry, Amoebas, and Polyhedra*. All authors thank the Institut Mittag-Leffler for inspiring working conditions. JR thanks Jürgen Hausen for helpful discussions.

2. IN SEARCH OF A POSITIVE-DIMENSIONAL TORUS

Throughout this section we fix an irreducible, closed subvariety $X \subseteq (\mathbb{C}^*)^n$. If $\dim_{\mathbb{R}} \mathcal{A}(X) < 2 \dim_{\mathbb{C}} X$, then we will find a one-dimensional torus $T \subseteq (\mathbb{C}^*)^n$ such that $T \cap (\mathbb{R}^*)^n$ preserves the Zariski-closure $|\overline{X}|$ and $\dim_{\mathbb{R}}(\mathcal{A}(X) + \mathcal{A}(T)) = \dim_{\mathbb{R}} \mathcal{A}(X)$.

Preliminaries. We write $S^1 \subseteq \mathbb{C}^*$ for the unit circle. Recall that this is a real form of the algebraic group \mathbb{C}^* : indeed, tensoring the coordinate ring $\mathbb{R}[c, s]/(c^2 + s^2 - 1)$ of S^1 with \mathbb{C} yields the coordinate ring $\mathbb{C}[c, s]/((c + is)(c - is) - 1)$, which we recognise als the coordinate ring of an algebraic torus with standard coordinate $c + is$; moreover, the the inverse morphism $S^1 \rightarrow S^1, (c, s) \mapsto (c, -s)$ complexifies to the inverse morphism $\mathbb{C}^* \rightarrow \mathbb{C}^*, (c + is) \mapsto 1/(c + is) = (c - is)$; and similarly for the multiplication morphism $S^1 \times S^1 \rightarrow S^1$. Both S^1 and the other real form of \mathbb{C}^* , the real-algebraic group \mathbb{R}^* , will play fundamental roles in our proof.

We write $(S^1)^n \subseteq (\mathbb{C}^*)^n$, where the former is a real form of the latter algebraic group. For $p \in (\mathbb{C}^*)^n$ and Q any subset of \mathbb{C}^n we write pQ for the for the result of coordinate-wise multiplication of p with each element of Q . Writing 1 for the unit element in $(\mathbb{C}^*)^n$ and T_{\bullet} for (real or complex) tangent spaces, we have

$$T_1(\mathbb{C}^*)^n = \mathbb{C}^n = \mathbb{R}^n \oplus_{\mathbb{R}} i\mathbb{R}^n = T_1\mathbb{R}^n \oplus_{\mathbb{R}} T_1(S^1)^n.$$

Component-wise multiplication by $p \in (\mathbb{C}^*)^n$ yields

$$T_p(\mathbb{C}^*)^n = p\mathbb{R}^n \oplus_{\mathbb{R}} ip\mathbb{R}^n = T_pp\mathbb{R}^n \oplus_{\mathbb{R}} T_pp(S^1)^n.$$

Note that $p^{-1}T_pp(S^1)^n$ is naturally identified with (the same) $i\mathbb{R}^n$ for all $p \in (\mathbb{C}^*)^n$, and $p^{-1}T_pp\mathbb{R}^n$ is identified with (the same) \mathbb{R}^n for all p .

Rather than directly with the amoeba of X , we will work with the algebraic amoeba $|X|$, the image of X under the semi-algebraic map

$$\text{abs} : (\mathbb{C}^*)^n \rightarrow \mathbb{R}_{>0}^n, (z_1, \dots, z_n) \mapsto (|z_1|, \dots, |z_n|).$$

The reason for this is that $|X|$ is, by real quantifier elimination, a semi-algebraic set, hence analysable with methods from real algebraic geometry. The following is immediate.

Lemma 8. *At $p \in (\mathbb{C}^*)^n$, the derivative $d_p \text{Log}$ (respectively, $d_p \text{abs}$) sends the real vector space $T_pp(S^1)^n$ to zero and an element pv with $v \in \mathbb{R}^n$ to v (respectively, to $|p|v$). \square*

Subvarieties of real tori. We prove an auxiliary result on subvarieties of real tori. We will use the term *real-Zariski* to refer to the real Zariski topology on a real algebraic variety or, more generally, on a semi-algebraic set. We write Z_{ns} for the nonsingular locus of a real algebraic variety.

Lemma 9. *Let Z be a real-Zariski-closed subset of $(S^1)^n \subseteq (\mathbb{C}^*)^n$. Then the real subspace $\sum_{p \in Z_{\text{ns}}} p^{-1}T_pZ \subseteq i\mathbb{R}^n$ is spanned by its intersection with $i\mathbb{Q}^n$.*

Proof. That subspace is additive under union of irreducible components, so we may assume that Z is irreducible. Let $Z_{\mathbb{C}} \subseteq (\mathbb{C}^*)^n$ be the complexification of Z , an irreducible algebraic variety. After multiplying with p^{-1} for any fixed $p \in Z_{\mathbb{C}}$ we may assume that $1 \in Z_{\mathbb{C}}$. By [Bor91, Proposition 2.2] there exist a natural number m and exponents $e_1, \dots, e_m \in \{\pm 1\}$ such that the image T of the multiplication map

$$\mu_Z : Z_{\mathbb{C}}^m \rightarrow (\mathbb{C}^*)^n, \mathbf{z} = (z_1, \dots, z_m) \mapsto z_1^{e_1} \cdots z_m^{e_m}$$

is a closed, connected algebraic subgroup T of $(\mathbb{C}^*)^n$, i.e., a sub-torus. Since Z_{ns} is Zariski-dense in $Z_{\mathbb{C}}$, there exists a point $\mathbf{z} = (z_1, \dots, z_m) \in Z_{\text{ns}}^m$ (no complexification!) such that the complex-linear map $d_{\mathbf{z}}\mu_Z : T_{\mathbf{z}}Z_{\mathbb{C}}^m \rightarrow T_{\mu(\mathbf{z})}T$ is surjective. Now μ_Z is the restriction to $Z_{\mathbb{C}}^m$ of the multiplication map $\mu : ((\mathbb{C}^*)^n)^m \rightarrow (\mathbb{C}^*)^n$ with the same definition. We have

$$\mu = L_{\mu(\mathbf{z})} \circ \mu \circ (L_{z_1^{-1}} \times \cdots \times L_{z_m^{-1}}),$$

where L_x is left multiplication with $x \in (\mathbb{C}^*)^n$, and accordingly,

$$d_{\mathbf{z}}\mu = d_1 L_{\mu(\mathbf{z})} \circ d_{(1, \dots, 1)}\mu \circ (d_{z_1} L_{z_1^{-1}} \times \cdots \times d_{z_m} L_{z_m^{-1}}).$$

Using that the derivative of multiplication is addition and the derivative of inverse is negation, we find

$$d_{\mu(\mathbf{z})} L_{\mu(\mathbf{z})^{-1}} \circ d_{\mathbf{z}}\mu_Z : T_{\mathbf{z}}Z_{\mathbb{C}}^m \rightarrow T_1T, (v_1, \dots, v_m) \mapsto e_1 z_1^{-1} v_1 + \cdots + e_m z_m^{-1} v_m;$$

and by the choice of \mathbf{z} this map is surjective. For each j we have $T_{z_j}Z_{\mathbb{C}} = T_{z_j}Z \oplus_{\mathbb{R}} iT_{z_j}Z$ and the complex-linear map $d_{\mu(\mathbf{z})} L_{\mu(\mathbf{z})^{-1}} \circ d_{\mathbf{z}}\mu_Z$ sends the real direct sum $\bigoplus_j T_{z_j}Z$ surjectively onto $T_1(T \cap (S^1)^n) = (T_1T) \cap (i\mathbb{R})^n =: Q$. Since T is an algebraic torus, Q is spanned by its intersection with $i\mathbb{Q}^n$, and the space in the lemma contains Q . Moreover, for all $z \in Z$ we have $z^{-1}T_{z_j}Z \subseteq Q$, so that the space in the lemma is in fact equal to Q . \square

A real torus action. We return to our irreducible variety $X \subseteq (\mathbb{C}^*)^n$. By standard results in real algebraic geometry, X is also irreducible when regarded as a real-algebraic variety of dimension $2 \dim_{\mathbb{C}} X$. Consequently, since $|X|$ is the image of X under a semialgebraic map, also $|X|$ is irreducible with respect to the Zariski topology induced from $(\mathbb{R}^*)^n$.

Proof of Proposition 5. For $q \in |X|$ write $Z_q := q^{-1}X \cap (S^1)^n$, a real Zariski-closed subset of $(S^1)^n$ such that $qZ_q = \text{abs}^{-1}(q) \cap X$. There is a subset U of $|X|$, open and dense in $|X|$ in the real Zariski-topology, such that the real algebraic variety Z_q has dimension equal to $c := 2 \dim_{\mathbb{C}} X - \dim_{\mathbb{R}} |X| = 2 \dim_{\mathbb{C}} X - \dim_{\mathbb{R}} \mathcal{A}(X)$. For each $q \in U$, define

$$Q_q := \sum_{p \in (Z_q)_{\text{ns}}} p^{-1}T_p Z_q \subseteq i\mathbb{R}^n,$$

which is a real vector space of dimension at least c , spanned by $Q_p \cap i\mathbb{Q}$ by Lemma 9.

Fix $q \in U$. For each $p \in Z_q$ we have $qp \in X$ and $qp(p^{-1}T_p Z_q) \subseteq T_{qp}X$ and hence, since X is a complex algebraic variety, also $(qp)(ip^{-1}T_p Z_q) \subseteq T_{qp}X$. The space on the left is contained in $qp\mathbb{R}^n$, and hence, by Lemma 8, $d_{qp} \text{abs}$ maps it onto $|qp|(ip^{-1}T_p Z_q) = q(ip^{-1}T_p Z_q)$. We conclude that the latter space is contained in $T_q U$ for each $p \in Z_q$. Therefore,

$$T_q U \supseteq q \sum_{p \in (Z_q)_{\text{ns}}} ip^{-1}T_p Z_q = q(iQ_q);$$

here $iQ_q \subseteq \mathbb{R}^n$ is spanned by its intersection with \mathbb{Q}^n . Now for each vector space $R \subseteq \mathbb{R}^n$ of dimension at least c and spanned by its intersection with \mathbb{Q}^n , the set

$$V_R := \{q \in U \mid T_q U \supseteq qR\}$$

is a real-Zariski-closed subset of U . There are only countably many such R , and the above discussion shows that the closed sets V_R cover the semialgebraic set U .

But then one of them must have dimension equal to that of U , and in fact, since U is irreducible, be equal to U . We conclude that there exists a real vector space $R \subseteq \mathbb{R}^n$, of dimension at least c and spanned by $R \cap \mathbb{Q}^n$, such that $qR \subseteq T_q U$ for all $q \in U$. But then $qR \subseteq T_q \overline{X}$ for all q in the real algebraic variety \overline{X} . Since R is spanned by its intersection with \mathbb{Q}^n , there exists a real-algebraic torus $R_{\mathbb{R}} \subseteq (\mathbb{R}^*)^n$ with Lie algebra R . The sub-bundle of the tangent bundle of $(\mathbb{R}^*)^n$ that arises by differentiating the action of $R_{\mathbb{R}}$ on $(\mathbb{R}^*)^n$ is tangent to \overline{X} . This implies that \overline{X} is $R_{\mathbb{R}}$ -stable. \square

3. PROOFS OF THE MAIN RESULTS

We begin with a lemma that was already used in the introduction (Example 4).

Lemma 10. *Let $X \subseteq (\mathbb{C}^*)^n$ be a closed, irreducible subvariety and $S \subseteq (\mathbb{C}^*)^n$ a subtorus. Then there exists a subtorus $T \subseteq S$ with $\dim_{\mathbb{C}} T = \dim_{\mathbb{C}} \overline{SX} - \dim_{\mathbb{C}} X$ such that $\overline{TX} = \overline{SX}$.*

Proof. We prove the statement by induction on $k = \dim_{\mathbb{C}} \overline{SX} - \dim_{\mathbb{C}} X$. If $k = 0$, then $\overline{SX} = X$ and $T = \{1\}$ will do. If $k > 0$, choose a one-dimensional subtorus $R \subset S$ such that $\dim_{\mathbb{C}} \overline{RX} > \dim_{\mathbb{C}} X$. Such R exists since otherwise X would be invariant under all such R and hence under S . Then the statement follows from the induction assumption applied to $X' = \overline{RX}$ and a torus S' such that $S = R \times S'$. \square

We now use Proposition 5 to establish our dimension formula for the (ordinary or algebraic) amoeba.

Proof of Theorem 3. Let $X \subseteq (\mathbb{C}^*)^n$ be Zariski-closed and irreducible. Since we have already proved the inequality \leq of the theorem, it suffices to establish the existence of subtori $T \subseteq S$ of $(\mathbb{C}^*)^n$ such that $\overline{T \cdot X}$ is S -stable and $\dim_{\mathbb{R}} \mathcal{A}(X) = 2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S$. We proceed by induction on n . For $n = 0$ we have $X = (\mathbb{C}^*)^0$ and we can take $S = T = \{1\}$. So we assume that $n > 0$ and that the statement holds for ambient tori of dimension $n - 1$.

If $\dim_{\mathbb{R}} \mathcal{A}(X) = 2 \dim_{\mathbb{C}} X$, then we may take $T = S = \{1\}$ and we are done. So we may assume that $\dim_{\mathbb{R}} \mathcal{A}(X) < 2 \dim_{\mathbb{C}} X$. Then, by Proposition 5, there exists a one-dimensional, real-algebraic torus $R_{\mathbb{R}} \subseteq (\mathbb{R}^*)^n$ which stabilises the Zariski-closure \overline{X} of the algebraic amoeba. Let $R \subseteq (\mathbb{C}^*)^n$ be the complexification of $R_{\mathbb{R}}$. Then we find an open subset $U \subseteq \mathcal{A}(X)$ whose complement has positive codimension such that U is a smooth manifold with $\mathcal{A}(R) \subseteq T_u U$ for each $u \in U$ (use Lemma 8 for the tangent vectors coming from the action of $R_{\mathbb{R}}$). We find that the fibres of the map $U \rightarrow \mathbb{R}^n / \mathcal{A}(R)$ have dimension 1, and this implies that

$$\dim_{\mathbb{R}} \overline{\mathcal{A}(R) + \mathcal{A}(X)} / \mathcal{A}(R) = -1 + \dim_{\mathbb{R}} \mathcal{A}(X).$$

Define

$$\tilde{X} := \overline{R \cdot X} / R \subseteq (\mathbb{C}^*)^n / R \cong (\mathbb{C}^*)^{n-1}.$$

Then the previous equation implies that the amoeba

$$\mathcal{A}(\tilde{X}) = \overline{\mathcal{A}(R) + \mathcal{A}(X)} / \mathcal{A}(R)$$

has real dimension equal to $-1 + \dim_{\mathbb{R}} \mathcal{A}(X)$.

By the induction hypothesis, there exist subtori $\tilde{T} \subseteq \tilde{S}$ of $(\mathbb{C}^*)^n / R$ such that $\overline{\tilde{T} \cdot \tilde{X}}$ is \tilde{S} -stable and

$$\dim_{\mathbb{R}} \mathcal{A}(\tilde{X}) = 2 \dim_{\mathbb{C}} \tilde{X} + 2 \dim_{\mathbb{C}} \tilde{T} - \dim_{\mathbb{C}} \tilde{S}.$$

We distinguish two cases. First, assume that X is not stable under R , so that $\dim_{\mathbb{C}} \tilde{X} = \dim_{\mathbb{C}} X$. Then let T, S be the pre-images in $(\mathbb{C}^*)^n$ of $\tilde{T}, \tilde{S} \subseteq (\mathbb{C}^*)^n / R$, respectively. Then $T \subseteq S$ are subtori such that $\overline{T \cdot X}$ is S -stable, and we find

$$\begin{aligned} \dim_{\mathbb{R}} \mathcal{A}(X) &= 1 + \dim_{\mathbb{R}} \mathcal{A}(\tilde{X}) \\ &= 1 + 2 \dim_{\mathbb{C}} \tilde{X} + 2 \dim_{\mathbb{C}} \tilde{T} - \dim_{\mathbb{C}} \tilde{S} \\ &= 1 + 2 \dim_{\mathbb{C}} X + 2(-1 + \dim_{\mathbb{C}} T) - (-1 + \dim_{\mathbb{C}} S) \\ &= 2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S. \end{aligned}$$

Second, assume that X is stable under R . As before, let S be the pre-image of \tilde{S} in $(\mathbb{C}^*)^n$, but now let T be any torus in $(\mathbb{C}^*)^n$ of complex dimension *equal* to $\dim_{\mathbb{C}} \tilde{T}$ that projects surjectively onto T . Using that X is R -stable and $\overline{T \cdot X}$ is \tilde{S} -stable, we find that $\overline{T \cdot X}$ is S -stable. Furthermore,

$$\begin{aligned} \dim_{\mathbb{R}} \mathcal{A}(X) &= 1 + \dim_{\mathbb{R}} \mathcal{A}(\tilde{X}) \\ &= 1 + 2 \dim_{\mathbb{C}} \tilde{X} + 2 \dim_{\mathbb{C}} \tilde{T} - \dim_{\mathbb{C}} \tilde{S} \\ &= 1 + 2(-1 + \dim_{\mathbb{C}} X) + 2 \dim_{\mathbb{C}} T - (-1 + \dim_{\mathbb{C}} S) \\ &= 2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S, \end{aligned}$$

as desired.

For the second formula, let T, S be subtori as in the first formula. We then have

$$2 \dim_{\mathbb{C}} \overline{S \tilde{X}} - \dim_{\mathbb{C}} S = 2 \dim_{\mathbb{C}} \overline{T \tilde{X}} - \dim_{\mathbb{C}} S \leq 2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S,$$

so the second formula is a lower bound to the first formula. Conversely, if S is any subtorus, then by Lemma 10 there exists a subtorus $T \subseteq S$ such that $\overline{T X} = \overline{S X}$ and $\dim_{\mathbb{C}} T = \dim_{\mathbb{C}} \overline{S X} - \dim_{\mathbb{C}} X$, and we find

$$2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S = 2 \dim_{\mathbb{C}} \overline{S X} - \dim_{\mathbb{C}} S,$$

hence the first formula is a lower bound to the second formula. \square

Example 11. We give an alternative proof of [NS18, Theorem 4.5], which says that if $\dim_{\mathbb{R}} \mathcal{A}(X) = \dim_{\mathbb{C}} X$, then X is a single orbit under a subtorus of $(\mathbb{C}^*)^n$. Take a subtorus $S \subseteq (\mathbb{C}^*)^n$ that achieves the minimum in the second formula of Theorem 3. Since we always have $\dim_{\mathbb{C}} \overline{S \cdot X} \geq \dim_{\mathbb{C}} S, \dim_{\mathbb{C}} X$, from our choice of S we have

$$\dim_{\mathbb{C}} X = 2 \dim_{\mathbb{C}} \overline{S \cdot X} - \dim_{\mathbb{C}} S \geq \dim_{\mathbb{C}} X.$$

Hence $\dim_{\mathbb{C}} S = \dim_{\mathbb{C}} X = \dim_{\mathbb{C}} \overline{S \cdot X}$. But $\overline{S \cdot X}$ is irreducible and contains both X and an orbit of S , so X must be equal to such an orbit. \clubsuit

Near Torus Actions. We start by reviewing Nisse-Sottile's notion of near torus actions [NS18, Definition 4.1].

Definition 12. Let $X \subseteq (\mathbb{C}^*)^n$ be an irreducible closed subvariety and $S \subseteq (\mathbb{C}^*)^n$. We set $Y := (SX)/S$. Then S has a near action on X if

$$2 \dim_{\mathbb{C}} X > \dim_{\mathbb{C}} S + 2 \dim_{\mathbb{C}} Y \quad \text{and} \quad n > \dim_{\mathbb{C}} S + 2 \dim_{\mathbb{C}} Y.$$

We now show that, as conjectured in [NS18], unexpected amoeba dimension is equivalent to a near torus action. The implication \Leftarrow is see [NS18, Theorem 4.3].

Proof of Corollary 6. For any subtorus $S \subseteq (\mathbb{C}^*)^n$, setting $Y := \overline{SX}/S$ we have $\dim_{\mathbb{C}}(Y) = \dim_{\mathbb{C}}(SX) - \dim_{\mathbb{C}}(S)$. Note that $2 \dim_{\mathbb{C}}(SX) - \dim_{\mathbb{C}}(S) < 2 \dim_{\mathbb{C}}(X)$ and $2 \dim_{\mathbb{C}}(SX) - \dim_{\mathbb{C}}(S) < n$ imply $S \neq \{1\}$ and $S \neq (\mathbb{C}^*)^n$, respectively. Hence the statement follows directly from the second formula of Theorem 3. In particular, in case of a dimension drop, a torus S providing the minimum in this formula has a near action on X . \square

Proof of Corollary 7. We start by presenting a well-known fact in tropical geometry.

Lemma 13. *Let $X \subseteq (\mathbb{C}^*)^n$ be an irreducible (in particular, reduced) closed subvariety and denote by $\text{Trop}(X) \subseteq \mathbb{R}^n$ its tropicalisation. Let $S \subseteq (\mathbb{C}^*)^n$ be a subtorus and $S = \text{Trop}(S) = \mathcal{A}(S) \subseteq \mathbb{R}^n$ the associated (rational) linear subspace. Then $SX = X$ if and only if $S + \text{Trop}(X) = \text{Trop}(X)$.*

Proof. By basic tropical geometry, $\text{Trop}(\overline{SX}) = S + \text{Trop}(X)$. Hence $SX = X$ implies $S + \text{Trop}(X) = \text{Trop}(X)$. Let us assume $S + \text{Trop}(X) = \text{Trop}(X)$ now. Note that for irreducible varieties $Y \subseteq (\mathbb{C}^*)^n$, we have $\dim_{\mathbb{C}} Y = \dim_{\mathbb{R}} \text{Trop}(Y)$, see [BG84, Theorem A]. Since both X and \overline{SX} are irreducible, it follows that $\dim_{\mathbb{C}} \overline{SX} = \dim_{\mathbb{C}} X$. Since $X \subseteq \overline{SX}$, this implies $X = \overline{SX} = SX$. \square

Proof of Corollary 7. By Lemma 13, the pairs of subtori $T \subseteq S \subseteq (\mathbb{C}^*)^n$ such that $S(\overline{TX}) = \overline{TX}$ are in bijection to the pairs of rational linear subspaces $T \subseteq S \subseteq \mathbb{R}^n$ such that $S + (T + \text{Trop}(X)) = T + \text{Trop}(X)$, via $T = \text{Trop}(T)$, $S = \text{Trop}(S)$. Using the relation $\dim_{\mathbb{C}} Y = \dim_{\mathbb{R}} \text{Trop}(Y)$ again, we have

$$2 \dim_{\mathbb{C}} X + 2 \dim_{\mathbb{C}} T - \dim_{\mathbb{C}} S = 2 \dim_{\mathbb{R}} \text{Trop}(X) + 2 \dim_{\mathbb{R}} T - \dim_{\mathbb{R}} S.$$

Hence the two minima agree. The second formula follows similarly as in the proof of Theorem 3. \square

We conclude this paper with a question on computability.

Question 14. *Does there exist an algorithm that, on input a balanced, pure-dimensional, rational polyhedral complex $\Sigma \subseteq \mathbb{R}^n$ which is connected in codimension 1, computes the expression*

$$\min\{2 \dim_{\mathbb{R}}(S + \Sigma) - \dim_{\mathbb{R}} S \mid S \subseteq \mathbb{R}^n \text{ rational subspace}\}$$

from Corollary 7?

The first term is the maximum, over all maximal cones C of Σ , of $\dim_{\mathbb{R}} \Sigma + \dim_{\mathbb{R}} S - \dim_{\mathbb{R}}((C)_{\mathbb{R}} \cap S)$, and hence it is minimised by an S have certain incidences with given linear subspaces of \mathbb{R}^n . If the rationality assumption is dropped, then real quantifier elimination answers the question in the affirmative. However, similar incidence problems often have real but no rational solutions. For instance, a classical result in enumerative geometry says that the number of two-dimensional subspaces in \mathbb{R}^4 (lines in projective three-space) that nontrivially intersect 4 given two-dimensional subspaces in general position is either zero (in which case there are two complex conjugate solutions) or two. In the latter case, even if the four given spaces are rational, the two solutions will typically not be. We do not know whether the existence of rational solutions for such incidence problems is decidable in general, nor whether the additional conditions on Σ force that real solutions imply rational solutions. On the other hand, if X is a variety given by equations with coefficients in, say, some number field, then of course, by real quantifier elimination, there does exist an algorithm for computing $\dim_{\mathbb{R}} \mathcal{A}(X) = \dim_{\mathbb{R}} |X|$.

REFERENCES

- [BG84] Robert Bieri and John R.J. Groves. The geometry of the set of characters induced by valuations. *J. reine angew. Math.*, 347:168–195, 1984.
- [Bor91] Armand Borel. *Linear algebraic groups. 2nd enlarged ed.* New York etc.: Springer-Verlag, 2nd enlarged ed. edition, 1991.
- [NS18] Mounir Nisse and Frank Sottile. Describing amoebas. 2018. Preprint, [arXiv:1805.00273](https://arxiv.org/abs/1805.00273).

UNIVERSITÄT BERN, MATHEMATISCHES INSTITUT, SIDLERSTRASSE 5, 3012 BERN, AND EINDHOVEN UNIVERSITY OF TECHNOLOGY

Email address: `jan.draisma@math.unibe.ch`

UNIVERSITÄT TÜBINGEN, FACHBEREICH MATHEMATIK, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

Email address: `johannes.rau@math.uni-tuebingen.de`

UNIVERSITÄT BERN, MATHEMATISCHES INSTITUT, ALPENEGGSTRASSE 22, 3012 BERN

Email address: `chi.yuen@math.unibe.ch`